

# ACTELIC<sup>®</sup> 5E INSECTICIDE

## LOW ODOR FORMULATION

**FOR CONTROL OF INSECTS ON STORED CORN (FIELD AND POP) AND GRAIN SORGHUM  
(Bulk or Bagged Field Corn, Popcorn, and Grain Sorghum Seed)**

**ACTIVE INGREDIENT:**

Pirimiphos-methyl  
0-(2-(diethylamino)-6-methyl-4-pyrimidinyl)0,0-dimethyl phosphorothioate .....57%

**OTHER INGREDIENTS:**.....43%

**TOTAL**.....100%

Actellic 5E Insecticide contains 5 pounds active ingredient per gallon.

### KEEP OUT OF REACH OF CHILDREN

## DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

#### FIRST AID

<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.  
**HOTLINE NUMBER:** In case of emergency call toll free 1-877-424-7452. **NOTE TO PHYSICIAN:** This product contains an organophosphate. Pirimiphos-methyl is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Give 2 mg. of atropine intramuscularly or subcutaneously and repeat if symptoms of poisoning reappear. Pralidoxime (2 PAM) may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe poisoning, use antidote immediately after establishing an open airway and respiration. Actellic 5E Insecticide contains petroleum distillate. Vomiting may cause aspiration pneumonia.

See side panel for additional precautionary statements.

EPA Reg. No. 1381-170  
**Distributed By:**  
**Winfield Solutions, LLC**  
**P.O. Box 64589**  
**St. Paul, MN 55164-0589**

EPA Est No. \_\_\_\_\_  
**Net Contents** \_\_\_\_\_

1/1202/4



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER:** Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes. Wear protective eyewear (goggles or face shield) when handling. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are barrier laminate or viton. If you want more options, follow the instructions for category G on an EPA chemical-resistant category selection chart.

Mixers, loaders, and other handlers supporting admixture seed and/or grain treatments must wear:

Long sleeve shirt and long pants

Shoes plus socks

Chemical-resistant gloves

Protective eyewear

See engineering controls for additional requirements.

All other mixers, loaders, applicators and other handlers must wear:

Coveralls over long sleeve shirt and long pants

Chemical-resistant footwear plus socks

Chemical-resistant gloves

Protective eyewear

In addition, mixers and loaders must wear:

Chemical-resistant apron

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENGINEERING CONTROLS**

**For products marketed in containers greater than 64 fluid oz. in size:**

For all seed and/or grain treatments, handlers must use a closed system/loading system designed by the manufacturer to enclose the pesticide in a manner that prevents it from contacting handlers. This product is formulated in a container designed for closed mixing and loading. In addition:

- handlers must wear the PPE specified for handlers supporting admixture seed and/or grain treatment.
- handlers must have available to them in case of accident or spill: coveralls and chemical resistant footwear.
- handlers must wear protective eyewear.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**STORAGE:** Store in a cool place and protect from freezing. If accidentally frozen, reconstitute by gently warming to room temperature and thoroughly mixing contents. Keep container closed when not in use.

**PESTICIDE DISPOSAL:** Pesticides are acutely toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Use label language appropriate for container size and type.

**Nonrefillable containers.** Do not reuse or refill this container. Clean container promptly after emptying.

**Nonrefillable container equal to or less than 5 gallons.** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or, by other procedures approved by state and local authorities.

**Nonrefillable container greater than 5 gallons.** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or, by other procedures approved by state and local authorities.

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or, by other procedures approved by state and local authorities.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call  
CHEMTREC 1-800-424-9300.**

### RESISTANCE MANAGEMENT RECOMMENDATIONS

Actellic 5E Insecticide contains a Group 1 Insecticide. Insect/mite biotypes with acquired resistance to Group 1 insecticides/acaricides may eventually dominate the insect/mite population if Group 1 insecticides/acaricides are used repeatedly in the same site or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Actellic 5E Insecticide or other Group 1 insecticides/acaricides.

To delay insecticide resistance consider:

- Avoiding the consecutive use of Actellic 5E Insecticide or Group 1 insecticides that have a similar target site of action, on the same insect/mite species.
- Using tank-mixtures or premixes with insecticides/acaricides from different target site of action Groups as long as the involved products are all registered for the same use and have different sites of action,
- Basing insecticide/acaricide use on a comprehensive IPM program.
- Monitoring treated insect/mite populations for loss of efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or Winfield Solutions, LLC representative for herbicide resistance management and/or integrated pest management recommendations for specific site and resistant pest problems.

### APPLICATION DIRECTIONS

Do not make more than one application per crop. Only treat corn or grain sorghum which has not previously been treated with any pirimiphos-methyl containing product. Treatment of seed is only allowed off-farm (for example, in a facility where seed is bagged).

### ADMIXTURE OR TOP DRESSING

The best practice to protect stored corn (field and pop) and grain sorghum from insect infestation is to use an admixture treatment of Actellic 5E Insecticide. However, as it may not always be possible to move grain, a surface treatment of Actellic 5E Insecticide will help prevent the establishment of migrating insects. The top dressing application will not substitute for a treatment of the entire amount of stored corn (field and pop) or grain sorghum. But within your current stored grain management system, it will help protect the surface from invasion by insects.

### ADMIXTURE TREATMENT

For products marketed in containers of 64 fl. oz. or less, use for treatment is prohibited.

Apply spray solution uniformly with suitable application equipment as grain enters storage or export containers. Select nozzle size and operating pressure to adjust the application rate with flow of grain. Prepare desired dilution of Actellic 5E Insecticide in spray tank and mix well. Make sure spray solution stays mixed, especially in large holding tanks. Apply the spray solution as a coarse, uniform low pressure spray in order to minimize drift. Shield nozzle if necessary. Apply the specified amount of Actellic 5E Insecticide in 5 gallons of water or 3-5 gallons of at least twice refined soybean oil to each 30 tons (60,000 pounds) of grain according to the following rates.

	Fluid Ounces	ppm Concentration
Shelled Corn (field corn, popcorn)	9.2-12.3	6-8
Grain Sorghum	9.2-12.3	6-8
	No. of Bushels Approximately Equal to 30 Tons of Grain	
Shelled Corn (field corn, popcorn)	1,071	
Grain Sorghum	1,071	

### TOP DRESSING TREATMENT

To each 1,000 square feet of grain surface, apply 3 fluid ounces of Actellic 5E Insecticide diluted in 2 gallons of water. The treatment should be split as follows: Spray half of the mixture evenly over the surface and rake into the grain to a depth of 4 inches. Apply the remaining half of the mixture (1 gallon) to the raked surface.

Before making a surface application of Actellic 5E Insecticide, it will be necessary to remove any webbing and break up crusting that exists on the grain surface. If high numbers of Indian Meal Moth worms (2-3 per sq ft) exist, it will be best to remove the webbing, fumigate, and then apply Actellic 5E Insecticide as a surface application. Mix the following amount of Actellic 5E Insecticide with 2 gallons of water for a total mixture to treat 1,000 sq ft of grain surface:

	Fluid Ounces	ppm Concentration
Shelled Corn (field corn, popcorn) and Sorghum	3.0	8

### INSECTS CONTROLLED WITH ACTELLIC 5E INSECTICIDE

COLEOPTERA		LEPIDOPTERA
Cigarette beetle	Maize weevil	Almond moth
Confused flour beetle	Merchant grain beetle	Angoumois grain moth
Corn sap beetle	Red flour beetle	Indian meal moth
Flat grain beetle	Rice weevil	
Granary weevil	Sawtoothed beetle	
Hairy fungus beetle		

- Grains treated with Actellic 5E Insecticide at rates indicated on this label may be used immediately for any food or feed purpose.
- Do not make more than one application per crop. Only use on treated corn (field and pop) or sorghum which has not previously been treated with any pirimiphos-methyl containing product.
- Calibrate spray equipment and periodically check in order to ensure proper spray rates.
- Do not store diluted Actellic 5E Insecticide in spray tank more than 48 hours before use.
- Avoid severely dusty application sites.
- Seed applications of Actellic 5E Insecticide are restricted to bulk or bagged seed corn (field and pop) and sorghum grain.
- Only to be used as grain enters storage or export containers. Not for use to control soil insects.
- Do not enter treated areas or have contact with treated grain or seed until sprays have dried.

#### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED “AS IS” BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

#### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER’S OR USER’S EXCLUSIVE REMEDY, AND MANUFACTURER’S OR SELLER’S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Actellic® is a registered trademark of a Syngenta Group Company.  
AgriSolutions™ is a trademark of Winfield Solutions, LLC.