Only apply this product to established ornamentals.

non-target plants unless they are listed on this label.

Treatment of Ornamental species not listed on this label:

under all possible growing conditions prior to large scale use.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

©2019 The Andersons, Inc. Maumee, OH 43537

10005667

Net Wt. 50 lbs. (22.68 kg)

SPREADER SETTINGS

LOW RATE | MEDIUM RATE | HIGH RATE | HEAVY RATE

sq. ft.

L 1/4

12

5 1/4 IIIC

23

3.4 lbs/1,000 | 3.6 lbs/1,000 | 4.6 lbs/1,000

sq. ft.

L 3/4

K 1/4

E 1/4

12 1/2

5 1/2 IIIC

26

sq. ft.

M 1/2

L 1/2

14

6 IIIC

29

Do not apply this product directly to bare root of ornamental plants as injury may result. Following transplanting, care must be taken that soil or planting mixes have settled firmly through irrigation, rainfall or packing and that there are no cracks that would allow direct contact of this product to the plant roots or

• Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may

Do not apply around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due

Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
 Do not apply this product on grasses grown for seed.

• Do not apply another preemergence herbicide within 4 weeks of application of this product.

not apply this product directly to plants that are grown for food (e.g., fruit trees or maple trees tapped

possible growing conditions. Users who wish to use this product on ornamental species not listed on this label may determine the suitability for use by treating a small number or ornamental plants at a listed rate. Prior to

n tolerance in field trials. However, this product has not been tested on all cultivars of each species or

treatment of larger areas, treated plants should be observed for any symptoms of herbicidal injury, such as foliar damage, reduced vigor or stand reduction, during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious.

When applied as directed under the conditions described on this label, ornamentals listed on back have

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE

LEGAL RIGHTS:THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

 ${\tt CONTRARY\ TO\ LABEL\ DIRECTIONS.\ Use\ contrary\ to\ label\ directions\ is\ not\ permitted.}$

to insects, nematodes or disease.

• Irrigate immediately after application to remove herbicide particles from the foliage.

• For use on Turf, Landscape Ornamentals, and Nursery Ornamentals • On DG Pro® Carrier

weeks apart and prior to crabgrass emergence.

For season long control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape

For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and areas, established lawns and ornamental turf (including golf course fairways, roughs and tee boxes and sports turf) and container and field grown nursery ornamentals. New York Use Restrictions: In New York State this product may only be used by professional applicators and at DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. no more than 4.6 pounds per 1000 square feet (equal to 0.5 pound active ingredient per acre); and is prohibited from use in Nassau and Suffolk Counties. ACTIVE INGREDIENTS:

CAS No. 97886-45-8 *This product contains 0.125 pounds of active ingredient, dithiopyr, per 50 pound bag. EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS KEEP OUT OF REACH OF CHILDREN

OTHER INGREDIENTS . .

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some

FIRST AID	
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
	ict container or label with you when calling a poison control center or doctor or

going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment

his product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters. e this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forest, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, tification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protection equipment (PPE), restricted-entry intervals. The requirements in this box only apply to uses of this product, that are covered by the Worker Protection Standard. Do no

enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours unless otherwise specified in the directions for use associated with each crop. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: 1. Coveralls over short sleeve shirt and short pants, 2. Chemical-resistant footwear, shoes plus socks, 3. Chemical-resistant gloves made of any waterproof material, and gloves, 4. Chemical-resistant headgear for overhead exposure. When the Restricted-Entry Interval for a crop is 7 days or longer, you must notify workers of the application by warning them orally and by posting warning signs at entrances to treated area. Non-Agricultural Use Requirements The requirements in this box apply to uses of this product that are NOT within the scope of the Worker

Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses. Do not enter or allow others to enter the treate area (except those involved in the watering-in) until the watering-in is completed and the area has dried. DIRECTIONS FOR USE

STORAGE AND DISPOSAL
: is a violation of Federal Law to use this product in any manner inconsistent with its labeli lead the entire label before using this product. Use only according to label instructions.
ARECTIONS FOR USE

STORAGE AND DISPOSAL STORAGE: Store this product only in its original container in a dry, cool, secured storage area. Do not contaminate water, food, or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies.
PESTICIDE DISPOSAL : Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal in accordance with applicable Federal, state or local procedures.
CONTAINER DISPOSAL : Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill by inciparation, or if allowed by State or local authorities by burning. If burned, stay out of smoke

for research purposes. This product is a herbicide that provides control of crabgrass and other annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes and greens. This product will not harm nearby established ornamentals when used according to label directions.

SPILL: In case of spill, sweep up material and dispose of material according to "Pesticide Disposal" directions

TURFGRASS SAFETY is product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established.

mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass

per year. species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product: Warm Season Grasses Bentgrass, creeping* Fescue, tall Bahiagrass Carpetgrass St. Augustinegrass

Bluegrass, Kentucky Ryegrass, perennial Bermudagrass Centipedegrass Zoysiagrass Fescue, fine** Ryegrass, perennial Buffalograss*** Kikuyugrass Zoysiagrass *Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. DO NOT apply this product to Colonial bentgrass (Agrostis tenuis) varieties. **Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf',

and 'Waldina'. Not all varieties of fine fescue have been tested.

***DO NOT use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots. NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties to this product on a limited area, at use rates, prior to initiating RESEEDING, OVERSEEDING, OR SPRIGGING

Reseeding, overseeding, or springing of treated areas with this product should be delayed until 12 weeks from the time of application. Reseeding, overseeding or springing before 12 weeks after application may prevent establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil overseed 8 weeks or later after a single application. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

SPREADER

Andersons Model 2000/ AccuPro 2000; (cone 6)

Andersons 2000 SR/ SR-2000 (cone 6)

Andersons LCO-1000

LESCO 020093 (0.25)

LESCO 705698 (0.25)

(PTO at 450 rpm) Vicon (03 Series)

Scotts R-8, R-8A (cone 6)

Lely Models WTR, WFR, HR, 125-

APPLICATION DIRECTIONS CONTROL OF CRABGRASS Preemergence and Early Postemergence Control

This product provides "preemergence" control of crabgrass (including the large, smooth, and southern species) when applied prior to the emergence of crabgrass from the ground in established lawns and ornamental turfs. It can also provide "early postemergence" control of crabgrass. Postemergence crabgrass control will be product's early postemergence activity is it can give the user a 2-4 week longer period of time to make

is product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall. DO NOT apply more than 4.6 lbs. per 1000 sq. ft. per application, and no more than 13.8 lbs. per 1000 sq. ft. per

It is impossible to evaluate tolerance to this product on all ornamental plant species or varieties or under all

For single applications made in the spring or early summer, this product should be applied at the appropriate rate listed in Table 1 below. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region. Use the higher rates where weed infestations are dense. Also use the higher rates if applications are made early and longer residual control is desired. Subsequent, sequential pre- and/or postemergence applications should be made where longer periods of Where split fertilizer applications are listed, the rates in Table 1 may be split across two application made 6-10

	108-160 lbs./acre 2.4-3.6 lbs./1,000 sq ft 0.27-0.40 lbs. ai/acre			
Barley, Brome, Crabgr. Dallisgrass (seedling),				
Barnyardgrass Chickweed Clover, hop Crabgrass, tropical Crowfootgrass Dichondra, Carolina Lespedeza, common Kikuyugrass Mary's grass Marestail Medic, black	Mulberry weed Mustard Oats, wild Oxalis Buttercup Creeping Woodsorrel Yellow Woodsorrel Pineappleweed Ryegrass, annual Ryegrass, perennial Sandbur	Smutgrass Speedwell, corn Spurge Garden Prostrate Spotted	160-200 lbs./acre 3.6-4.6 lbs./1,000 sq. ft. 0.40-0.50 lbs. ai/acre	
Bluegrass, annual Bittercress Carpetweed Chickweed Geranium Carolina Goosegrass	Henbit Knotweed, prostrate Parsley-piert Pigweed, redroot Purslane, common Rocket, London		200 lbs./acre 4.6 lbs./1,000 sq. ft. 0.50 lbs. ai/acre	

This product can also be applied in the late summer or early fall (late August through November) at the use Table 2: Application Use Rates for Fall Applications (lbs. of product/1,000 square feet)

Use the higher rates where weed infestations are dense. Also use the higher rates if applications are made early 081622

*DO NOT apply more than 4.6 lbs. per application, and no more than 13.8 lbs. per year. Postemergence control is limited; see "Crabgrass Control" and "Precautions" sections above. Tips For Improved Control For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result. Use of split (half-rate) applications spaced 6-8 weeks apart may provide improved weed control.

APPLICATION EQUIPMENT AND INSTRUCTIONS Apply this product with suitable spreaders such as drop or rotary-type designed to apply granular herbicides. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate listed above. Apply this product uniformly over the lawn or ornamental turf area. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform ORNAMENTALS (Landscape, Field Grown and Container Grown) This specialty herbicide provides preemergence control of listed annual grass and broadleaf weeds in areas planted with tolerant ornamental plants listed on this label. It is intended for use on plants grown for aesthetic

purposes in landscaped areas or in container or field grown production nurseries. When applied as directed, the product may be applied as a single application or as a split application. Refer to the following table for product DO NOT apply more than 200 lbs. per acre per application and no more than 3 applications (600 lbs.) per acre In New York State, this product may only be used by Professional Applicators at no more than 200 lbs. per acre (4.6 pounds per 1,000 sq. ft./ 0.5 lbs. active ingredient per acre) per year; and is prohibited from use in Nassau

Pounds of Product Pounds of Product Ounces of Product per Acre per 1,000 square feet per 100 Square Feet 4.6 7.36

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

Suggested Spreader Settings These suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product. 50 lbs treats 20,800 sq. ft. (0.48 acre) at the LOW RATE (2.4 lbs/1,000 sq .ft.)

50 lbs treats 14,700 sq. ft. (0.34 acre) at the MEDIUM RATE (3.4 lbs/1,000 sq. ft.)

50 lbs treats 13,800 sq. ft. (0.32 acre) at the HIGH RATE (3.6 lbs/1,000 sq.f t.)

50 lbs treats 10,800 sq. ft. (0.25 acre) at the HEAVY RATE (4.6 lbs/1,000 sq. ft.)

11 feet

11 feet

7 feet

7 feet

34 feet

30 feet

10

4 1/4 IIIC

SPREADER | COVERAGE |

3 mph

3 mph

3 mph

3 mph

4.5 mph

4.5 mph

GROUND | WIDTH OF | 2.4 lbs/1,000 |

Annual Grasses Barley Barnyardgrass Bluegrass, annual Brome Crabgrass, Iarge Crabgrass, smooth Crabgrass, southern Crowfootgrass Dallisgrass (seedling) Foxtail, green Foxtail, green Gosegrass Kikuyugrass Kikuyugrass Ryegrass (annual & perennial) Sandbur Smutgrass Smutgrass Smutgrass	Broadleaf Weeds Bittergrass Carpetweed Geranium, Carolina Henbit Knotweed prostrate Lespedeza, common Marestail Medic, black Mulberry weed Mustard Oxalis, buttercup	Parsley-piert Pigweed, redroot Pineappleweed Purslane, common Rocket, London Shepherdspurse Speedwell, corn Spurge, garden Spurge, prostrate Spurge, spotted Woodsorrel, creeping Woodsorrel, yellow
---	--	--





Acceptable Application Method Noted by an (X)

Ajuga carpet bugle (Ajuga reptans)

almond, flowering (Prunus gladulosa)

(Thuja occidentalis)

arborvitae, dwarf golden (Thuja orientalis)

ash, autumn purple

(Fraxinus americana)

(Rhododendron spp.)

barberry, Japanese (Berberis thunbergii)

(Berberis thunbergii var

(Gaillardia spp.)

(Vaccinium spp.)

wood, welleri

(Camellia japonica)

(Camellia sasanqua)

ress, Hinoki false

(Hemerocallis spp.)

(Delphinium spp.)

ogwood, American

Pseudotsuga menziesi

(Euonymus japonicus)

(Euonymus fortunei)

an palm, European (Chamaerops humilis)

fan palm, Mexican (Washingtonia robusta)

Manhattan

(Cornus sericea)

nthus (sweet William)

ressocyparis leyland

dwarf pygmy

Professional	Turf ^m Products
	© (C)

F
ATI

				(i or sy triid suspensu)	I	1 '	×		I v I	(Osteospermum muticosum)						4
<	Χ	X	X		weeping .		-		X	pachysandra			×		X	l -
			 	fountain grass, purple (Pennisetum setaceum)	rubrum	X	X	×	×	(Pachysandra terminalist)	ļ	ļ			 /	4
-	X		X	fuchsia (Fuchsia spp.)			Х		×	palm, King (Archontophoenix cuninghamiana)			X		X	
-	X		×	galium (Galium ordoratum)			Х		Х	palm, queen (Syagrus romanzoffianum)		Х	Х	×	X	
(Х	gardenia (Gardenia jasminoides)	mystery radicans	,	X		X X	pampas grass (Cortaderia selloana)			Х		Х	
(X X X	Garlic, variegated society	white gem variegata	X	X	X	X	pansy (Viola x wittrockiana)			Х		Х	
			x	(Thulbaghia violacea) geranium (Pelargonium x hortorum)			×		X	paper flower (Bougainvillea glabra)	Barbara Karst	Х	X	X	Х	
			X X X	gum (Eucalyptus citriodora)			×		Х	peach† (Prunus persica)			Х		Х	
			x	hawthorn (Crataegus spp.)	cockspur white crimson cloud		X		X X	periwinkle, dwarf (Vinca minor)			×		X	
			X X X	(IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	enchantress Jack Evans		X X		X X	petunia (Petunia x hybrida)	picoti		Х	<u> </u>	Х	
			X X	havubhann ladiss	Washington white		X		X	photinia, red tip (Photinia x fraseri)			X	<u> </u>	X	
			×	hawthorn, Indian (Rhaphiolepsis indica)	ballerina		X		X	pieris (Pieris taiwanensis)			X		X	
			X X X	heather, twisted (Erica cinerea)	Mediterranean pink		X		X	pieris, Japanese (Pieris japonica)	mountain fire		X	<u> </u>	X	
\dashv	Х		X	hemlock, Canada (Tsuga canadensis)	blue bird		X		X	pine, Afghan (Pinus eldarica)		X	X	×	X	
\dashv	X		X	hisbiscus (Hibiscus spp.)	blue bird brilliant hula girl		X X X		X X X	pine, Austrian black (pinus nigra)		Х	X	×	X	
	X		X	holly (llex x 'Nellie R. Stevens')	Nellie R.	×	Х	×	х	pine, Japanese black (Pinus Thunbergiia) pine, loblolly			X	 	X	
	X		X	(llex x attenuata)	Stevens fosteri Savannah		X X		X X	(Pinus taeda) pine, longleaf			X	 	X	
	X X		X X	holly, blue (llex x meserveae)	blue bird brilliant		X		X	(Pinus palustris) pine, mugo or Swiss Mt.			X	 	X X	
	X X X		X X X	holly, cassine	hula girl		X		x	(Pinus mugo) pine, Scotch			X			
\neg	Х		Х	(llex cassine) holly, Chinese	burfordii		x		x	(Pinus sylvestris) pine, slash			X	 	X	
\dashv	X		Х	(llex cornuta) holly, Japanese	needlepoint compacta		X		x	(Pinus elliottii) pine, Virginia			X			
\dashv	X		Х	(llex crenata)	hellerie Japanese		×		X X	(Pinys virginiana) pine, white		X	X	X		
	X		Х	hally your on	northern beauty steeds	Х	×	Х	X	(Pinus strobus) pineapple guava†			X		X	
\dashv	X		Х	holly, yaupon (Ilex vomitoria)			Х		X	(Feijoa sellowiana) pittosporum, tobira			^			
\dashv	Х		X	honeysuckle (Lonicera xylosteum) (Lonicera japonica)	Claveyi dwarf halliana		X X		X X	(Pittosporum tobira)	Wheeler's dwarf	Х	Х	×	Х	
\dashv	Х		X	(Lonicera tatarica)	Canadian white Zabelli		X X		X X	plumbago, cape (Plumbago auriculata)	royal cape	Х	Х	Х	Х	
\dashv	Х		Х	hosta (Hosta sieboldii) (Hosta lancifoliai)	albo marginata		X X		××	potentilla (Potentilla fruticosa) (Potentilla nepalensis)	abbotswood		X X		X X	
	Х		Х	ice plant (Carpobrotus edulis)	aibo marginata	×	X	Х	X	privet (Ligustrum x vicaryii)	golden vicary		X	<u> </u>	X	
	Х		Х	ice plant (Drosanthemum floribundum)		×	Х	Х	X	(Ligustrum japonicum)	regal texanum	Х	X X	×	X	
	X X X		X X X	impatiens (Impatiens spp.)			Х		Х	privet, glossy (Ligustrum lucidum)	yellow tipped		X	 	X	
I	^	I	1	1.2.	alore of laters	$\overline{}$, , , , , , , , , , , , , , , , , , ,	i	i , ,	(2.gusti aiii iucidaiii)	-	-	-	+	+	4

uince, Japanese †

(Rhododendron spp)

nododendron, rhodie max-

e, knockout shrub (Rosa spp. hybrid)

semary, bog

smoketree, royal purple

(Cotinus coggygria)

pruce, Black Hills

oruce, Norway

(Liquidambar styraciflua)

camore, American

tea tree, New Zealand

rumpet flower or Carolina

tree fern (tiki fern)

Verbena, St. Paul's

viburnum (Viburnum spp.)

(Taxus cuspidata)

(Verbena peruviana)

nonroe white

marigold (Tagetes patula)

mock orange† (Philadelphus spp)

(Ophiopogon japonicus)

(Lagerstroemia indica)

X

ruce, Colorado blue

Container Field/ Land-Grown scape Grown Nam

A cceptable Application Method Noted by an (X)

Field/Landscapes or Container Grown Ornamentals Noted by an (X)

Acceptable Application Method Noted by an (X)

(Forsythia x 'Meadowlark

(Forsythia x intermedia)

	E
	E
	E