


**GUARANTEED ANALYSIS:** 



### ELEMENTAL **GUARANTEED ANALYSIS:**

TOTAL NITROGEN (N)12.00%	TOTAL NITROGEN (N) 12.00%
5.00% Ammoniacal Nitrogen	5.00% Ammoniàcal Nitrogen
7.00% Urea Nitrogen*	7.00% Urea Nitrogen**
AVAILABLE PHOSPHATE ( $P_2O_5$ )4.00%	TOTAL PHOSPHORUŠ (P)1.75%
SOLUBLE POTASH (K <sub>2</sub> O)	TOTAL POTASSIUM (K)
Magnesium (Mg)3.00%	Magnesium (Mg)
Sulfur (S)	Sulfur (S)
Copper (Cu)0.05%	Copper (Cu)0.05%
Iron (Fe)	Iron (Fe)
Manganese (Mn)0.20%	
Molybdenum (Mo)0.001%	Molybdenum (Mó)0.001%
Zinc (Zn)0.05%	Zinc (Zn) 0.05%
Derived from Polymer-Coated Urea, Monoammonium Phosphate, Ammonium Sulfate, Sulfate of Potash, Sulfate of Potash-Magnesia, Magnesium Carbonate, Magnesium Oxide, Magnesium Sulfate, Iron Oxysulfate, Ferric Oxide, Ferrous Sulfate, Copper Oxide, Copper Sulfate, Manganese Oxide, Manganese Oxysulfate, Manganese Sulfate, Sodium Molybdate, Zinc Oxide and Zinc Sulfate.	Derived from Polymer-Coated Urea, Monoammonium Phosphate, Ammonium Sulfate, Sulfate of Potash, Sulfate of Potash-Magnesia, Magnesium Carbonate, Magnesium Oxide, Magnesium Sulfate, Iron Oxysulfate, Ferric Oxide, Ferrous Sulfate, Copper Oxide, Copper Sulfate, Manganese Oxide, Manganese Oxysulfate, Manganese Sulfate, Sodium Molybdate, Zinc Oxide and Zinc Sulfate.
Manganese (Mn)0.20% Molybdenum (Mo)0.001% Zinc (Zn)0.05% Derived from Polymer-Coated Urea, Monoammonium Phosphate, Ammonium Sulfate, Sulfate of Potash, Sulfate of Potash-Magnesia, Magnesium Carbonate, Magnesium Oxide, Magnesium Sulfate, Iron Oxysulfate, Ferric Oxide, Ferrous Sulfate, Copper Oxide, Copper Sulfate, Manganese Oxide, Manganese Oxysulfate, Manganese	Manganese (Mn)0.2 Molybdenum (Mo)0.00 Zinc (Zn)0.0 Derived from Polymer-Coated Urea, Monoammonium Phosphate, Ammonium Sulf Sulfate of Potash, Sulfate of Potash-Magnesia, Magnesium Carbonate, Magnes Oxide, Magnesium Sulfate, Iron Oxysulfate, Ferric Oxide, Ferrous Sulfate, Cop Oxide, Copper Sulfate, Manganese Oxide, Manganese Oxysulfate, Mangar



**U.S. STANDARD** 

## See reverse side for Application Rates and Application Precautions.



J.R. Simplot Company Professional Products Simplot. P.O. Box 198 · Lathrop. CA 95330 · (208) 336-2110 IN CASE OF EMERGENCY CALL CHEMTREC 1-800-424-9300





J.R. Simplot Company Professional Products



Professional Products P.O. Box 198 • Lathrop, CA 95330 • (208) 336-2110 IN CASE OF EMERGENCY CALL CHEMTREC +1-703-527-3887 • CCN# 12097





APPLICATION RATES: (Call for rates on larger containers.) Use LOW rate for low feeding, sensitive plants or under high soil temperatures. Use MEDIUM rate for medium to moderately heavy feeding plants.

Use HIGH rate only for heavy feeding hardy plants. These application rates are based on the average temperature at the fertilizer location of 70°F. Increase fertilizer application rates by 20% if average monthly temperatures are lower than 60°F. Lower application rates by 20% if average monthly temperatures are greater than 80°F. CONVERSION TABLE

DRY MEASURE			1	APEX <sup>®</sup> S	SPOONS				
Level Measure	Grams	Oz.(Wt.)		Size	Grams	Oz.(Wt.)	Size	Grams	Oz.(Wt.)
1 teaspoon (tsp.)	6.8	0.24		1	11.7	0.41	5	41.8	1.47
1 tablespoon (tbsp.)	22.3	0.79		2	17.8	0.63	6	63.0	2.22
1/4 cup	68.5	2.42		3	24.8	0.87	7	81.9	2.89
1/2 cup	142.7	5.03		4	30.8	1.09	8	98.0	3.46
TOPDRESS CO	NTAINER	: vo	LUN	ME (gal.)	DIAME	TER L	.ow	MEDIUM	HIGH
Plant Nutrient Requir	rements /		1 g	allon	6 incl	nes 3	.5 g	7 g	10.5 g
Uniformly apply (top		uct	2 g	allons	8 incl	nes	8 g	15 g	23 g
			3 g	allons	10 inc	hes	13 g	26 g	39 g
onto the container su	unace using	j trie	5 g	allons	12 inc	hes	20 g	40 g	60 g
amounts listed.			7 g	allons	14 inc	hes	30 g	60 g	90 g
		1	0 g	allons	17 inc	hes	50 g	100 g	150 g
		1	5 g	allons	18 inc	hes	70 g	140 g	210 g

INCORPORATION: Plant Nutrient Requirements / Uniformly mix (incorporate) nursery fertilizer into potting media as follows:

POUNDS PE	ER CUBIC YARD	LOW 3.5	MED 7	HIGH 10.5				
PLANTING BED: Plant Nutrient Requirements (incorporate if possible or use lower rates) as follows:								
POUNDS PI	ER 100 SQ. FT.	LOW 1.5	MED 3	HIGH 4.5				
SURFACE APPLICATION: Uniformly apply around drip line of tree.								
CUPS	PALM SIZE	LOW	MEDIUM	HIGH				
PER	2-3 ft.	1/2	1	2				
TREE	4-6 ft.	1	2	3				
	7-10 ft & up	2	3	4+				

# APPLICATION PRECAUTIONS:

• Trial before use of this product under your local growing conditions, application methods, and desired rates. Avoid application to plants under stress.

If mixed media is not used within one week, leach thoroughly before using.

- · Product left in media for more than one week will lose longevity resulting in reduced release time and wasted controlled release fertilizer.
- · Avoid the use of media processing equipment that could change the integrity of the polymer coating.
- Avoid mounding of fertilizer against base of plant.
  Iron and other plant nutrients can cause staining of cement.
- · Keep away from pools, ponds, and other bodies of water
- · When using potting media with higher cation exchange capacities use lower recommended rates of this formulation.
- · When using supplemental liquid feed reduce the rate of this formulation accordingly.
- Do not incorporate into media prior to steam sterilization.
   This product is not recommended for dibble applications.
- To avoid buildup of soluble salts, occasional leaching may be necessary.
   WARNING: Application of fertilizer materials containing Molybdenum (Mo) may result in forage crops containing levels of Molybdenum (Mo) that are toxic to ruminant animals.

DISCLAIMER OF WARRANTIES J.R. Simplot Company warrants that the composition of this product conforms to the description and is reasonably fit for the purpose as stated on the label only when used in accordance with label directions under normal conditions of use. This warranty is expressly in lieu of all other warranties and representations expressed, implied, or statutory, including warrante so of merchantability and fitness for a particular use. Timing, rate and method of application, weather and crop conditions, and mixtures not specifically recommended on this label or an accompanying written recommendation, are beyond the control of the seller. Buyer assumes all risks of use, storage event of damages resulting from, but not limited to, weather or soil conditions, presence of other materials, or failure to follow label directions, responsibility will be assumed by the buyer or user; and to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions presences of celler's liability. No net is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions scale and no such warranties, guarantees or directions concerning this product, and no such warranties, guarantees or directions concerning this product. Buyer Such S-23-2016 R4-25-2017

U.S.A. Trademark designation:

DUAL\_GHS\_5-23-2016\_R4-25-2017

, SDS and other on APEX products

Simplot®, APEX®, and GAL-XeONE® are registered trademarks of the J.R. Simplot Company. 1218A

J.R. Simplot Company Professional Products

Simplot P.O. Box 198 • Lathrop, CA 95330 • (208) 336-2110 IN CASE OF EMERGENCY CALL CHEMTREC 1-800-424-9300



# 12-1.75-9.96 PALM K

# with GAL-Xe ONE™

APPLICATION RATES: (Call for rates on larger containers.) Use LOW rate for low feeding, sensitive plants or under high soil temperatures. Use MEDIUM rate for medium to moderately heavy feeding plants. Use HIGH rate only for heavy feeding hardy plants.

These application rates are based on the average temperature at the fertilizer location of 21.0°C. Increase fertilizer application rates by 20% if average monthly temperatures are lower than 15.5°C. Lower application rates by 20% if average monthly temperatures are greater than 26.5°C. CONVERSION TABLE

001	 101014	

DRY MEASURE			7	<b>APEX</b> <sup>™</sup>	SPOONS				
Level Measure	Grams	Oz.(Wt.)		Size	Grams	Oz.(Wt.)	Size	Grams	Oz.(Wt.)
1 teaspoon (tsp.)	6.8	0.24		1	11.7	0.41	5	41.8	1.47
1 tablespoon (tbsp.)	22.3	0.79		2	17.8	0.63	6	63.0	2.22
1/4 cup	68.5	2.42		3	24.8	0.87	7	81.9	2.89
1/2 cup	142.7	5.03		4	30.8	1.09	8	98.0	3.46
		DIAMETER (mm) 100 mm 125 mm 150 mm 175 mm 200 mm 250 mm 300 mm		LOW 0.7 g 1.5 g 2.0 g 4.0 g 6.0 g 13.0 g 18.0 g	1 3 4 8 12 26	DIUM .4 g .0 g .0 g .0 g .0 g .0 g .0 g	HIGH 2.1 g 4.5 g 6.0 g 12.0 g 18.0 g 39.0 g 54.0 g		
INCORPORATION: Diant Nutriant Description ante / Uniformity (incorporate) auroau fastilizar inte activita									

**INCORPORATION:** Plant Nutrient Requirements / Uniformly mix (incorporate) nursery fertilizer into potting media as follows:

KILOGRAMS	PER CUBIC METRES	LOW 2	MED 4	HIGH 6				
PLANTING BED: Plant Nutrient Requirements (incorporate if possible or use lower rates) as follows:								
KILOGRAMS I	PER 100 SQ. METRES	LOW 7	MED 14	HIGH 21				
SURFACE APPLICATION: Uniformly apply around drip line of tree.								
GRAMS	PALM SIZE	LOW	MEDIUM	HIGH				
PER	0.5-1 metre	130 g	260 g	520 g				
TREE	1-2 metres	260 g	520 g	780 g				
	2-3 metres & up	520 g	780 g	1,040 g				

### APPLICATION PRECAUTIONS:

- Trial before use of this product under your local growing conditions, application methods, and desired rates. Avoid application to plants under stress.
- If mixed media is not used within one week, leach thoroughly before using
- Product left in media for more than one week will lose longevity resulting in reduced release time and wasted controlled release fertilizer.
- Avoid the use of media processing equipment that could change the integrity of the polymer coating.
- Avoid mounding of fertilizer against base of plant. Iron and other plant nutrients can cause staining of cement.
- Keep away from pools, ponds, and other bodies of water
- · When using potting media with higher cation exchange capacities use lower recommended rates of this formulation.
- When using supplemental liquid feed reduce the rate of this formulation accordingly.
- Do not incorporate into media prior to steam sterilization.
- This product is not recommended for dibble applications.
- Da avoid buildup of soluble salts, occasional leaching may be necessary.
   WARNING: Application of fertilizer materials containing Molybdenum (Mo) may result in forage crops containing levels of Molybdenum (Mo) that are toxic to ruminant animals.

DISCLAIMER OF WARRANTIES J.R. Simplot Company warrants that the composition of this product conforms to the description and is reasonably fit for the purpose as stated on the label only when used in accordance with label directions under normal conditions of use. This warranty is expressly in lieu of all other warranties and representations expressed, implied, or statutory, including warrante is of merchantability and fitness for a particular use. Timing, rate and method of application, weather and crop conditions, and mixtures not specifically recommended on this label or an accompanying written recommendation, are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herwith. Buyer further agrees that, in the event of damages resulting from, but not limited to, weather or soil conditions, presence of other materials, or failure to follow label directions, responsibility will be assumed by the buyer or user; and to accept a replacement of the product as they rate or directions as full discharged seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions the valid or binding upon seller. International Trademark designations. DUAL\_GHS\_5-23-2016\_R4-25-2017

### International Trademark designation:

Simplot<sup>™</sup>, APEX<sup>™</sup> and GAL-Xe<sup>ONE™</sup> are trademarks of J.R. Simplot Company.



### J.R. Simplot Company Professional Products



IN CASE OF EMERGENCY CALL CHEMTREC +1-703-527-3887 • CCN# 12097