

**RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO FISH AND AQUATIC ORGANISMS.**
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



FIFRA Section 2(ee) Recommendation

THIS RECOMMENDATION IS ONLY FOR THE STATES OF CONNECTICUT, DELAWARE, GEORGIA, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, NEW JERSEY, NORTH CAROLINA, OHIO, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, VERMONT, VIRGINIA, WEST VIRGINIA AND WASHINGTON



EPA Reg. No. 59639-209

**FOR CONTROL OF SPOTTED WING DROSOPHILA
IN APPLE, PEAR AND STONE FRUIT**

This FIFRA section 2(ee) recommendation expires on December 31, 2025.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using Asana® XL Insecticide. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

For Control of Spotted Wing Drosophila in Apple, Pear and Stone Fruit				
Crop	Application Rate fl oz/A	Preharvest Interval (PHI)	Field Re-entry Interval	Maximum Seasonal Rate ai lb/A
Apple	4.8 to 14.5	21 days	12 hours	0.525
Pear	4.8 to 14.5	28 days	12 hours	0.375 (0.225 between bloom and harvest)
Stone Fruit	4.8 to 14.5	14 days	12 hours	0.375 (0.3 between bloom and harvest)
Application Timing				
Applications to protect against Spotted Winged Drosophila (SWD) damage should begin as fruit starts to ripen or when local monitoring indicates the first presence of SWD adults in your area. Repeat applications may be required to protect the crop up until harvest. Post harvest orchard floor applications can also be used to control SWD adults laying eggs on fruit that has fallen to the ground.				

Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating Spotted Winged Drosophila in your area.

Precaution: Do not graze livestock on treated orchard floors.

Please contact Valent U.S.A. LLC at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2020 Valent U.S.A. LLC

Asana is a registered trademark of Sumitomo Chemical Company, Ltd.

Registrant: **Valent U.S.A. LLC**
P.O. Box 5075
San Ramon, CA 94583