



Water Soluble Bio Insecticide from Neem

For Agricultural and Commercial Use

Keep out of reach of children
CAUTION

Active Ingredient: Azadirachtin.....6%
Other Ingredients:.....94%
Total:.....100%

Net Contents: See individual packet EPA Reg. No. 81899-4 EPA Est. No. 87465-IND-001

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

Available from Arborjet, Inc. 99 Blueberry Hill Road, Woburn, MA 01801
Manufactured for SoluNeem, Inc. 1050 Bridgeway, Sausalito, CA 94965

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **For medical emergencies, phone 24 hours a days, National Pesticide Telecommunication Network at 1-800-858-7378.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates. Terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any irrigation system unless the chemigation instructions on this label are followed.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirement for the protection of agriculture workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Shoes and socks
- Chemical resistant gloves

NON-AGRICULTURAL USE REQUIREMENTS

These requirements apply to uses of this product that are NOT within the WPS for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses including golf courses and other non-agricultural uses, do not enter treated areas without protective clothing until sprays have dried.

PRODUCT DESCRIPTION

AzaSol is a pale yellow/white, amorphous powder containing 6% by weight azadirachtin. It will instantly dissolve in water to give a solution that is ready for spray application for pest control. Non-Oil based and highly effective as a powder.

Use AzaSol for pre-harvest treatment of fruits and vegetables in case of sudden pest infestations.

AzaSol is effective on a very wide spectrum of insects and pests as listed on this label.

Use AzaSol on a wide variety of plants as listed indoors and outdoors. If plans are made to use AzaSol on plants not listed on this label, it is recommended that a small area such as a leaf, stem, or branch be "test sprayed", first and checked several days later to make sure that leaf wilting or damage does not occur.

When used as directed, AzaSol will destroy targeted insect larvae when they, (1), eat sprayed plants, or (2), come in contact with the spray. AzaSol eliminates insects by stopping the insect's growth, and is effective on all insects listed, insect larval stages and pupae.

MODE OF ACTION

AzaSol controls insects in the larval, pupal, and nymphal stages by interfering with the metabolism of ecdysone. Insects typically die between larval to larval, larval to pupal, nymph to nymph molts, or during adult eclosion.

COMPATIBILITY

AzaSol has been found to be compatible with the most commonly used non-alkaline insecticides, fungicides and water soluble fertilizers in the neutral pH range. Check compatibility by using the correct proportion of each the products application rate in a quart or gallon container. Solubilize AzaSol first in the mixture. Test the tank-mix combinations for possible adverse effects (such as settling out, flocculation, etc.) and for phytotoxic effects on a small sample of plants prior to use. As environmental conditions can alter the interactions between compounds, test compatibility for both new and previously used combinations. Avoid mixtures of several materials and very concentrated spray mixtures.

Do not use AzaSol with Bordeaux mixture, triphenyltin hydroxide, lime sulfur, Rayplex iron or other highly alkaline materials. Use mildly alkaline mixtures immediately after mixing to prevent loss of insecticidal activity.

When using AzaSol in combination with other products, use AzaSol at the rate, or half the rate, specified in the Use Rate table. Follow the directions for use, precautions and limitations for use on the entire product labels used in the combination.

Some suggested tank mix combinations are as follows: AzaSol plus endosulfan*, AzaSol plus chlorpyrifos*, AzaSol plus acephate*, AzaSol plus Bacillus thuringiensis* (BT), AzaSol plus bifenthrin*, AzaSol plus esfenvalerate*, AzaSol plus abamectin*, AzaSol plus diflubenzuron*, AzaSol plus pyrethrum + piperonyl butoxide (for fogging use)*

Always follow the manufacturer's Directions for Use and Precautionary Statements. Use AzaSol on vegetables, coconut palms and other food crops with such chemicals as Endosulfan.

APPLICATION INSTRUCTIONS

AzaSol is exempt from tolerances and may be applied as directed to any food crop up to and including the day of harvest at a rate not exceeding 0.75 lb (20 grams active ingredient) per acre per application.

READ ALL DIRECTIONS AND PRECAUTIONS BEFORE USE

To apply AzaSol select a suitable power or pump pressure trigger sprayer or a hand held trigger type sprayer that will deliver a forceful, fine, leaf, fruit covering, wetting, spray mist. To get thorough spray coverage on waxy or pubescent plant surfaces the addition of small amount of a suitable sticker agent (such as NuFilm P) added to the spray mix, at the recommended rates may give better foliage, insect coverage and control.

APPLICATION METHOD AND EQUIPMENT: Apply AzaSol as a foliar spray or a drench to soil or soil less media (e.g., greenhouses and mushroom houses) to control insects and nematodes. When needed, drench soil to control soil borne pests, including soil-borne larvae of foliar insect pests. When applying as a drench, avoid excessive leaching. Apply AzaSol through sub-surface soil treatment equipment (e.g. turf grass). To repel adult flies, apply through fogging equipment.

Always follow equipment manufacturer's use directions.

Apply AzaSol by using any powered or manual pesticide application equipment, which includes but is not restricted to: high-volume, low-volume, ultra-low volume, electrostatic, fogging, and chemigation. Follow the original manufacturer's recommendations when using these types of equipment.

For optimum results, 2 to 3 applications made at 7 to 10 day intervals is recommended, unless otherwise specified. Foliar applications should be made to both side of leaves. In addition, a sticker agent used as per the manufacturer's recommendations may improve product performance.

AZASOL USE RATE RECOMMENDATIONS FOR KEY PESTS BY USE SITE

AzaSol label rates specify dry ounce (weigh/acre (high rate) and tsp or tbsp/1000 sq. ft. (low rate). These label rates provide a high and low dose application of Azasol.

Abbreviation & Conversion Table

tsp	teaspoon
tbsp	tablespoon
A	acre
3 tsp = 1 tbsp	
1 A = 43,560 sq. ft.	

AZASOL USE RATE RECOMMENDATIONS FOR KEY PESTS BY USE SITE (continued)

High Rate 6oz/50 gal water/Acre (6oz. is approximately 170 tsp/50 gal water/acre) (approximately 4 tsp/1000 sq ft.)

Low Rate 1 tsp/1 gal water/1000 sq. ft. (1 tsp. is approximately 44 tsp/ 44 gal water/ acre)

USE RECOMMENDATION:

Low Rate (Recommended for preventative treatments before signs of infestation.) 1 tsp/1000 sq. ft.

Medium Rate (Recommended for most treatments. For preventative to medium infestations when pests are present.) 2-3 tsp/1000 sq. ft.

High Rate (Recommended for difficult to manage pests or high infestations.) 4 tsp/1000 sq. ft.

Use the tables below to determine the appropriate use rate for your site/pest combination.

AZASOL PEST CONTROL CHART:

USE RATES for indoor and outdoor plants including, FOOD CROPS, TREES, TURFGRASS, NURSERY, GREENHOUSE, INTERIORSCAPE & LANDSCAPE PLANTS

PEST	RATES: AzaSol oz's./acre-tsp./1,000 sq. ft.	COMMENTS For Spray, Drench or Chemigation.
WHITEFLIES, such as: Green-house whiteflies, Silverleaf white flies, Woolly whiteflies.	6 oz in 50 gal water/A 1 tsp/1 gal water /1000 sq.ft.	Make sure that spray covers upper, lower and all surfaces of leaves fruit and twigs.
LEAF MINERS, such as Azalea leafminers, Birch leafminers, Citrus Leafminers, Serpentine leafminers.	6 oz in 50 gal water/A 1 tsp/1 gal water /1,000 sq.ft.	Apply to new growth in spring before new larvae enter plant foliage. Repeat application at 10 to 14 day intervals if new infestations are expected.
SCALE, Crawlers: such as Brown Soft scale, California red scale, Coffee Scale, Olive Scale, San Jose Scale	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	Make sure to thoroughly spray upper, lower and all surfaces of leaves and twigs.
MEALY BUGS Such as Citrus Mealybugs	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq ft	Spray to thoroughly cover twigs and leaves.
THRIPS, such as: Citrus thrips, Onion thrips, thrips palmi.	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq ft	Spray in spring when young nymphs first appear on foliage.
APHIDS, such as: Cotton aphids, Green peach aphids, Pea aphids, Potato aphids	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	Spray to wet lower side of leaves when "leaf curl" first appears.
PSYLLIDS, such as : Pear psylla	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	Spray for new "instar" nymphs appearing on new discolored foliage.
BUGS, Nymphs of: such as Box elder bugs, Chinch bugs, Lygus bugs, spittle bugs, stink bugs	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	Spray early when nymphs are young. AzaSol will control "instar" growth until they die.
FLIES, Larvae of: such as Blueberry Maggot, Cherry Maggot, Crane Flies, Fruit flies, Midges, Onion Maggots, Tip worms, Walnut husk fly larvae.	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	For food and Non food crops spray when larvae first appear
SAWFILES, Larvae of: such as: European Pine Sawflies, Yellow Headed pine sawflies	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	Spray when first larvae appear when plants start new growth
CATERPILLARS, such as: Armyworms, Artichoke plume moth, Bagworms, Bollworms, Budworms, Cabbage butterflies, Cabbage loopers, Cankerworms, Caseworms, Corn Earworms, Cutworms, Diamond back moths, Fireworms, Fruitworms, Grapeleaf, skeletonizer, Gypsy moths, Hornworms, Imported cabbage worm, leaf perforators, Leafrollers, Melonworms, Navel orangeworms, Oblique banded Leafrollers, oriental fruit moths, Pickleworms, Pine tip moths, Pinworms, Red banded leaf rollers, Sod webworms, Soybean loopers, Tent Caterpillars, Tobacco bud worms, Tussocks moth larvae.	6 oz in 50 gal water/A 1 tsp/1 gal water/1000 sq.ft.	Spray when first larvae worms appear. Repeat applications in 7 to 10 days. For continued pest control in the spring or fall when insect infestations are expected spray ornamentals and other plants at intervals of 2 to 3 weeks.
BEETLES, larvae of : such as Bark beetles, Blueberry Flea beetles, Boll weevils, Colorado potato beetles, Flea beetles, Japanese beetles, Leaf beetles, Mexican bean beetles, Phylloxera, Rose Chafers, Twig girdlers	6 oz in 50 gal water/A 1 tsp/ 1 gal water/1000 sq.ft.	Spray when pests first appear. For Food crops, Repeat application after 7 to 10 days. Do not use with oil. Make sure that all plant surfaces are thoroughly spray treated. Repeat in 5 to 7 days if required.

WEEVILS, such as Black vine weevils, Pepper weevils, Strawberry vine weevils.	6 oz in 50 gal water/A 1 tsp/ 1 gal water/1000 sq.ft.	Foliar anti-feedant sprays will stop adult feeding. Make at least 3 to 4 applications 10 days apart.
BORERS, Larvae of: Peach twig borer, Peach tree borers, Cranberry borers.	6 oz in 50 gal water/A 1 tsp/ 1 gal water/1000 sq.ft.	Thoroughly spray in spring after egg hatch to control young larvae.
MOLE CRICKETS, nymphs and young "instars". Turf Treatment	6 oz in 50 gal water/A 1 tsp/ 1 gal water/1000 sq.ft.	For turfgrass, spray to drench turf for young cricket nymphs in spring. Stops young from growth to adults.
MUSHROOM FLIES Nematodes and Phorid Flies	Mix ½ oz. in 1 to 2 gal of water and mist over, (or drench) 1,000 sq. ft.	See "For Mushrooms" section on this label.

USE SITES FOR AZASOL

AzaSol can be used on Green-house: food crops, such as: Brassica (cole) crops, cucurbits, eggplants, herbs and spices, legumes, peppers, tomatoes.

MUSHROOMS, Varieties such as: Agaricus, enoki, maitake, oyster, shitake and other specialty mushrooms

Food crops including, Root, and tuber vegetables such as: Artichoke, beets, carrots, ginger, horseradish, potatoes, radishes, rutabagas, sweet potatoes, turmeric, tunips, yams.

Leafy vegetables (including Brassica leafy vegetables) such as: Amaranth, broccoli, Brussels sprouts, cabbage, cauliflower, celery, Chervil, Chinese cabbage, collards, cress, endives, fennel, kale, kohlrabi, lettuce, mizuna, mustard greens, parsley, purslane, rape greens, rhubarb, spinach, Swiss chard.

Legume vegetables such as: Beans (field, kidney etc..) chick-peas, cowpeas, guar, jackbeans, lablab beans, lentils, peas, pigeon peas, soybeans, sword beans.

Fruiting vegetables such as: Eggplants, ground-cherries, pepinos, peppers, pimentos, tomatillos, tomatoes.

Cucurbit vegetables such as: Bitter melons, Chayotes, Chinese wax gourds, citron melons, cucumbers gherkins, gourds, muskmelons (such as cantaloupes, casabas cranshaw etc..) pumpkins, squash, watermelons.

Citrus Fruits such as: Calamondins, citrus citrons, citrus hybrids, Grapefruits, Kumquats, Lemons, Limes, Mandarins, Oranges, pumellos, satusuma mandarins.

Pome fruits such as: Apples, crabapples, loquats, mayhaws, oriental pears, pears, quinces.

Stone fruits such as: Apricots, cherries, nectarines, peaches, plums, prunes.

Berries such as: Blackberries, cranberries, blueberries, currants, cranberries, elderberries, gooseberries, huckleberries, loganberries, raspberries, strawberries, youngberries.

Cereal grains such as: Barley, buckwheat, corn, millet, oats, popcorn, rice, rye, sorghum, teosinites, triticale hybrids, wheat, wild rice,

Herbs and spices including but not limited to: Allspice, ahgelica, anise, annatto, balm, basil, black and white peppers, borage, burnet, camomile, caper buds, cardamom, caraway, cassia, catnip, celery seeds, chervil, chives, cinnamon, caraway, cloves, coriander (cilantro), costmary, cumin, curry leaf, dills, fennels, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, mace, marigolds, marjoram, mustard seeds, nasturtium, nutmeg, parsley, pennyroyal, pepper (black & white), poppy seeds, rosemary, rue, saffron, sage, savory, sweet bay (bay leaf), tansy, tarragon, thyme, vanilla, wintergreen, woodruff, wormwood.

Bulb vegetables such as: Garlic, leek, onions, shallots.

Nuts such as: Almonds, beechnuts, Brazil nuts, butternuts, cashews, chestnuts, chinquapin, coconuts, filberts, hickorynuts, imacadamia, pecans, pistachios, walnuts.

Oilseed crops such as: Canola, castor, crambe, guar, jojoba, peanuts, rape, safflower, sesame, soybean, sunflower.

Tropical fruits such as: Atemoyas, bananas, breadfruits, cherimoyas, durians, guavas, malangas, mangos, papayas, passionfruits, starfruits.

Other food & non-food crops such as: Asparagus, avocados, birdseed, cacao, coffee, edible flowers, feijoa, figs, ginseng, grapes, guayule, hops, kiwis, okras, olives, palms, papayas, pawpaws, persimmons, pineapples, rambutans, sugarcane, tamarillos, tea, tobacco, water chestnuts, watercress.

Ornamental Plants such as: African violets, ageratum, aster, aucuba, begonia, cacti, calendula, calla, carnation, ceanothus, chrysanthemum, cineraria, coleus, cyclamen, daffodil, dahlia, delphinium, ficus, foliage plants, fuschia, gardenia, geranium, gloxinia, hyacinth, hydrangea, iris, ivy, lily, maidenhair fern, marigold, narcissus, orchid, pansy, pelargonium, peony, phlox, pittisporum, poinsettia, pyracantha, rubber plant, snapdragon, stock, tulip, wandering jew, yucca, zinnia.

Ornamental Trees and Shrubs such as: Andromeda, Arbovitae, ash, Austrian pine, azalea, beech, birch, birdsnest spruce, blue spruce, bougainvillea, boxwood, butternut, camellia, cedar, chamaecyprus, dogwood, Douglas fir, elm, euonymus, firethorn, forsythia, hackberry, hawthorn, hemlock, hickory, holly, honeylocust, horsechestnut, ilex, juniper, larch, laurel, lilac, linden, London plane, magnolia, manvillea, maple, mimosa, mountain ash, myrtle, oak, pachysandra, peach, pine, Pinota, pines, plane tree, poplar, privet, quince, rhododendron, roses, spruce, sycamore, white cedar and white pine.

(continued on next page)

USE SITES FOR AZASOL(cont.)

TURF GRASSES such as: Bent grass, Bermuda grass, Fescue, Bluegrass, annual & perennial; Ryegrass, annual & perennial; Buffalo grass, St. Augustine grass, Centipede grass, Wheat grass, Zoysia grass.

For control of Sod Webworms, Cutworms, Aphids, Leafhoppers, ants and chiggers: use a suitable pressure sprayer and mix 1 tbsp. in 2 to 3 gal of water and apply to 2,500 sq ft of turf. Apply when insect larvae first appear and if necessary repeat application in 10 to 14 days. The use of an approved "spreader sticker" may help the spray to penetrate turf down to the larvae/ worm feeding area.

AzaSol For Mushrooms and the Mushroom House

For Mushroom Flies, Nematode and Phorid Flies use AzaSol at the rate indicated on the PEST CONTROL CHART as a drench to the casing layer, media or compost. Make 4 to 5 applications 7 to 10 days apart. To repel fly adults, apply with fogging equipment at the first sign of activity. Can be applied between breaks up to the final flush.

AZASOL APPLICATION BY TRUNK INJECTION

Directions for Use

Inject into the trunk flare or within 36" of the soil level. Place the injection sites in the first few sapwood elements (growth rings). Drill holes using a clean sharp drill bit (brad point drill bits are recommended). Drill through the bark and into the sapwood. When using the Arborjet Arborplug®, drill a minimum of 16 mm (5/8") into the sapwood. Trunk inject product into the tree's sapwood, the conductive tissue that moves water to the canopy.

Calculating Application Rate

The dosage and number of application sites are based on tree diameter (DBH). To determine the number of application sites and dose rate per tree:

1. Determine the Tree Diameter (DBH):

Measure the tree diameter in inches (or centimeters) at chest height [54" (135 cm) from the ground] to find the diameter at breast height (DBH). If measuring tree circumference, divide the circumference by 3 to obtain the DBH.

2. Calculate # of Injection Sites:

By Micro-injection (QUIK-jet®, Air Hydraulic): Calculate the number of injection sites by dividing the DBH in inches by 2 (or cm DBH by 5). This is equivalent to one drill hole for every 6" (15 cm) of tree circumference. By Micro-infusion® (TREE I.V.): Calculate the number of injection sites by dividing the DBH in inches by 3 (or cm DBH by 7.5). This is equivalent to one drill hole for every 8" (20 cm) of tree circumference.

3. Determine the Dose:

Measure the amount of AzaSol needed following the table:
Use Rate Recommendations for Tree Injection.

4. Determine Dose per Injection Site:

Divide the total dose by the number of injection sites to determine the dosage per injection site.

In resinous conifers (such as pine and spruce) you may inject each site shortly after drilling to avoid slow uptake on account of resin flow.

In palms, only one injection site is generally required.

1. Locate the application site 1-3' (30-90 cm) from the soil level.
2. Drill depth is 1/3 the total diameter or 4" (10 cm) deep into the stem (whichever is less).
3. Refer to table: *Use Rate Recommendations for Palm Injection* for dosages to apply.

Application Equipment

AzaSol may be used with the Arborjet Tree Injection Systems or with other tree injection devices that meet the label requirements. For all injection systems, read carefully and follow the manufacturer's direction for use.

Recommendations for Tree Injection.

Application in Trees

1. Inject 4 mLs of solution every 6" (15 cm) of stem circumference in trees <8" DBH (20cm).
2. Inject 6 mLs of solution every 6" (15 cm) of stem circumference in trees 8-16" DBH (15- 40cm).
3. Inject 8 mLs of solution every 6" (15 cm) of stem circumference in trees >16" DBH (>40cm).

USE RATE RECOMMENDATIONS FOR TREE INJECTION

DBH"	cm DBH	level tsp. Azasol	Milliliters of water	ml/2.5 cm (inch) DBH	Average Number of Injects	mL/Inject
5	12.5	2	12	2	3	4
6	15.0	2	12	2	3	4
7	17.5	2	16	2	4	4
8	20.0	4	24	3	4	6
9	22.5	4	30	3	5	6
10	25.0	4	30	3	5	6
11	27.5	6	36	3	6	6
12	30.0	6	36	3	6	6
13	32.5	6	42	3	7	6

USE RATE RECOMMENDATIONS FOR TREE INJECTION (cont.)

14	35.0	6	42	3	7	6
15	37.5	8	48	3	8	6
16	40.0	8	48	3	8	6
17	42.5	10	72	4	9	8
18	45.0	12	72	4	9	8
19	47.5	12	80	4	10	8
20	50.0	12	80	4	10	8
21	52.5	14	88	4	11	8
22	55.0	14	88	4	11	8
23	57.5	14	96	4	12	8
24	60.0	16	96	4	12	8
25	62.5	16	104	4	13	8

For trees larger than 25" (62.5 cm) DBH apply 4 mL/2.5 cm (inch) DBH.

USE RATE RECOMMENDATIONS FOR PALM INJECTION

Canopy or Tree Size	Tsp AzaSol	Milliliters of water	Minimum Number of Injection Points Needed
Small Canopy or Tree	2	10	1
Medium Canopy or Tree	4	20	1
Large Canopy or Tree	6	30	1

DOSE/RATE SPRAY APPLICATIONS

Directions for Use

AzaSol is measured in dry ounces (weight) and approximate teaspoons for each packet size. Packet sizes come in depending on the type of pest and timing of treatment you may use low, medium, or high rates of application.

NOTE: Low Rates are recommended for preventative treatments before signs of insects. **Medium Rates** are recommended for most treatments for preventative to medium infestations when pests are present. **High Rates** are recommended for difficult to manage pests or for heavy infestations.

AzaSol packet recommendations for mixing, and dosing in spray applications.

Packet sizes are recommended by using the most economical size and no partial packets.

Sq. Ft.	Low Rate		Medium Rate		High Rate	
	Azasol	Water	Azasol	Water	Azasol	Water
1,000			(1) 2 tsp packet	1-2 gal	(2) 2 tsp packets	1-4 gal
5,000	(3) 2 tsp packets	5-10 gal	(5) 2 tsp packets	5-10 gal	(1) 0.75 oz packet	10-20 gal
10,000	(5) 2tsp packets	10-20 gal	(1) 0.75 oz Packet	10-20 gal	(2) 0.75 oz packets	20-40 gal
20,000	(1) 0.75oz packet	20-40 gal	(2) 0.75 oz packets	20-40 gal	(4) 0.75 oz. packets	50-100 gal
(1 Acre) 43,560	(2) 0.75 oz packets	50-100 gal	(4) 0.75 oz. packets	50-100 gal	4) 0.75 oz. packets	100-200 gal

Low Rate: 0.07 oz. (approx. 2 tsp) / 2-4 gal of water/ 2,000 sq ft. or

Medium Rate: 0.07 oz. (approx. 2 tsp) / 1-2 gal of water/ 1,000 sq ft.

High Rate: 0.14 oz. (approx. 4 tsp) / 2-4 gal of water/ 1,000 sq ft.

MIXING

Re-sealable Mylar Packets:

AzaSol is sealed in mylar packets to ensure air tight and water tight seal to protect powdered AzaSol. Carefully open mylar packet and dispense the powder into the mixing tank. You can use the entire contents or you can measure a partial amount to meet mixing and rates specified for application. Airborn powder: You can reduce the amount of air born powder by avoiding windy conditions, using the entire contents of packet, and by emptying contents into a partially filled tank. Agitate tank mix.

It's recommended to use a sticking agent to increase product coverage on plant tissue

(continued on reverse side)

CHEMIGATION OF AZASOL

General Information

Apply this product only through drip (trickle) or sprinkler (center pivot, lateral move, end tow, side roll, traveler, big gun, solid set, or hand move), flood (basin) irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Solubilize AzaSol with water before introduction into the system; use the diluted mixture within 8 hours. Do not apply in irrigation water if the pH exceeds 7.0. The optimum pH for application is a range of 5.5 to 6.5. If needed, the pH of the irrigation water can be adjusted by use of a suitable buffering agent. Agitation is necessary. Apply at the rate stated in the Directions for Use using sufficient water to achieve an even distribution within an 8 hour period. Do not apply AzaSol at a rate that exceeds 20 grams active ingredient per acre. If applying AzaSol in combination with other products refer to the compatibility statement in the Directions for Use section.

OBSERVE THE FOLLOWING PRECAUTIONS IF YOUR CHEMIGATION SYSTEM IS CONNECTED TO A PUBLIC WATER SYSTEM

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.

Chemigation systems connected to the public water systems must contain a functional, reduced-pressure zone (RPZ), backflow prevention the functional equivalent in the water supply line upstream from the point of pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in the cases where there is not a water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock,

Do not apply when wind speeds favor drift beyond the area intended for treatment.

STATEMENTS CONCERNING THE OPERATION OF SPRINKLER CHEMIGATION; DRIP (TRICKLE), UTILIZING A PRESSURIZED WATER AND PESTICIDE INJECTION SYSTEM

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick – closing check valve to prevent the flow of fluid back toward the injecting pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid – operated valve located on the intake side of the injection pump and connected to the system, interlock to prevent fluid from being withdrawn from the supply tank when the irrigating system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STATEMENTS CONCERNING THE OPERATION OF FLOOD (BASIN) IRRIGATION UTILIZING GRAVITY FLOW OR PRESSURIZED WATER AND PESTICIDE INJECTION SYSTEM.

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drip structure or weir box to decrease potential for water source contamination from back flow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements.

- The system must contain a functional interlocking check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of the fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Do not store this product above 100°F or below 20°F for extended periods of time. Store product in the original labeled container in a cool, dry, locked place out of reach of children. Keep containers tightly closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

IMPORTANT: PLEASE READ BEFORE USE

To the extent consistent with applicable laws, SoluNeem, Inc. warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to herein, when used in accordance with its directions; and (c) that the directions, cautions and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, and upon reports of field experience. Testing has not been performed on all varieties of food crops, and plants, in all states, or under all application, weather and crop conditions. There are no express warranties other than those set forth herein. SoluNeem, Inc. neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. SoluNeem, Inc. expressly excludes and disclaims all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality of performance.

This warranty does not extend to, and the user shall be solely responsible for, any loss or damage that results from the use of this product in any manner that is inconsistent with this label's directions, or cautions.



Available from:

ARBORJET
Revolutionary Plant Health Solutions