BIOMIN® BOOSTER 126

1-0-0

Amino Acid Chelated Minerals for Soil and Foliar Application

Product Information

BIOMIN BOOSTER 126 contains chelated minerals designed for soil and foliar application. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct micronutrient deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizer.

GUARANTEED ANALYSIS

Total Nitrogen (N)	1.0%
1.0% Water Soluble Nitrogen	
Magnesium (Mg)	0.8%
0.8% Chelated Magnesium	
Boron (B)	0.025%
Copper(Cu)	0.1%
0.1% Chelated Copper	
Iron (Fe)	0.5%
0.5% Chelated Iron	
Manganese(Mn)	1.0%
1.0% Chelated Manganese	
Zinc (Zn)	3.0%
3.0% Chelated Zinc	

Derived from cupric sulfate, ferrous sulfate, magnesium sulfate, manganese sulfate, sodium borate, zinc sulfate and hydrolyzed vegetable protein.

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

PREVENTS AND CORRECTS
MINERAL DEFICIENCIES

NET CONTENTS: 2.5 Gallons (9.45 liters) Weight per Gallon: 9.8 lb (4.5 kg) @ 68°F

U.S. Pat.# 5.504.055

Lot #:



Directions and General Recommendations

SOIL APPLICATION

BIOMIN BOOSTER 126 can be applied before or at planting as banding or broadcast. The application rate is 1 quart to 1 gallon per acre depending on soil conditions and application methods. For maintenance, 1 quart per acre is recommended. BIOMIN BOOSTER 126 should be diluted with enough water to cover the entire acre.

FOLIAR APPLICATION

FIELD CROPS: Apply 16-32 oz. per acre (1.25 to 2.5 liters/hectare) of BIOMIN BOOSTER 126 when crops are about 6-12 inches (15-30 cm.) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

FRUIT TREES AND GRAPES: Apply 16-32 oz. per acre (1.25 to 2.5 liters/hectare) of BIOMIN BOOSTER 126 starting at bud break or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

VEGETABLE CROPS: Apply 16-32 oz. per acre (1.25 to 2.5 liters/hectare) of BIOMIN BOOSTER 126 at about 2 - 4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1-3 weeks if necessary.

ALMONDS, WALNUTS, PISTACHIOSAND OTHER NUT CROPS: Apply 16-32 oz. per acre (1.25 to 2.5 liters/hectare) before bloom. Repeat the application if necessary.

For best results, apply **BIOMIN BOOSTER 126** when the crop is in active growing stage, after irrigation or natural rain fall. For foliar application, it should be diluted with at least 100 times of water. **BIOMIN BOOSTER 126** will lower high pH of water and decrease the hydrolysis of most pesticides.

BIOMIN BOOSTER 126 can be applied to: Strawberry, Lettuce, Tomato, Celery, Stone Fruits, Vegetables, Grapes, Commercial Turf, And Ornamental Gardens.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm