BIOMIN® Cal-Boro CALCIUM with BORON



1-0-0
Minerals for Soil and Foliar Application

Product Information

BIOMIN CAL-BORO contains calcium and boron designed for soil and foliar application. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct micronutrient deficiencies and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizer.

GUARANTEED ANALYSIS

Total Nitrogen (N)	1.0%
1.0% Water Soluble Nitrogen	
Calcium (Ca)	5.0%
5.0% Chelated Calcium	
Boron (B)	1.0%

Derived from calcium carbonate, sodium tetraborate and hydrolyzed vegetable protein.

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

PREVENTS AND CORRECTS
MINERAL DEFICIENCIES

NET CONTENTS: 2.5 Gallons (9.45 liters) Weight per Gallon: 9.3 lb (4.2 kg) @ 68°F

U.S. Pat.# 5,504,055

Lot #:



Directions and General Recommendations

SOIL APPLICATION

BIOMIN CAL-BORO can be applied before or at planting as banding or broadcast. The application rate is 1 quart to 1 gallon per acre (2.5 to 10 liters/hectare) depending on soil conditions and application methods. For maintenance, 1 quart per acre (2.5 liters/hectare) is recom-mended. Biomin Cal-Boro should be diluted with enough water to cover the entire acre.

FOLIAR APPLICATION

FIELD CROPS: Apply 2-3 quarts per acre (5-7.5 liters/hectare), when crops are about 6-12 inches (15-30 cm)high or when deficiency symptoms appear. Repeat the application in 2-4 weeks if necessary.

VEGETABLE CROPS: Apply 2-3 quarts per acre (5-7.5 liters/hectare) at about 2-4 weeks after emergence, or when deficiency symptoms appear. Repeat the application in 1-3 weeks if necessary.

FRUIT TREES AND GRAPES: Apply 2-3 quarts per acre (5-7.5 liters/hectare) in early spring, when new leaves are emerged or when deficiency symptoms appear. Repeat the application in about 4 weeks if necessary.

ALMONDS, WALNUTS, PISTACHIOS AND OTHER NUT CROPS: Apply 2-3 quarts per acre (5-7.5 liters/hectare) before bloom. Repeat the application if necessary.

For best results, apply when the crop is in active growing stage, after irrigation or natural rainfall. For foliar application, it should be diluted with at least 100 times of water.

CAUTION - This product contains Boron. Do not apply to Boron sensitive crops. Use of the product on any crop other than those recommended may result in serious injury to the crop(s).

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm