SASOUTHERN 5 0 AG_®

BRUSH

For Outdoor Residential Use Only

Active Ingredient(s):
Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), as the triethylamine salt*

8.8%
Other Ingredients
91.2%
Total
100.0% Contains 6.36% Triclopyr acid - 0.6 lb/ gallon.

Keep out of reach of children **CAUTION PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Se label booklet for additional PRECAUTIONARY STATEMENTS, FIRST AID, and DIRECTIONS FOR USE.



Net Contents Liquid: 1 Pint (.474 liters)

Distributed by: Southern Agricultural Insecticides, Inc. Palmetto, Fl 34220 Hendersonville, NC 28793 Boone, NC 28607 F1 EPA Est. No. 829-FL-1 EPA Reg. No. 62719-226-829



Keep out of reach of children CAUTION PRECAUCION

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Precautionary Statements Hazards to Humans and Domestic Animals CAUTION

Causes Moderate Eye Irritation * Harmful If Swallowed

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Wear protective clothing such as a long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material, and shoes and socks when applying this product.

User Safety Recommendations

Users should: • Wash hands before eating, drinking. chewing gum, using tobacco, or using the toilet. • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in original container and keep out of reach of children. Store in a cool, dry place. Avoid contamination of food and foodstuffs.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container

If empty: Place in trash or offer for recycling if avail-

If partly filled: Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

General Information

For Outdoor Residential Use Only

SA-50 brand BRUSH KILLER is recommended for the control of unwanted woody plants, vines and broadleaf weeds around homes, cabins, fences, walkways and other non-crop areas.

General Use Precautions and Restrictions

*Do not use on or around edible food or food crops. *Avoid contact with eyes or clothing during and after application.

*Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. *Entry Restrictions: Do not enter or allow others to

enter the treated area until sprays have dried. *Do not spray when the wind is blowing toward desirable plants.

*Do not apply this product directly to, or allow to drift to grapes, vegetable crops, flowers, ornamentals or other desirable foliage. Applications should be made when wind is low and there is little chance of spray drift. *Do not apply with nozzles that produce a fine-droplet

*Measuring utensils and spray equipment should not be reused for purposes other than herbicide applica-

consequential or incidental damages or losses.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no such case shall Seller be liable for

The terms of the Warranty Disclaimer. Terms and Conditions of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

IN CASE OF EMERGENCY CALL: CHEMTREC 1-800-424-9300

For information or questions you may contact us by email us at sales@southernag.com

Inherent Risks of Use

Wild Lettuce, Wild Violet.

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buver.

Woody Plants and Vines Controlled Include:

Alder, Arkansas Rose, Arrowwood, Ash, Aspen, Beech,

Birch, Blackberry, Blackgum, Box Elder, Brazilian Pep-

per, California Rose, Cascara, Ceanothus, Cherry,

Chinquapin, Choke Cherry, Cottonwood, Crataegus

(Hawthorn), Douglas Fir, Dogwood, Elderberry, Elm, Ha-

zel, Honeysuckle, Hornbeam, Kudzu, Locust, Madrone,

Maples, Mesquite, Mimosa, Mulberry, Oaks, Persimmon,

Pine, Poison Ivy, Poison Oak, Poplar, Raspberry, Salm-

onberry, Sassafras, Scotch Broom, Sumac, Sweetbay

Magnolia, Sweetgum, Sycamore, Tanoak, Thimbleberry,

Trumper Creeper, Tulip Poplar, Virginia Creeper, Western

Hemlock, Willow, Winged Elm, Wild Grape, Wild Rose,

Include: Bindweed, Burdock, Canada Thistle, Chickory,

Clover, Curly Dock, Dandelion, Field Bindweed, Field

Ragweed, Smartweed, Spurge, Tansy Ragwort, Vetch,

Horsetail Rush, Henbit, Lambsquarter, Oxalis, Plantain.

Perennial Broadleaf Weeds Controlled

Limitations of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

APPLICATION METHODS

For best results, apply as a full coverage foliar spray when brush and weeds are fully leafed out and actively growing. Wilting or other symptoms will occur in 1-6 weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

Using a hose-end or tank sprayer, mix 8-16 tablespoons (4-8 ounces) in enough water to make on gallon mixed spray. For mature or hard-to-control plants (such as ash, blackgum, choke cherry, elm, maple, oak, pine or winged elm), or during drought conditions, use the highest rate of application indicated. Resprouting may occur the year following treatment when lower rates are used. The addition of a surfactant/spreader/sticker (such as SA-50 brand Surfactant for Herbicides) enhances herbicidal action and improves distribution of the spray on hard-to-wet plant surfaces. Apply at the

rate of one gallon mixed spray to approximately 500 square feet of area.

Coverage may vary depending on the amount of foliage to be treated. Using coarse spray pattern and low pressure, apply as thorough foliage spray to wet all leaves. See General Use Precautions and Restrictions.

Stump Treatment

To kill stumps and prevent sprouting, apply the undiluted product with a paint brush or backpack sprayer to completely cover the freshly cut surface. Do not reuse paint brush. Wrap in several layers of newspaper and discard in trash.

Vine Control

When treating vines such as poison oak, poison ivy, etc. that grow on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat as directed for stump treatment.


Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY





Active Ingredient(s):
Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid).as the triethylamine salt* ... Other Ingredients Contains 6.36% Triclopyr acid - 0.6 lb/ gallon.


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) Agricultural Chemical: Do not ship or store with


food, feeds, drugs or clothing. Refer to inside of label booklet for additional precautionary information including Directions

Distributed by: Southern Agricultural Insecticides, Inc.

P.O. Box 218 Palmetto, FL 34220 EPA Reg. No. 62719-226-829 EPA Est. No. 829-FL-1







14