## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye and skin irritation. Avoid breathing vapor or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

### FIRST AID STATEMENT

IF INHALED: Move person to fresh air. If person is not breathing call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for advice.

IF IN EYES: Hold eye(s) open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, and continue rinsing. Call a Poison Control Center or doctor for advice.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

#### HOT LINE NUMBER

For emergency information on product, uses, and treatment, etc., call the National Pesticides Information Center at 1-800-858-7378, 6:30 am - 4:30 pm Pacific Time (PT), 7 days a week. Otherwise, call the Poison Control Center at 1-800-222-1222.

## ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

## PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F may cause bursting.

WARRANTY: To the extent consistent with applicable law, Suterra® LLC warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose stated in the Directions for Use. To the extent consistent with applicable law, Suterra® LLC neither makes nor authorizes any agent or representative to make any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material. To the extent consistent with applicable law, Suterra® LLC's maximum liability for breach of this warranty shall not exceed the purchase price of this product. To the extent consistent with applicable law, buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage, and use of this material contrary to label instructions.



#### For the control of INDIAN MEAL MOTH (Plodia interpunctella), MEDITERRANEAN FLOUR MOTH (Ephestia kuehniella), RAISIN MOTH (Cadra figuiliella), ALMOND MOTH (Cadra cautella) and TOBACCO MOTH (Ephestia elutella)

CheckMate<sup>®</sup> Puffer<sup>®</sup> IMM contains a behavior modifying biochemical (pheromone) that disrupts the mating behavior of stored product moths: Indian Meal Moth (*Plodia interpuctella*), Mediterranean Flour Moth (*Ephesita kuehniella*), Raisin Moth (*Cadra figuiliella*), Almond Moth (*Cadra cautella*) and Tobacco Moth (*Ephestia elutella*). CheckMate<sup>®</sup> Puffer<sup>®</sup> IMM is used with the CheckMate<sup>®</sup> Puffer<sup>®</sup> Aerosol Cabinet, an automatic dispenser that delivers the pheromone to disrupt the mating behavior of stored product moths.

## ACTIVE INGREDIENT\*:

(Z,E)-9,12-tetradecadien-1-yl acetate	
OTHER INGREDIENTS:	
TOTAL	

\*Contains: 12.2 g a.i. per canister Net Contents: 11.8 oz (334 grams)

KEEP OUT OF REACH OF CHILDREN CAUTION MANUFACTURED BY: Suterra®

Suterra LLC 20950 NE Talus Place Bend, Oregon U.S.A. 97701 Phone: 1 (866) 326-6737

EPA Reg. No. 73479-16 EPA Est. No. 56336-OR-001 Batch Code #. See bottom of can.

## DIRECTIONS FOR USE IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

# Do not exceed 3.5 grams of pheromone per 1,000 ft<sup>2</sup> per year. (700 g of formulated product per 7,300 ft<sup>2</sup> per year)

METHOD OF APPLICATION: For use in food/feed storage areas of silos, warehouses, food-processing plants, storage areas within retail stores and any areas where stored product moths are a problem. Use a CheckMate® Puffer® Aerosol Cabinet to automatically dispense the contents of the CheckMate® Puffer® IMM canister (herein referred to as "Puffer") In situations where adults do not occur year-round, deploy at the start of adult activity or when males are first captured in monitoring traps. In situations where adults occur year-round, the application can be started at any time. Replace cans when empty.

To avoid accidental spraying during the placing and re-placing of the canisters, ensure that canister nozzles are pointed away during the performance of these tasks.

PRODUCT PLACEMENT: Suspend an activated Puffer Aerosol Cabinet containing a full Puffer canister from secure supports such as walls, pallets, columns or ceiling hangers, ideally 7 to 15 feet from the floor. Deploy the Puffers at a rate of 1 Puffer per 7,300 feet<sup>2</sup> and consider internal airflows: in case of constant internal air flows, ensure that the Puffers are oriented in the same direction as the main airflow. Place Puffers where spray is directed away from walls and open food sources.

APPLICATION NOTES: This is a mating disruption product that interferes with the ability of stored product moths' males to find females and mate. It does not directly affect eggs, larvae or mated females. If first application occurs in highly infested areas, it is recommended to adequately treat the area with insecticides that will effectively control larvae and eggs already present in the area. Previously mated females, larvae and eggs introduced with contaminated raw materials are not affected by the product and may reduce the level of control. Always monitor insect infestation with traps. Complement the mating disruption with effective insecticides whenever it is necessary.

For best results apply at the time or shortly before moths begin to emerge, or when moths are detected in monitoring traps.

# STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal PESTICIDE STORAGE: Store container in cool, dry area away from heat or open flame.

**PESTICIDE DISPOSAL**: Wastes resulting from use of this product may be disposed of on-site or at an approved disposal facility.



CONTAINER HANDLING: Do not puncture or incinerate. Nonrefillable container. Do not reuse or refill this container. If can is empty: Remove from cabinet, place in trash or offer for recycling. If partly filled: Call your local solid waste agency for disposal instructions.