



# Cheetah®

## Herbicide

Cheetah Herbicide will be referred to as Cheetah throughout the label. Cheetah is a non-selective herbicide that provides control of a broad spectrum of broadleaf weeds and grassy weeds.

Cheetah is registered for use: as a burndown treatment prior to planting or prior to emergence of canola, corn, cotton, sweet corn, soybean and sugar beets; post emergence weed control herbicide to be applied on crops containing the glufosinate-resistant gene; post emergence weed control in cotton when applied with a hooded sprayer in-crop; post emergence weed control in listed tree, olives, vine, and berry crops; applied for potato vine desiccation; as a nonselective postemergent herbicide

**ACTIVE INGREDIENT:**

Glufosinate-ammonium\* ..... 24.5%\*\*

**OTHER INGREDIENTS:** ..... 75.5%

**TOTAL:** ..... 100.0%

\*CAS Number 77182-82-2

\*\*Equivalent to 2.34 pounds of active ingredient per U.S. gallon.

Shake Well, Agitate or Recirculate Before Use

## KEEP OUT OF REACH OF CHILDREN

# CAUTION

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle. (If you DO NOT understand the label find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND PRECAUTIONARY STATEMENTS

EPA Reg. No. 71368-112

Manufactured for  
Nufarm Inc.  
11901 S. Austin Avenue  
Arling, IL 60803



Grow a better tomorrow

**FIRST AID**

<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>DO NOT induce vomiting unless told to do so by a poison control center or doctor.</li> <li>DO NOT give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**NOTE TO PHYSICIAN**

If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

For Chemical Spill, Leak, Fire, or Exposure,  
Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840



Net Contents  
**2.5 Gal.**  
(9.46 L)

**Nonrefillable Container**

15404000

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Harmful if absorbed through skin, swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing and breathing vapor. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### Personal Protective Equipment (PPE)

- All handlers must wear long-sleeved shirts, long pants, shoes and socks.
- Applicators using groundboom equipment with open cabs to treat cotton must wear long-sleeve shirts, long pants, shoes, and socks plus chemical-resistant gloves.
- Mixer/loaders supporting ground boom applications to corn, canola, soybean, cotton, citrus fruit, pome fruit, stone fruit, and olives must wear long-sleeve shirts, long pants, shoes, and socks plus chemical resistant gloves.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Applicators using ground boom equipment with open cabs to treat cotton must wear long-sleeve shirts, long pants, shoes, and socks plus chemical-resistant gloves.

Mixer/loaders supporting ground boom applications corn, canola, soybean, cotton, citrus fruit, pome fruit, stone fruit, and olives must wear long-sleeve shirts, long pants, shoes, and socks plus chemical-resistant gloves.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling Cheetah. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## ENVIRONMENTAL HAZARDS

DO NOT apply directly to water or to areas where surface water is present. DO NOT apply to intertidal areas below the mean high water mark. DO NOT contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate.

This pesticide is toxic to vascular plants and must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions, Cheetah may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, including no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur to minimize water run-off is recommended.

### PHYSICAL AND CHEMICAL HAZARDS

**DO NOT** mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

Combustible. **DO NOT** use or store near heat or open flame.

## DIRECTIONS FOR USE

**It is a violation of Federal law to use Cheetah in a manner inconsistent with its labeling.**

**DO NOT** use Cheetah until you have read the entire label.

**DO NOT** apply Cheetah in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**In the State of New York Only: Not For Use In Nassau and Suffolk Counties.**

## AGRICULTURAL USE REQUIREMENTS

Use Cheetah only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of Cheetah that are covered by the Worker Protection Standard.

### Restricted entry-interval (REI) 12 hours for all post-application activities

**Restricted entry-interval (REI) 12 hours for all post-application activities, with the following exception: Workers engaged in scouting or irrigation activities in corn, canola, and soybean is 4 days. The REI for workers to move irrigation piping is 7 days for all crops.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants;
- Chemical resistant gloves including barrier laminate, butyl rubber 14 mils, nitrile rubber >14 mils, neoprene rubber > 14 mils, polyvinyl chloride (PVC) > 14 mils, or Viton > 14 mils, and
- Chemical resistant footwear plus socks;
- Protective eyewear (goggles, face shield or safety glasses).

## IMPORTANT CROP SAFETY INFORMATION READ BEFORE USING CHEETAH

Cheetah may be applied as a burndown treatment **prior to planting or prior to crop emergence** of any canola, sweet corn<sup>[1]</sup>, corn, cotton, olive, soybean or sugar beet.

### POST EMERGENT TREATMENTS

Post emergence row crop applications of Cheetah may be made only to crops resistant to glufosinate. Tank mixtures of Cheetah with other products may impact crop resistance and increase risk of crop injury. The basis of selectivity of Cheetah herbicide in glufosinate-resistant crops is the presence of a gene that makes crops not sensitive to glufosinate. Crops not containing the glufosinate-resistant gene will be sensitive to Cheetah and severe crop injury and/or death may occur. **DO NOT** allow spray to contact foliage or green tissue of desirable vegetation other than crops containing the glufosinate-resistant trait.

Post emergent applications of Cheetah may be applied to crops not containing the glufosinate-resistant gene using a hooded sprayer.

### TREE, NUT, VINE AND BERRY TREATMENTS

Applications to trees, vines and berries must avoid contact of Cheetah solution, spray drift or mist with green bark, stems or foliage as injury may occur to trees, berries and vines. Only trunks with callused mature brown bark must be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of Cheetah with parts of trees, berries or vines other than mature brown bark can result in serious damage.

[<sup>1</sup> – Not for use in California]

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or greenhouses.

**DO NOT** enter or allow others to enter treated areas until sprays have dried.

## PRODUCT INFORMATION

Cheetah is a water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds in a variety of crops.

Cheetah is registered for use:

- as a burndown treatment prior to planting or prior to emergence of canola, corn, cotton, sweet corn, olive, soybean and sugar beets
- post emergence weed control herbicide to be applied on crops containing the glufosinate-resistant gene, including canola, soybean, corn, sweet corn and cotton
- post emergence weed control in cotton when applied with a hooded sprayer in-crop
- post emergence weed control in listed tree, olives, vine, and berry crops
- applied for potato vine desiccation.

**Many glufosinate-resistant seed trade names are available. Contact the seed manufacturer or seed distributor to determine if the seed variety is designated and supported as containing the glufosinate-resistant trait.**

It is important to always follow a responsible integrated weed management program.

Contact your local agronomic advisor for more specific information on integrated weed management in your area.

Cheetah is only foliar-active with little or no activity in soil. Weeds that emerge after application will not be controlled.

Apply Cheetah to actively growing weeds as described in the **WEED CONTROL FOR ROW CROPS** section to get maximum weed control. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Necrosis of leaves and young shoots occur within 2 to 4 days after application under good growing conditions.

- Cheetah is rainfast 4 hours after application to most weed species; therefore, rainfall within 4 hours may necessitate retreatment or may result in reduced weed control.
- Make applications between dawn and 2 hours before sunset to avoid the possibility of reduced lambsquarters and velvetleaf control.
- Consult your local Cooperative Extension Service or Nufarm, Inc. representative for guidelines on the optimum application timing for Cheetah in your region.
- Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to environmental conditions including drought, cool temperatures, or extended periods of cloudiness.
- To maximize weed control, **DO NOT** cultivate from 5 days before an application to 7 days after an application.

## ROTATIONAL CROP RESTRICTIONS\*

Rotational crop planting intervals following application of Cheetah are listed below. Failure to comply with these restrictions may result in illegal residues in rotated crops.

Rotational Crop	Plant Back Interval (Minimum Rotational Crop Planting Interval from Last Application)
Canola, Corn, Cotton, Soybeans, Sweet Corn, and Sugar beets	May be planted at any time
Root and Tuber Vegetables, Leafy Vegetables, Brassica Leafy Vegetables and Small Grains (Barley, Buckwheat, Oats, Rye, Teosinte, Triticale, and Wheat)	70 days
All other crops	180 days

\*See **Application Directions for Potato Vine Desiccation** for Rotational Crop Restrictions specifically after application of Cheetah to potatoes.

## WEED RESISTANCE MANAGEMENT

For resistance management, Cheetah contains a Group 10 herbicide –Glufosinate-ammonium. Any weed population may contain or develop plants naturally resistant to Cheetah and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Cheetah or other Group 10 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage ( or other mechanical control methods), cultural ( e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report non-performance or suspected resistance, contact Nufarm at 1-800-345-3330

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. **DO NOT** assume that each listed weed is being controlled by these mechanisms of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredient in this product.

Suspected herbicide-resistant weeds may be identified by these indicators:

- \* Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- \* A spreading patch of non-controlled plants of a particular weed species; and
- \* Surviving plants mixed with controlled individuals of the same species.

#### INTEGRATED PEST MANAGEMENT

Nufarm advises the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program which can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product needs to be based on IPM principles and practices including field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop or site systems in your area.

### WEED CONTROL FOR ROW CROPS

Rates in fluid ounce of formulated product per acre for the control of weeds as shown in the weed control tables. In weed populations with mixed species, apply at a rate needed for the species targeting less than three inch weeds.

**Table 1. Broadleaf Weeds Controlled  
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22.0 Fl Oz/A (0.40 lb ai/A)	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression	C=Control S = Suppression
Amaranth, Palmer	<i>Amaranthus palmeri</i>	Not Advised	C
Anoda, spurred	<i>Anoda cristata</i>	C	C
Beggarweed, Florida	<i>Desmodium tortuosum</i>	C	C
Black medic	<i>Medicago lupulina</i> L.	C	C
Blueweed, Texas	<i>Helianthus ciliaris</i> DC.	C	C
Buckwheat, wild	<i>Polygonum convolvulus</i>	C	C
Buffalobur	<i>Solanum cornutum</i>	C	C
Burcucumber	<i>Sicyos angulatus</i>	C	C
Canola, volunteer <sup>2</sup>	<i>Brassica</i> spp.	C <sup>2</sup>	C <sup>2</sup>
Catchweed bedstraw (cleavers)	<i>Galium aparine</i> L.	C	C
Carpetweed	<i>Mollugo verticillata</i>	C	C
Chickweed, common	<i>Stellaria media</i>	C	C
Cocklebur, common	<i>Xanthium strumarium</i>	C	C
Copperleaf, hophornbeam	<i>Acalypha ostryaefolia</i>	C	C
Cotton, volunteer <sup>2</sup>	<i>Gossypium</i> spp.	C <sup>2</sup>	C <sup>2</sup>
Croton, tropic	<i>Croton glandulosus</i>	C	C
Croton, woolly	<i>Croton capitatus</i>	C	C
Eclipta	<i>Eclipta alba</i>	C	C
Devil's claw	<i>Proboscidea Louisiana</i>	C	C
Fleabane, annual	<i>Erigeron annuus</i>	C	C
Galinsoga, hairy	<i>Galinsoga ciliata</i>	C	C
Galinsoga, small flower	<i>Galinsoga parviflora</i>	C	C

(continued)

**Table 1. Broadleaf Weeds Controlled (continued)**  
**(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22.0 Fl Oz/A (0.40 lb ai/A)	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression	C=Control S = Suppression
Groundcherry, cutleaf	<i>Physalis angulate</i>	C	C
Geranium, cutleaf	<i>Geranium dissectum L.</i>	C	C
Hempnettle	<i>Galeopsis spp.</i>	C	C
Horsenettle, Carolina <sup>3</sup>	<i>Solanum carolinense</i>	C <sup>3</sup>	C <sup>3</sup>
Jimsonweed	<i>Datura stramonium</i>	C	C
Knotweed	<i>Polygonum spec.</i>	C	C
Kochia	<i>Kochia scoparia</i>	C	C
Ladysthumb	<i>Polygonum persicaria</i>	C	C
Lambsquarters, common	<i>Chenopodium album</i>	C	C
Mallow, common	<i>Malva spec.</i>	C	C
Mallow, Venice	<i>Hibiscus trionum</i>	C	C
Marestail <sup>3</sup>	<i>Conyza Canadensis</i>	S	C
Marsh-elder, annual	<i>Iva annua</i>	C	C
Morningglory, entireleaf	<i>Ipomoea hederacea var. intergriuscula</i>	C	C
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	C	C
Morningglory, pitte	<i>Ipomoea lacunose</i>	C	C
Morningglory, sharppod	<i>Ipomoea cordatotriloba</i>	C	C
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>	C	C
Morningglory, tall	<i>Lpomoea purpurea</i>	C	C
Mustard, wild	<i>Sinapis arvensis</i>	C	C
Nightshade, black	<i>Solanum nigrum</i>	C	C
Nightshade, eastern black	<i>Solanum ptycanthum</i>	C	C
Nightshade, hairy	<i>Solanum sarrachoides</i>	C	C
Pennycress	<i>Thlaspi arvense</i>	C	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C	C
Pigweed, prostrate	<i>Amaranthus blitoides</i>	C	C
Pigweed, spiny	<i>Amaranthus spinosus</i>	C	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C	C
Pigweed, tumble	<i>Amaranthus albus</i>	C	C
Puncturevine	<i>Tribulus terrestris</i>	C	C
Purslane, common	<i>Portulaca oleracea</i>	C	C
Pusley, Florida	<i>Richardia scabra</i>	S	C
Ragweed, common	<i>Ambrosia artemisiifolia</i>	C	C
Ragweed, giant	<i>Ambrosia trifida</i>	C	C
Senna coffee	<i>Cassia occidentalis</i>	C	C
Sesbania, hemp	<i>Sesbania herbacea</i>	C	C
Shepherd's-Purse	<i>Capsella bursa-pastoris</i>	C	C
Sicklepod (java bean)	<i>Senna obtusifolia</i>	C	C

(continued)

**Table 1. Broadleaf Weeds Controlled (continued)**  
**(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22.0 Fl Oz/A (0.40 lb ai/A)	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression	C=Control S = Suppression
Sida, prickly	<i>Sida spinosa L.</i>	C	C
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	C	C
Smell melon	<i>Cucumis melo L. var. Dudaim</i>	C	C
Sowthistle, annual	<i>Sonchus oleraceus L.</i>	C	C
Soybeans, volunteer <sup>2</sup>	<i>Glycine max</i>	C <sup>2</sup>	C <sup>2</sup>
Spurge, prostrate	<i>Euphorbia humifusa</i>	C	C
Spurge, spotted	<i>Euphorbia maculate L.</i>	C	C
Starbur, bristly	<i>Acanthospermum hispidum</i>	C	C
Sunflower, common	<i>Helianthus annuus</i>	C	C
Sunflower, prairie	<i>Corythucha pura</i>	C	C
Sunflower, volunteer	<i>Girassol</i>	C	C
Thistle, Russian <sup>3</sup>	<i>Salsola kali</i>	S <sup>3</sup>	C <sup>3</sup>
Velvetleaf	<i>Abutilon theophrasti</i>	C	C
Waterhemp, common	<i>Amaranthus rudis</i>	Not Advised	C
Waterhemp, tall	<i>Amaranthus tuberculatos</i>	Not Advised	C

<sup>1</sup> Maximum rate on canola, field corn, sweet corn and soybean in California is 36 fl oz/A (0.66 lb ai/A).

<sup>2</sup> Volunteer glufosinate-resistant crops from the previous year will not be controlled.

<sup>3</sup> May require sequential applications for control.

<sup>4</sup> For optimum control apply Cheetah on 6" marestalk

**Table 2. Grass Weeds Controlled**  
**(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22.0 Fl Oz/A (0.40 lb ai/A)	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression	C=Control S = Suppression
Barley, volunteer <sup>3</sup>		C <sup>3</sup>	C <sup>3</sup>
Barnyardgrass	<i>Echinochloa spec.</i>	C	C
Bluegrass, annual	<i>Poa annua L.</i>	C	C
Corn, volunteer <sup>2</sup>	<i>Zea mays L.</i>	C <sup>2</sup>	C <sup>2</sup>
Crabgrass, large <sup>4</sup>	<i>Digitaria sanguinalis</i>	C <sup>4</sup>	C <sup>4</sup>
Crabgrass, smooth <sup>4</sup>	<i>Digitaria ischaemum</i>	C <sup>4</sup>	C <sup>4</sup>
Cupgrass, woolly	<i>Eriochloa villosa</i>	C	C
Foxtail, bristly	<i>Setaria verticillata</i>	C	C
Foxtail, giant	<i>Setaria faberi</i>	C	C
Foxtail, green	<i>Setaria viridis</i>	C	C
Foxtail, robust purple	<i>Setaria viridis</i>	C	C
Foxtail, yellow <sup>4</sup>	<i>Pennisetum glaucum</i>	C <sup>4</sup>	C <sup>4</sup>
Goosegrass <sup>3</sup>	<i>Eleusine indica</i>	C <sup>3</sup>	C <sup>3</sup>
Johnsongrass, seedling	<i>Sorghum halepense</i>	C	C

(continued)

**Table 2. Grass Weeds Controlled (continued)**  
**(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

Common Name	Scientific Name	22.0 Fl Oz/A (0.40 lb ai/A)	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression	C=Control S = Suppression
Junglerice	<i>Echinochloa coloum</i>	C	C
Millet, wild-proso	<i>Panicum millaceum L.</i>	C	C
Millet, proso volunteer	<i>Milium vernale</i>	C	C
Oat, wild <sup>4</sup>	<i>Avena fatua</i>	C <sup>4</sup>	C <sup>4</sup>
Panicum, fall	<i>Panicum dichotomiflorum</i>	C	C
Panicum, Texas	<i>Panicum texanum</i>	C	C
Rice, red	<i>Oryza sativa L.</i>	C	C
Sandbur, field <sup>4</sup>	<i>Cenchrus pauciflorus</i>	S <sup>4</sup>	C <sup>4</sup>
Shattercane	<i>Sorghum vulgare PERS.</i>	C	C
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	C	C
Sprangletop	<i>Leptochloa spec.</i>	C	C
Sorghum, volunteer	<i>Sorghum spp.</i>	C	C
Stinkgrass	<i>Eragrostis ciliaris</i>	C	C
Wheat, volunteer <sup>3,4</sup>	<i>Triticum spec.</i>	C <sup>3,4</sup>	C <sup>3,4</sup>
Witchgrass	<i>Panicum virgatum L.</i>	C	C

<sup>1</sup> Maximum rate on canola, field corn, sweet corn and soybean in California is 36 fl oz/A (0.66 lb ai/A).

<sup>2</sup> Volunteer glufosinate-resistant crops from the previous year will not be controlled. A timely cultivation 7 to 10 days after an application and/or retreatment 10-21 days after the first application is needed for controlling dense clumps of volunteer corn.

<sup>3</sup> May require sequential applications for control.

<sup>4</sup> For best control of yellow foxtail, field sandbur, crabgrass, and wild oats, treat prior to tiller initiation.

**Table 3. Biennial and Perennial Weeds Controlled**  
**(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)**

**For control of the biennial and perennial weeds listed below, tank mix partners or sequential applications of Cheetah are specified by crop (see crop sections)**

Common Name	Scientific Name	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression
Alfalfa	<i>Medicago sativa L.</i>	C
Bermudagrass	<i>Cynodon dactylon</i>	C
Bindweed, field	<i>Convolvulus arvensis L.</i>	C
Bindweed, hedge	<i>Calystegia sepium</i>	C
Bluegrass, Kentucky	<i>Poa pratensis L.</i>	C
Blueweed, Texas	<i>Helianthus ciliaris DC.</i>	C
Bromegrass, smooth	<i>Bromus inermis</i>	C
Burdock	<i>Arctium spp.</i>	C
Bursage, woollyleaf	<i>Ambrosia grayi</i>	C
Chickweed, Mouse-ear	<i>Cerastium vulgatum L.</i>	C
Clover, red	<i>Trifolium pretense L.</i>	C
Dandelion	<i>Taraxacum officinale</i>	C

(continued)



**Table 3. Biennial and Perennial Weeds Controlled (continued)**  
(including Glyphosate-, Triazine-, PPO-, ALS-, HPPD-, and Auxin-Resistant Biotypes)

**For control of the biennial and perennial weeds listed below, tank mix partners or sequential applications of Cheetah are specified by crop (see crop sections)**

Common Name	Scientific Name	29.0 – 43.0 Fl Oz/A <sup>1</sup> (0.53-0.79 lb ai/A)
		C=Control S = Suppression
Dock, smooth	<i>Rumex spec.</i>	C
Dogbane, hemp	<i>Apocynum cannabinum</i>	S
Goldenrod, gray	<i>Solidago nemoralis</i>	S
Johnsongrass, rhizome	<i>Sorghum halepense</i>	C
Milkweed, common	<i>Asclepias syriaca</i>	S
Milkweed, honeyvine	<i>Ampelamus albidus</i>	S
Muhly, wirestem	<i>Muhlenbergia frondosa</i>	S
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>	C
Nutsedge, purple	<i>Cyperus rotundus</i>	S
Nutsedge, yellow	<i>Cyperus ferax</i>	S
Orchardgrass	<i>Dactylis glomerata L.</i>	C
Poinsettia, wild	<i>Euphorbia heterophylla L.</i>	C
Pokeweed	<i>Phytolaccaceae</i>	C
Quackgrass	<i>Agropyron repens</i>	S
Sowthistle, perennial	<i>Sonchus arvensis L.</i>	C
Thistle, bull	<i>Cirsium vulgare</i>	C
Thistle, Canada	<i>Cirsium arvense</i>	C
Timothy	<i>Phleum pretense L.</i>	S
Wormwood, biennial	<i>Artemisia biennis</i>	S

<sup>1</sup> Maximum rate on canola, field corn, sweet corn and soybean in California is 36 fl oz/A (0.66 lb ai/A).

## APPLICATION AND MIXING PROCEDURES

**Uniform, thorough spray coverage is important to achieve consistent weed control.**

### Ground Application:

- Refer to the **Rate Tables** for proper application rates.
- Apply early, when weeds are small.
- To avoid drift and insure consistent weed control, apply Cheetah with the spray boom as low as possible while maintaining a uniform spray pattern.
- Apply Cheetah broadcast in a minimum of 15.0 gallons of water per acre. Increase to 20 gallons of water per acre if dense weed canopy exists.
- Apply at ground speed of less than 15 mph to attain adequate coverage.
- See the **SPRAY DRIFT MANAGEMENT** section of this label for additional information on proper application of Cheetah.

### Aerial Application:

- Refer to the **Rate Tables** for proper application rates.
- Apply early, when weeds are small.
- Apply Cheetah by air in a minimum of 10.0 gallons of water per acre.
- See the **SPRAY DRIFT MANAGEMENT** section of this label for additional information on proper application of Cheetah.

### Application and Mixing Restrictions:

- **DO NOT** use flood jet nozzles, controlled droplet application equipment, or air-assisted spray equipment.

### Compatibility Testing:

If Cheetah is to be mixed with pesticide products not listed on this label, test the compatibility of the intended tank mixture prior to mixing the products in the spray tank. The following procedure assumes a spray volume of 25.0 gallons per acre. For other spray volumes, adjust the amount of the water used accordingly. Check compatibility as follows:

1. Place 1.0 pint of water from the source that will be used to prepare the spray solution in a clear 1.0 quart jar.
2. For each pound of a dry tank mix partner to be applied per acre, add 1.5 teaspoons to the jar.
3. For each 16.0 fluid ounces of a liquid tank mix partner to be applied per acre, add 0.5 teaspoon to the jar.
4. For each 16.0 fluid ounces of Cheetah to be applied per acre, add 0.5 teaspoon to the jar.
5. After adding all the ingredients, place a lid on the jar and tighten. Invert 10 times to mix.
6. Let the mixture stand for 15 minutes, and evaluate the solution for uniformity and stability. Look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. If the tank mix partners are not compatible, **DO NOT** use the mixture in a spray tank.
7. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **STORAGE AND DISPOSAL** section of this label.

## MIXING INSTRUCTIONS

**Tank Mix Instructions:** Cheetah may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Cheetah cannot be mixed with any product containing a label prohibition against such mixing. Refer to the specific crop section for rates and other restrictions.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Cheetah must be applied with properly calibrated and clean equipment. Cheetah is formulated to mix readily in water.

Prior to adding Cheetah to the spray tank, ensure that the spray tank is thoroughly clean, particularly if an herbicide with the potential to injure crops was previously used (see **Cleaning Instructions**).

Mix Cheetah with water to make a finished spray solution as follows:

1. Properly calibrated and clean equipment
2. Fill the spray tank half full with water.
3. Start agitation.
4. If mixing with a flowable/wettable powder tank mix partner, prepare a slurry of the proper amount of the product in a small amount of water. Add the slurry to the spray tank.
5. Add the appropriate amount of ammonium sulfate (AMS) to the spray tank.
6. If mixing with a liquid tank mix partner, add the liquid mix partner next.
7. Complete filling the spray tank with water before adding Cheetah, as foaming may occur.
8. Add the proper amount of Cheetah and continue agitation.
9. If foaming occurs, use a silicone-based antifoam agent.

Ensure that all spray system lines including pipes, booms, etc. have the correct concentration of spray solution by flushing out the spray system lines before starting the crop application.

If tank mix partners specified on this label are added, maintain good agitation at all times until contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep bypass line on or near bottom of tank to minimize foaming. Screen size in nozzles or line strainers must be 50 mesh or larger.

### Cleaning Instructions:

Before using Cheetah, thoroughly clean bulk storage tank, refillable tank, nurse tanks, spray tank, lines, and filter, particularly if a herbicide with the potential to injure crops was previously used. Thoroughly rinse equipment using a commercial tank cleaner and as instructed on the prior herbicide label.

After using Cheetah, triple rinse the spray equipment and clean with a commercial tank cleaner before using the equipment for crops not containing glufosinate-resistant gene. Make sure any rinsate or foam is thoroughly removed from spray tank and boom. Rinsate may be disposed following the pesticide disposal directions on this label.

## SPRAY DRIFT MANAGEMENT

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For all non-aerial applications, wind speed must be measured adjacent to the application site, on the upwind side, immediately prior to application.

### MANDATORY SPRAY DRIFT MITIGATION

- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, applicators must use ½ swath displacement upwind at the downwind edge of the field.
- DO NOT apply when wind speeds exceed 10 miles per hour at the application site.
- DO NOT apply during temperature inversions.
- For aerial applications, DO NOT release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.
- For ground applications and aerial applications, select nozzle and pressure that deliver medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1.
- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but DO NOT exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- For non-crop vegetation management ground applications, apply with the nozzle height no more than 4 feet above the ground or target vegetation, unless necessitated by the application equipment. Examples would include roadside, railroad, utility rights of way, forestry and other industrial vegetation management applications where safety or natural barriers obstruct application.

### SPRAY DRIFT ADVISORIES

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

#### POLLINATOR ADVISORY

This product contains an herbicide. Follow all label directions and precautions to minimize potential off-target exposure in order to prevent effects to non-target plants adjacent to the treated site which may serve as habitat or forage for pollinators.

#### IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

#### Controlling Droplet Size – Ground Boom

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

#### Controlling Droplet Size – Aircraft

- Number of Nozzles - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations. **AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**
- Nozzle Type - Solid stream nozzles (including disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- Boom Length - Longer booms increase drift potential. Therefore a shorter boom length is recommended.
- Application Height - Application more than 10 ft. above the canopy increases the potential for spray drift.

#### BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

#### Drift Reduction Technology (DRT)

The EPA Drift Reduction Technology (DRT) Program was developed to encourage the manufacturer, marketing, and use of spray technologies scientifically verified to significantly reduce pesticide drift. The use of DRTs should result in significantly less pesticide from spray applications drifting and being deposited in areas not targeted by those applications, compared to spray technologies that DO NOT meet the minimum DRT standard. EPA-verified drift reduction technologies (DRTs) and their ratings will be added to the following webpage when they become available: <https://www.epa.gov/reducing-pesticide-drift/epa-verified-and-rated-drift-reduction-technologies>

#### WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.**

Note: Local terrain can influence wind patterns. Every applicator needs to be familiar with local wind patterns and how they affect spray drift.

## TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

## TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

## APPLICATION DIRECTIONS FOR BURNDOWN USE

Cheetah may be applied as a **burndown treatment prior to planting or prior to emergence** of any variety of canola, corn, sweet corn, cotton, soybean or sugar beet.

### Application Timing:

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section. Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. For optimum results on lambsquarters, Palmer amaranth and velvetleaf make applications between dawn and 2 hours before sunset. Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

### Application Rates:

Apply 29.0 – 43.0 fluid ounces (1.32 lbs ai/A) per acre of Cheetah depending on crop, weed species and intention of post application use. Please see application charts below.

### Restrictions (also see specific crop use directions):

- In **cotton**, if environmental conditions prevent timely applications, a single application may be made of up to 43.0 fluid ounces (0.79 lbs ai/A) per acre of Cheetah. **If more than 29.0 fluid ounces (0.53 lbs ai/A) per acre are used in any single application, the annual total may not exceed 72.0 fluid ounces (1.32 lbs ai/A) per acre, including all application timings.**
- In **canola, corn (sweet and field) and soybean**, if environmental conditions prevent timely applications, a single application may be made of up to 43.0\* fluid ounces (0.79 lbs ai/A) per acre of Cheetah. The year total may not exceed 43.0\* fluid ounces (0.79 lbs ai/A) per acre, including all application timings, for non-Glufosinate resistant crops. \*Maximum rate in California is 36 fl oz/A (0.66 lb ai/A).

### Adjuvant:

Ammonium sulfate (AMS) may be used at 1.5 to 3 pounds per acre. Adjuvant rates are dependent on tank mix partners, temperatures, environmental conditions and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (low relative humidity) or hard water.

### Surfactants / Crop Oils:

The use of surfactants may be included. Please refer to the surfactant label for more detailed information.

Table 4. APPLICATION DIRECTIONS FOR NON-GLUFOSINATE RESISTANT CROPS			
Crop	Burndown	In Season Applications	Annual Max
Canola, Soybean, Sweet Corn, Field Corn	29 – 43* fl oz/A (0.53 – 0.79 lbs ai/A)	None	43* fl oz/A (0.79 lbs ai/A)
Sugar beets	29 - 36 fl oz/A (0.53 – 0.66 lbs ai/A)	None	36 fl oz/A (0.66 lbs ai/A)
Cotton Use Pattern 1	29 fl oz/A (0.53 lbs ai/A)	2 applications at 29 fl oz/A** (0.53 lbs ai/A) Make second application 10-14 days after the first application.	87 fl oz/A (1.59 lbs ai/A)
Cotton Use Pattern 2	30-43 fl oz/A (0.55 – 0.79 lbs ai/A)	1 application at 29 fl oz/A** (0.53 lbs ai/A)	72 fl oz/A (1.32 lbs ai/A)

\* Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A).

\*\* Cotton containing the glufosinate-resistant gene OR with hooded sprayer for all varieties (see **COTTON** use directions).

**Table 5. APPLICATION DIRECTIONS FOR CROPS CONTAINING THE GLUFOSINATE RESISTANT GENE TRAIT**

<b>GLUFOSINATE RESISTANT Crop</b>	<b>Burndown</b>	<b>In Season Applications of Crops Containing the GLUFOSINATE RESISTANT GENE</b>	<b>Annual Max</b>
Soybean	29 – 43* fl oz/A (0.53 – 0.79 lbs ai/A)	1 to 2 applications at 29 – 43* fl oz/A (0.53 – 0.79 lbs ai/A) For soybeans, make second application at least 5 days after the first application. For field corn, make second application at least 7 days after first application.	87* fl oz/A (1.59 lbs ai/A)
Field Corn	29 – 43** fl oz/A (0.53 – 0.79 lbs ai/A)	1 to 2 applications at 29 – 43** fl oz/A (0.53 – 0.79 lbs ai/A) Make second application at least 7 days after the first application.	87** fl oz/A (1.59 lbs ai/A)
Sweet Corn***	22 fl oz/A (0.40 lbs ai/A)	1 to 2 applications at 22 fl oz/A (0.4 lbs ai/A) Make second application at least 7 days after the first application.	44 fl oz/A (0.8 lbs ai/A)
Canola	29 – 43* fl oz/A (0.53 – 0.79 lbs ai/A)	1 to 2 applications at 29 fl oz/A (0.53 lbs ai/A) Make second application at least 10 days after the first application	87* fl oz/A (1.59 lbs ai/A)
Cotton Use Pattern 1	29 fl oz/A (0.53 lbs ai/A)	1 to 2 applications at 29 fl oz/A**** (0.53 lbs ai/A) Make second application 10-14 days after the first application.	87 fl oz/A (1.59 lbs ai/A)
Cotton Use Pattern 2	30 - 43 fl oz/A (0.55 – 0.79 lbs ai/A)	1 application at 29 fl oz/A**** (0.53 lbs ai/A)	72 fl oz/A (1.32 lbs ai/A)

\* Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A) with annual maximum of 72 fl oz/A (1.32 lbs ai/A).

\*\* Maximum rate in California is 22 fl oz/A (0.40 lbs ai/A) with annual maximum of 44 fl oz/A (0.8 lbs ai/A).

\*\*\* Not for use in California

\*\*\*\* Cotton containing the glufosinate-resistant gene OR with hooded sprayer for all varieties (see **COTTON** use directions).

## **APPLICATION DIRECTIONS FOR USE ON CANOLA CONTAINING THE GLUFOSINATE-RESISTANT GENE**

Apply Cheetah only to canola containing the Glufosinate-resistant gene. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

### **Application Timing:**

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

Applications of Cheetah on canola containing the glufosinate-resistant gene may be made from the cotyledon stage up to the early bolt stage of the canola. Slight discoloration of the canola may be visible after application. This effect is temporary and will not influence crop growth, maturity, or yield.

Cheetah is a foliar-active material with little or no soil-residual activity.

Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

For best result:

- On lambsquarters, Palmer amaranth and velvetleaf control, make applications of Cheetah between dawn and 2 hours before sunset.
- Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah.

### **Application Rates:**

Apply Cheetah at 22.0 to 29.0 fl oz/A (0.4 to 0.53 lbs ai/A) per application, depending on weed species, size and density per weed chart.

If a second application is needed, make the second application in a minimum of 7 days after the first application.

The maximum annual rate of Cheetah on canola is 87.0 fl oz/A (1.59 lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

### **Application Rates with Tank Mix Partners:**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Apply Cheetah at 22.0 to 29.0 fl oz/A (0.4 to 0.53 lbs ai/A) per application, depending on weed species, size and density per weed chart. Tank mix partners advised to enhance grass control, including quizalofop p-ethyl, sethoxydim, and clethodim.

If a second application is needed, make the second application in a minimum of 7 days after the first application.

Tank mixes may aid in the performance of Cheetah. Please refer to weed chart tables for a listing of weed species controlled at this rate.

No additional surfactant is needed with any tank mix partner.

The tank mix partner must be used in accordance with the label limitations, restrictions and precautions.

**DO NOT** exceed any labeled dosage rates.

**DO NOT** mix Cheetah mix with any product containing a label prohibition against such mixing.

#### **Adjuvants:**

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (including temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (including low relative humidity) or hard water.

The use of an anti-foam agent is advised.

#### **Surfactants / Oils:**

The use of additional surfactants or crop oils in tank mixes with Cheetah may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

#### **Restrictions to the Directions For Use on Canola Containing the Glufosinate-Resistant Gene:**

- **DO NOT** exceed the maximum single application rate of 43 fl oz/A (0.79 lb ai/A).
- **DO NOT** apply more than 87.0 fl oz/A (1.59 lbs ai/A) of Cheetah per year.
- **DO NOT** use on canola containing the glufosinate-resistant gene in the states of Alabama, Delaware, Georgia, Kentucky, Maryland, New Jersey, North Carolina, South Carolina, Tennessee, Virginia and West Virginia.
- **DO NOT** apply more than 2 applications of Cheetah per year. Sequential applications must be at least 10 days apart.
- **DO NOT** apply Cheetah within 65 days of harvesting canola.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** apply Cheetah if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply Cheetah through any type of irrigation system.
- Refer to the **ROTATIONAL CROP RESTRICTIONS** section under the **PRODUCT INFORMATION** heading of this label for the appropriate rotational crop plant back intervals.

## **APPLICATION RATE AND TIMING FOR CANOLA CONTAINING GLUFOSINATE RESISTANT GENE SEED PROPAGATION**

### **Not for use in California**

Up to 3 applications of Cheetah at up to 29.0 fl oz/A (0.53 lbs ai/A) per application may be made to canola containing the Glufosinate-resistant gene for seed propagation. Applications may be made from the cotyledon stage up to the early bolting stage (e.g., BBCH 18-30, between just prior to stem elongation/bolting, 8 or more leaves and beginning of stem elongation, no internodes).

#### **Restrictions to the Directions for Canola Containing the glufosinate-resistant gene for Seed Propagation:**

- Maximum single application is 29.0 fl oz/A (0.53 lbs ai/A).
- **DO NOT** apply more than 3 applications.
- Sequential applications must be made more than 10 days apart.
- **DO NOT** apply more than 87.0 fl oz/A (1.59 lbs ai/A) of Cheetah per year.
- **DO NOT** apply Cheetah beyond the early bolting stage or within 65 days of harvesting canola seed.
- **DO NOT** use treated canola seed for food, feed or oil purposes.
- **DO NOT** apply Cheetah if canola shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply this product through any type of irrigation system.

## **APPLICATION DIRECTIONS FOR USE ON SWEET CORN CONTAINING THE GLUFOSINATE RESISTANT GENE**

### **Not for use in California.**

Apply Cheetah only to sweet corn containing the glufosinate-resistant gene. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

#### **Application Timing:**

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

Applications for Cheetah on sweet corn may be made from emergence until the V-6 stage of growth.

Cheetah is a foliar-active material with little or no soil-residual activity.

Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

For best result:

- On lambsquarters, Palmer amaranth and velvetleaf control, make applications of Cheetah between dawn and 2 hours before sunset.
- Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah.

#### **Application Rate:**

Apply Cheetah at 22.0 fl oz/A (0.4 lbs ai/A), depending on weed species, size and density per weed chart.

If a second application is needed, make the second application in a minimum of 7 days after the first application.

The maximum annual rate of Cheetah on sweet corn is 44.0 fl oz/A (0.8 lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

#### **Application Rates with Tank Mix Partners:**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Apply Cheetah at 22.0 fl oz/A (0.4 lbs ai/A) per application, depending on weed species, size and density per weed chart.

Advised tank mix partners, including atrazine, tembotrione, thiencazabone-methyl, and dicamba, (DGA salt).

If a second application is needed, make the second application in a minimum of 7 days after the first application.

Tank mixes may aid in the performance of Cheetah. Please refer to weed chart tables for a listing of weed species controlled at this rate.

No additional surfactant is needed with any tank mix partner.

The tank mix partner must be used in accordance with the label limitations, restrictions and precautions.

**DO NOT** exceed any labeled dosage rates.

**DO NOT** mix Cheetah mix with any product containing a label prohibition against such mixing.

#### **Adjuvants:**

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (including temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (including low relative humidity) or hard water.

The use of an anti-foam agent is advised.

#### **Surfactants / Oils:**

The use of additional surfactants or crop oils in tank mixes with Cheetah may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

#### **Restrictions to the Directions For Use on Sweet Corn Containing the Glufosinate-Resistant Gene:**

- **DO NOT** apply Cheetah within 50 days of harvesting sweet corn ears and within 55 days of harvesting stover.
- **DO NOT** apply more than 44.0 fl oz/A (0.8 lbs ai/A) of Cheetah on sweet corn per year.
- **DO NOT** apply more than 2 applications of Cheetah to sweet corn per year. Sequential applications must be at least 7 days apart.
- **DO NOT** exceed the the maximum single application rate of 22 fl oz/A (0.40 lb ai/A).
- If Cheetah was used in a burndown application, no post emergence applications may be applied to the crop.
- **DO NOT** use nitrogen solutions as spray carriers.
- **DO NOT** apply Cheetah if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply Cheetah through any type of irrigation system.

Refer to the **ROTATIONAL CROP RESTRICTIONS** section under the **PRODUCT INFORMATION** heading of this label for the appropriate rotational crop plant back intervals.

See **APPLICATION DIRECTIONS FOR USE ON FIELD CORN AND SILAGE CORN for APPLICATION METHODS, MIXING INSTRUCTIONS, and WEED CONTROL TABLES.**

## **APPLICATION DIRECTIONS FOR USE ON FIELD CORN AND SILAGE CORN CONTAINING THE GLUFOSINATE RESISTANT GENE**

Apply Cheetah only to corn containing the glufosinate-resistant gene. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

#### **Application Timing:**

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

Applications for Cheetah on corn may be made from emergence until the V-6 stage of growth.

Cheetah is a foliar-active material with little or no soil-residual activity.

Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

For best result:

- On lambsquarters, Palmer amaranth and velvetleaf control, make applications of Cheetah between dawn and 2 hours before sunset.
- Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah.

#### **Application Rate:**

Apply Cheetah at 29 – 43\* fl oz/A (0.53 – 0.79 lbs ai/A), depending on weed species, size and density per weed chart. \*Maximum rate in California is 22 fl oz/A (0.4 lb ai/A).

If a second application is needed, make the second application at up to 29 fl oz/A (0.53 lbs ai/A) with a minimum of 7 days after the first application. \*Maximum rate in California is 22 fl oz/A (0.4 lb ai/A).

The maximum annual rate of Cheetah on field corn and silage corn is 87.0\* fl oz/A (1.59 lbs ai/A). \*Maximum annual rate in California is 44 fl oz/A (0.8 lbs ai/A)

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

#### **Application Rates with Tank Mix Partners:**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Apply Cheetah at 29.0 – 43.0\* fl oz/A (0.53 – 0.79 lbs ai/A), depending on weed species, size and density per weed chart. \* Maximum rate in California is 22 fl oz/A (0.4 lb ai/A).

Advised tank mix partners, including atrazine, tembotrione, thien carbazono-methyl, and dicamba, DGA salt.

If a second application is needed, make the second application in a minimum of 7 days after the first application.

Tank mixes may aid in the performance of Cheetah. Please refer to weed chart tables for a listing of weed species controlled at this rate. No additional surfactant is needed with any tank mix partner.

The tank mix partner must be used in accordance with the label limitations, restrictions and precautions.

**DO NOT** exceed any labeled dosage rates.

**DO NOT** mix Cheetah mix with any product containing a label prohibition against such mixing.

#### **Adjuvants:**

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (including temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (including low relative humidity) or hard water.

The use of an anti-foam agent is advised.

#### **Surfactants / Oils:**

The use of additional surfactants or crop oils in tank mixes with Cheetah may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

#### **Restrictions to the Directions For Use on Field Corn and Corn Silage Containing the Glufosinate-Resistant Gene:**

• **DO NOT** apply Cheetah within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder.

• **DO NOT** apply more than 2 applications per year. Sequential applications must be at least 10 days apart.

• **DO NOT** apply more than 87.0\* fl oz/A (1.59 lbs ai/A) of Cheetah on corn per year. \* Maximum annual rate in California is 44 fl oz/A (0.8 lbs ai/A).

• **DO NOT** exceed the maximum single application rate of 43 fl oz/A (0.79 lb ai/A).

• **DO NOT** use nitrogen solutions as spray carriers.

• **DO NOT** apply Cheetah if corn shows injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.)

• **DO NOT** apply Cheetah through any type of irrigation system.

Refer to the **ROTATIONAL CROP RESTRICTIONS** section under the **PRODUCT INFORMATION** heading of this label for the appropriate rotational crop plant back intervals.

## **APPLICATION DIRECTIONS FOR USE ON COTTON CONTAINING THE GLUFOSINATE RESISTANT GENE**

Uniform, thorough spray coverage is necessary to achieve consistent weed control. Cheetah may be applied as a broadcast, over-the-top, post-emergence spray or as a directed spray only to cotton containing the glufosinate-resistant gene.

#### **Application Timing:**

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

Cheetah is a foliar-active material with little or no soil-residual activity.

Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.



For best result:

- On lambsquarters, Palmer amaranth and velvetleaf control, make applications of Cheetah between dawn and 2 hours before sunset.
- Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah.

Apply Cheetah to cotton from emergence up to the early bloom stage at 29.0 fl oz/A (0.53 lbs ai/A). If environmental conditions prevent a timely herbicide application, a single application of up to 43.0 fl oz/A (0.79 lbs ai/A) of Cheetah may be made to cotton. If more than 29.0 fl oz/A (0.53 lbs ai/A) are used in any single application, the annual total may not exceed 72.0 fl oz/A (1.32 lbs ai/A), including all application timings. See **Restrictions to the Directions for use on Cotton Containing the Glufosinate-Resistant Gene** below for additional information.

#### Application Rates:

##### Option 1 3 post applications

Apply 29 fl oz/A (0.53 lbs ai/A) per application depending on weed species, size and density per weed chart. If required a second application of 29 fl oz/A (0.53 lbs ai/A) can be applied, followed by a third application of 29 fl oz/A (0.53 lbs ai/A). The sequential applications must be made 10-14 days after each other. The maximum annual rate of Cheetah on cotton is 87.0 fl oz/A (1.59 lbs ai/A). Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

##### Option 2 2 post applications

Apply 32 - 43 fl oz/A (0.59 – 0.79 lbs ai/A) per application depending on weed species, size and density per weed chart. If required a second application of 29 fl oz/A (0.53 lbs ai/A) can be applied. The sequential applications must be made 10-14 days after each other. The maximum annual rate of Cheetah on cotton is 72 fl oz/A (1.32 lbs ai/A). Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

Use Pattern	1 <sup>st</sup> Application	2 <sup>nd</sup> Application Minimum 10 Days Up to 14 Days After 1 <sup>st</sup> Application	3 <sup>rd</sup> Application Minimum 10 Days Up to 14 Days After 2 <sup>nd</sup> Application	Annual Maximum
Option 1	29 fl oz/A (0.53 lbs ai/A)	29 fl oz/A (0.53 lbs ai/A)	29 fl oz/A (0.53 lbs ai/A)	87 fl oz/A (1.59 lbs ai/A)
Option 2	32-43 fl oz/A (0.59 – 0.79 lbs ai/A)	29 fl oz/A (0.53 lbs ai/A)	None	72 fl oz/A (1.32 lbs ai/A)

#### Tank Mix on Cotton Containing the Glufosinate-Resistant Gene:

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Certain herbicide tank mixes may aid in the performance of Cheetah. Cheetah may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Cheetah cannot be mixed with any product containing a label prohibition against such mixing.

#### Adjuvants:

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (including temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (including low relative humidity) or hard water.

The use of an anti-foam agent is advised.

#### Surfactants / Oils:

The use of additional surfactants or crop oils in tank mixes with Cheetah may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

#### Restrictions to the Directions For Use on Cotton Containing the Glufosinate-resistant gene:

- **DO NOT** apply Cheetah to cotton **containing** the Glufosinate-resistant gene in Florida, South of Tampa (Florida Route 60), or in Hawaii, except for test plots or breeding nurseries.
- **DO NOT** apply Cheetah within 70 days prior to cotton harvest.
- Up to 3 applications of Cheetah may be made to cotton per year at a maximum application rate of 29.0 fl oz/A (0.53 lbs ai/A). **DO NOT** apply more than 87.0 fluid ounces (including all application timings) to cotton (1.59 lbs ai/A) per year under this application scenario. Sequential applications must be at least 10 days apart.

- If environmental conditions prevent timely applications resulting in large weeds or heavy infestations, a single application of Cheetah at up to 43.0 fl oz/A (0.79 lb ai/A) may be made to cotton. **DO NOT** apply more than 43.0 fluid ounces (0.79 lb ai/A) of Cheetah in a single application under this use scenario. If a single application greater than 29.0 fluid ounces (0.53 lb ai/A) is made, a subsequent application not to exceed 29.0 fluid ounces (0.53 lb ai/A) may be made to cotton. The annual total use rate under this scenario may not exceed 72.0 fluid ounces (1.32 lb ai/A) of Cheetah. Sequential applications must be at least 10 days apart.
- **DO NOT** apply Cheetah through any type of irrigation system.
- Refer to the **ROTATIONAL CROP RESTRICTIONS** section under the **PRODUCT INFORMATION** heading of this label for the appropriate rotational crop plant back intervals.

## APPLICATION DIRECTIONS FOR USE ON COTTON

Application of Cheetah to cotton varieties not containing the Glufosinate-resistant gene requires the use of hooded spray equipment designed to minimize exposure of the spray to the cotton stand. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

### Application Timing:

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

Applications for Cheetah on cotton may be made from emergence up to early bloom.

Cheetah is a foliar-active material with little or no soil-residual activity.

Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

For best result:

- On lambsquarters, Palmer amaranth and velvetleaf control, make applications of Cheetah between dawn and 2 hours before sunset.
- Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah.

### Application Rates:

#### Option 1 3 post applications

Apply 29 fl oz/A (0.53 lbs ai/A) per application depending on weed species, size and density per weed chart.

If required a second application of 29 fl oz/A (0.53 lbs ai/A) can be applied, followed by a third application of 29 fl oz/A (0.53 lbs ai/A).

The sequential applications must be made 10-14 days after each other.

The maximum annual rate of Cheetah on cotton is 87.0 fl oz/A (1.59 lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

#### Option 2 2 post applications

Apply 32 - 43 fl oz/A (0.59 - 0.79 lbs ai/A) per application depending on weed species, size and density per weed chart.

If required a second application of 29 fl oz/A (0.53 lbs ai/A) can be applied.

The sequential applications must be made 10-14 days after each other.

The maximum annual rate of Cheetah on cotton is 72 fl oz/A (1.32 lbs ai/A).

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

Use Pattern	1 <sup>st</sup> Application	2 <sup>nd</sup> Application Minimum 10 Days Up to 14 Days After 1 <sup>st</sup> Application	3 <sup>rd</sup> Application Minimum 10 Days Up to 14 Days After 2 <sup>nd</sup> Application	Annual Maximum
Option 1	29 fl oz/A (0.53 lbs ai/A)	29 fl oz/A (0.53 lbs ai/A)	29 fl oz/A (0.53 lbs ai/A)	87 fl oz/A (1.59 lbs ai/A)
Option 2	32-43 fl oz/A (0.59 - 0.79 lbs ai/A)	29 fl oz/A (0.53 lbs ai/A)	None	72 fl oz/A (1.32 lbs ai/A)

### Adjuvants:

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (including temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (including low relative humidity) or hard water.

The use of an anti-foam agent is advised.

### Surfactants / Oils:

The use of additional surfactants or crop oils in tank mixes with Cheetah may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

### Application Methods to Cotton:

Application of Cheetah to cotton varieties not containing the Glufosinate-resistant gene requires the use of hooded spray equipment designed to minimize exposure of the spray to the cotton stand. A hooded sprayer directs the spray onto weeds, while shielding the cotton stand from contact. Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Extreme care must be exercised to avoid exposure of the desirable vegetation to the spray.

With a hooded sprayer, the spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground. If the hoods are raised, spray particles may escape and come into contact with the cotton, causing damage or destruction of the crop.

Herbicide rates and spray volume instructions are presented as broadcast equivalents and must be reduced in proportion to the area actually treated. Use the following formulas to calculate the correct rate and volume per planted (field) acre:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Amount of banded product needed per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast spray VOLUME per acre} = \text{Banded spray volume needed per acre}$$

### Post-Harvest – Fall Burndown:

Cheetah may be applied as a post-harvest burndown treatment to fields (after cotton harvest). Up to 43.0 fl oz/A (0.79 lb ai/A) of Cheetah may be applied in a single application to control larger weeds growing in the crop at the time of harvest. If more than 29.0 fl oz/A (0.53 lb ai/A) is used in a single application, the annual total may not exceed 72.0 fl oz/A (1.32 lb ai/A), including all application timings.

### Tank Mix on Cotton:

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Certain herbicide tank mixes may aid in the performance of Cheetah. Cheetah may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the cotton to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Cheetah cannot be mixed with any product containing a label prohibition against such mixing.

### RESTRICTIONS TO THE DIRECTIONS FOR USE ON COTTON

- **DO NOT** apply to cotton in Florida, South of Tampa (Florida Route 60), or in Hawaii, except for test plots or breeding nurseries.]
- **DO NOT** apply within 70 days prior to cotton harvest.
- Up to 3 applications may be made to cotton per year at a maximum application rate of 29 fl oz/A (0.53 lb ai/A).
- **DO NOT** apply more than 87 fl oz (1.59 lb ai/A) (including all application timings) to cotton per year.
- Minimum retreatment interval is 10 days.
- If environmental conditions prevent timely applications resulting in large weeds or heavy infestations, a single application at up to 43 fl oz/A (0.79 lb ai/A) may be made to cotton. **DO NOT** apply more than 43 fl oz/A (0.79 lb ai/A) in a single application under this use scenario.
- **DO NOT** apply this product through any type of irrigation system.
- Refer to the "**Rotational Crop Restrictions**" section under the "**Product Information**" heading of this label for the appropriate rotational crop plant back intervals.

## APPLICATION DIRECTIONS FOR USE ON SOYBEANS CONTAINING THE GLUFOSINATE-RESISTANT GENE

Apply Cheetah only to soybeans containing the Glufosinate-resistant gene. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

### Application Timing:

Apply to small and actively growing weeds, targeting less than 3 inch weeds in height. For additional information on weed heights refer to the **WEED CONTROL FOR ROW CROPS** section.

Weed control may be reduced if application is made when heavy dew, fog, and mist/rain are present; or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

Applications for Cheetah on soybeans may be made from emergence up to bloom or R1 growth stage.

Cheetah is a foliar-active material with little or no soil-residual activity.

Cheetah is rainfast 4 hours after application, therefore, rainfall within 4 hours may necessitate retreatment.

For best result:

- On lambsquarters, Palmer amaranth and velvetleaf control, make applications of Cheetah between dawn and 2 hours before sunset.
- Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah.

### Application Rate:

Apply Cheetah at 29 – 43\* fl oz/A (0.53 – 0.79 lbs ai/A), depending on weed species, size and density per weed chart. \*Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A)

If a second application is needed, make the second application of 29 – 43\* fl oz/A (0.53 – 0.79 lbs ai/A), can be applied up to a yearly maximum of 87.0\* fl oz/A (1.59 lbs ai/A). \*Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A) and maximum annual rate in California is 72 fl oz/A (1.32 lbs ai/A).

The maximum annual rate of Cheetah on soybeans is 87.0\* fl oz/A (1.59 lbs ai/A). \*Maximum annual rate in California is 72 fl oz/A (1.32 lbs ai/A)

Use a minimum spray volume of 15 gallons per acre, unless there is a difficult to control situation (including dense canopy, large weeds or unfavorable growing conditions are present). In difficult to control situations use a minimum spray volume of 20 gallons per acre.

Use Pattern Rate Ranges		
1 <sup>st</sup> Application	2 <sup>nd</sup> Application Minimum of 5 Days After 1 <sup>st</sup> Application	Annual Maximum
29.0 to 43.0* fl oz/A (0.53 – 0.79 lbs ai/A)	29.0 to 43.0* fl oz/A (0.53 – 0.79 lbs ai/A)	87.0* fl oz/A (1.59 lbs ai/A)

\*Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A) with annual maximum of 72 fl oz/A (1.32 lbs ai/A).

#### Adjuvants:

Ammonium sulfate (AMS) may be used at 1.5 to 3.0 pounds per acre. Adjuvant rates are dependent on a variety of factors including tank mix partners, environmental conditions (including temperature) and potential for leaf burn.

AMS has shown to improve weed control of difficult-to-control weeds, like lambsquarters and velvetleaf, under difficult environmental conditions (including low relative humidity) or hard water.

The use of an anti-foam agent is advised.

#### Surfactants / Oils:

The use of additional surfactants or crop oils in tank mixes with Cheetah may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

#### Restrictions to the Directions For Use on Soybeans Containing the Glufosinate-resistant gene:

- **DO NOT** apply Cheetah within 70 days of harvesting soybean seed.
- **DO NOT** apply more than 87.0 fl oz/A (1.59 lbs ai/A) of Cheetah on soybeans per growing year.
- **DO NOT** apply more than 43.0\* fl oz/A (0.79 lbs ai/A) of Cheetah in a single application. \*Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A)
- **DO NOT** make more than 3 applications per year.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone-based antifoam agent may be added if needed.
- **DO NOT** apply Cheetah if soybeans show injury from prior herbicide applications or environmental stress (drought, excessive rainfall, etc.).
- **DO NOT** apply Cheetah through any type of irrigation system.
- Refer to the **ROTATIONAL CROP RESTRICTIONS** section under the **PRODUCT INFORMATION** heading of this label for the appropriate rotational crop plant back intervals.
- Sequential applications must be at least 5 days apart.

#### Soybean Tank Mix Instructions:

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Certain herbicide tank mixes may complement Cheetah. No additional surfactant is needed with any tank mix partner. Cheetah may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the soybean to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Cheetah cannot be mixed with any product containing a label prohibition against such mixing.

## APPLICATION DIRECTIONS FOR CANOLA, CORN, COTTON, AND SOYBEAN SEED PROPAGATION

Cheetah may be applied to select out susceptible "segregates", i.e., canola, corn, cotton, and soybean plants that **DO NOT** contain the Glufosinate-resistant gene during seed propagation.

#### • Canola Containing as Glufosinate-resistant gene:

Cheetah may also be used in canola seed propagation as a foliar spray to selectively eliminate canola plants that **DO NOT** carry a gene that imparts resistance to glufosinate-ammonium and as such, can be applied to remove susceptible segregates during canola seed propagation. Breeding material not possessing the glufosinate-ammonium resistance gene will be severely injured or killed if treated with this herbicide. See **APPLICATION DIRECTIONS FOR USE ON CANOLA CONTAINING THE GLUFOSINATE-RESISTANT GENE** for use rates and application timing.

• **Corn Containing the Glufosinate-resistant gene:**

Inbred lines, plants not containing the Glufosinate-resistant gene, will be severely injured or killed if treated with this herbicide. A hooded sprayer may be used to protect plants from coming into contact with the herbicide application. For the selection of resistant corn “segregates,” Cheetah may be applied at 22.0 fl oz/A (0.4 lbs ai/A) plus AMS at 3.0 pounds per acre (17.0 pounds per 100 gallons) when corn is in the V-3 to V-4 stage of growth, i.e., 3 to 4 developed collars. A second treatment of 22.0 fl oz/A (0.4 lbs ai/A) plus AMS at 3.0 pounds per acre may be applied when the corn is in the V-6 to V-7 stage of growth or up to 24 inches tall. Sequential applications must be at least 10 days apart. When temperatures exceed 85 °F, the rate of AMS can be reduced to 1.5 pounds per acre (8.5 pounds per 100 gallons) to reduce potential leaf burn.

• **Cotton Containing the Glufosinate-resistant gene:**

Cheetah may also be used in cotton seed propagation as a foliar spray to selectively eliminate cotton plants that **DO NOT** carry the Glufosinate-resistant gene and as such, can be applied to remove susceptible segregates during cotton seed propagation. Breeding material not containing the Glufosinate-resistant gene will be severely injured or killed if treated with this herbicide. See **APPLICATION DIRECTIONS FOR USE ON COTTON CONTAINING THE GLUFOSINATE-RESISTANT GENE** for use rates and application timing.

• **Soybeans Containing the Glufosinate-resistant gene:**

For the selection of resistant soybean “segregates,” Cheetah may be applied at up to 29.0 to 43.0\* fl oz/A (0.53 – 0.79 lbs ai/A) when soybean is in the third trifoliate stage. A second treatment of 29.0 to 43.0\* fl oz/A (0.53 – 0.79 lbs ai/A) may be applied up to but not including the bloom growth stage of soybean. Sequential applications must be at least 5 days apart. \* Maximum rate in California is 36 fl oz/A (0.66 lbs ai/A)

## **APPLICATION DIRECTIONS FOR USE ON LISTED TREE, VINE, AND BERRY CROPS**

Apply this to the tree, vine, and berry crops listed below. Uniform, thorough spray coverage is necessary to achieve consistent weed control.

### **REGISTERED CROPS**

#### **BERRIES:**

Crop Subgroup 13-B: Bushberry subgroup

Blueberry, highbush; blueberry, lowbush; currant; elderberry; gooseberry; huckleberry

Juneberry; lingonberry; salal

#### **CITRUS CROP GROUP 10-10:**

Orange or tangerine/mandarin, Calamondin; citron, citrus hybrids; Mediterranean Mandarin; orange, sour;

orange, sweet; satsuma darin; tachibana orange; tangerine (mandarin); tangelo; tangor, trifoliate orange;

cultivars, varieties and/or hybrids of these

Lemon or lime – Australian desert lime; Australian finger lime; Australian round lime; brown river finger lime;

kumquat; lemon; lime; mount white lime; New Guinea wild lime; Russel River lime; sweet lime; Tahiti lime;

cultivars, varieties and/or hybrids of these Grapefruit – Grapefruit; Japanese summer grapefruit; pummelo; tangelo; uniq fruit; cultivars,

varieties and/or hybrids of these.

**OLIVES:** all olive varieties

#### **POME FRUIT (CROP GROUP 11-10):**

##### **Crop Group 11. Pome Fruits Group**

Apple; crabapple; loquat; mayhaw; pear; pear, oriental; quince; azarole; hook; medlar; quince, Chinese; quince, Japanese; tejocote;

cultivars, varieties and/or hybrids of these

#### **STONE FRUIT (CROP GROUP 12-12):**

##### **Crop Group 12. Stone Fruits Group**

Apricot; cherry, sweet; cherry, tart; nectarine, peach; plum; plum, chicksaw; damson; plum, Japanese; plumcot; prune; capulin; jujube

and sloe; cultivars, varieties and/or hybrids of these.

#### **TREE NUTS (CROP GROUP 14 INCLUDING PISTACHIOS):**

Crop Group 14. Tree Nuts Group

Almond; beech nut; Brazil nut; butternut; cashew; chestnut; chinquapin; filbert (hazelnut), hickory nut, macadamia nut (bush nut), pecan,

pistachios, and walnut, black and English

**GRAPES:** all grape varieties (table, wine, and raisins)

#### **Application Rate and Timing:**

For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Cheetah. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. **DO NOT** retreat these weeds with Cheetah until sufficient regrowth has occurred.

Apply Cheetah as a directed spray to control undesirable vegetation in tree, vine, and berries listed on this label. Apply as a broadcast, banded, or spot treatment application depending on the situation to control weeds listed under the heading **Weeds Controlled in Tree, Vine and Berry Crops**. Avoid direct spray or drift to desirable vegetation. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of Cheetah may be necessary to control plants generating from underground parts or seed.

Avoid contact of Cheetah solution, spray, drift or mist with green bark, stems, or foliage, as injury may occur to trees, vines, and berries. **Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Contact of Cheetah with parts of trees, vines, or berries other than mature brown bark can result in serious damage.**

#### Application Methods for Broadcast Applications:

Apply Cheetah at the rates listed below for broadcast applications based on weed size and stage of growth.

Weed Size and Stage	Rate of this product
Weeds < 3 in height	48 fl oz/A (0.88 lbs ai/A)
Weeds < 6 in height pre tiller grasses	56 fl oz/A (1.02 lbs ai/A)
Weeds > 6 in height and/or grasses that have tillered	56-82 fl oz/A (1.02 – 1.5 lbs ai/A)

#### Application Methods for Banded Spray Applications:

Banded applications may be used using the following formula to calculate the amount of herbicide needed for orchard or vineyard strip sprays:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate per acre broadcast} = \text{Amount of herbicide needed for treatment}$$

#### Application Methods for Spot or Directed-Spray Applications:

For spot or directed spray applications: mix Cheetah at 1.7 fluid ounces of product (0.031 lbs ai) per gallon of water. Apply to undesirable vegetation foliage until wet but prior to runoff. Ensure uniform and complete coverage. Thoroughly clean the sprayer following use. **DO NOT** make spot or directed spray applications to tree or vine trunk as injury may occur.

#### Weeds Controlled in Tree, Vine and Berry Crops:

##### Broadleaf Weeds

Alkali sida	Fleabane, annual	Morningglory, ivyleaf	Smartweed, Pennsylvania
Ammannia, purple	Goosefoot	Morningglory, pitted	Sowthistle, annual
Arrowhead, California	Gromwell, field	Mullein, turkey	Spurge, prostrate
Buckwheat, wild	Groundcherry, cutleaf	Mustard, wild	Starthistle, yellow
Buffalobur	Groundsel, common	Nettle	Sunflower, common
Burclover, California	Herbit	Nightshade, black	Sunflower, prairie
Carpetweed	Jimsonweed	Nightshade, eastern black	Sunflower, volunteer
Chickweed, common	Knotweed	Nightshade, hairy	Swinecress
Chinese thornapple	Kochia	Pennycress	Thistle, Russian
Cocklebur, common	Lambsquarters, common	Pigweed, redroot	Turnip, wild
Copperleaf, Virginia	Lettuce, miner's	Pineapple-weed	Velvetleaf
Cudweed	Lettuce, prickly	Puncturevine	Vervain
Cutleaf eveningprimrose	London rocket	Purslane, common	Vetch
Dodder	Mallow, common	Radish, wild	Virginia copperleaf
Eclipta	Malva (little mallow)	Ragweed, common	Willowherb, panicle
Fiddleneck	Marestail	Ragweed, giant	
Filaree	Mayweed	Redmaids	
Filaree, redstem	Morningglory, entireleaf	Shepherd's-purse	

##### Grass Weeds

Barneyardgrass	Crabgrass, smooth	Junglerice	Shattercane
Bluegrass, annual	Cupgrass, woolly	Oat, wild	Sprangletop
Brome, rigput	Foxtail, giant	Panicum, fall	Stinkgrass
Bromegrass, downy	Foxtail, green	Panicum, Texas	Wheat, volunteer
Canarygrass	Foxtail, yellow	Rush, toad**	Windgrass
Chess, soft	Goosegrass	Ryegrass, annual*	Witchgrass
Crabgrass, large	Johnsongrass, seedling	Sandbur, field	

##### Biennial and Perennial Weeds

Aster, white heath	Clover, red	Horsetail	Paragrass	Thistle, musk
Bindweed, field	Clover, white	Lovegrass	Plantain	Torpedograss
Bindweed, hedge	Dallisgrass	Mugwort	Poison ivy/oak	Vaseygrass
Bluegrass, Kentucky	Dandelion	Mullein, common	Quackgrass	Woodsorrel
Bromegrass, smooth	Dock, curly	Mustard, tansy	Rocket, yellow	Yarrow, common
Bulrush**	Dogbank (hemp)	Nutsedge, purple	Rose, wild	
Burdock	Fescue	Nutsedge, yellow	<i>Rubus</i> spp.	
Canada thistle	Golden rod, gray	Onion, wild	Spurge, leafy	
Clover, Alsike	Guineagrass	Orchardgrass	Thistle, bull	

\* apply to annual ryegrass prior to 3 inches in height

\*\*indicates suppression

### Restrictions to the Directions For Use on Tree, Vine, and Berry Crops:

- **DO NOT** apply more than 164 fluid ounces of Cheetah per acre (3.0 lbs ai/A) to berry bushes and stone fruit in a 12 month period.
- **DO NOT** exceed the maximum single application rate of 82 fl oz/A (1.50 lb ai/A)
- **DO NOT** make more than 2 applications at a maximum application rate of 82 fl oz/A (1.5 lbs ai/A) per application to berry bushes and stone fruit.
- **DO NOT** apply more than 246 fl oz (4.5 lbs ai/A) of Cheetah per acre to tree, nuts, vines, pome fruit, citrus and olives in any calendar year.
- **DO NOT** make more than 3 applications at a maximum rate of 82 fl oz/A (1.50 lb ai/A) per application to tree nuts, vines, pome fruits, citrus and olives.
- **DO NOT** graze harvest, and/or feed treated orchard cover crops to livestock.
- **DO NOT** apply Cheetah through any type of irrigation system.
- **DO NOT** apply Cheetah aerially to tree, berry, or vine crops.
- **DO NOT** apply Cheetah within 14 days of nut, fruit, berry, or grape harvest.
- Applications to citrus fruits, pome fruits, olives and berry bushes must be a minimum of 14 days apart.
- Applications to stone fruit must be a minimum of 28 days apart.
- **DO NOT** make spot spray applications to suckers, as tree injury may occur.

### Sucker Control with Cheetah:

Cheetah will reduce or eliminate sucker growth when applied to suckers that are young, green, and uncalled. For sucker control, apply a split application approximately 4 weeks apart at 56.0 fluid ounces of product per acre (1.02 lbs ai/A). Coverage of all sucker foliage is necessary for optimum control. Suckers must not exceed 12 inches in length.

### Tank Mix Partner Instructions:

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Cheetah does not provide residual weed control or control of unexposed plant parts. Certain herbicide tank mixes may aid in the performance of Cheetah or be added to provide residual herbicide activity. No additional surfactant is needed with any tank mix partner. Cheetah may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. The tank mix partner must be used in accordance with the label limitations and precautions. No label dosage rates may be exceeded. Cheetah cannot be mixed with any product containing a label prohibition against such mixing.

diuron	napropamide	oryzalin	terbacil
flumioxazin	norfluzon	simazine	

### APPLICATION DIRECTIONS FOR POTATO VINE DESICCATION

#### Application Rates and Timing:

Apply Cheetah at the beginning of natural senescence of potato vines. Apply 21.0 fl oz/A (0.38 lbs ai/A). **DO NOT** split this application or apply more than 1 application (See restrictions section). Potato varieties with heavy or dense vines may require an application of another desiccation product to complete vine desiccation.

Thorough coverage of the potato vines to be desiccated is essential. Use a sufficient volume of water (20.0 to 100 gallons per acre) to obtain a thorough coverage of the potato vines. Vary the gallons of water per acre and the spray pressure as indicated by the density of the potato vines to assure thorough spray coverage. Increase the spray volume to at least 30.0 gallons of water per acre when the potato vine canopy is dense or under cool and dry conditions. Apply Cheetah with the spray boom as low as possible to achieve thorough coverage of the potato vines for best control and to minimize drift potential.

#### Restrictions to the Directions For Use in Potato Vine Desiccation:

- **DO NOT** apply more than 21.0 fl oz/A (0.38 lbs ai/A) to potato vines per year or per single application.
- **DO NOT** harvest potatoes until 9 days or more after application of Cheetah.
- **DO NOT** apply to potatoes grown for seed.
- Potatoes, canola, corn, cotton, soybean, and sugar beets may be planted at any time after the application of Cheetah as a potato vine desiccant.
- **DO NOT** plant treated areas to wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale until 30 or more days after an application of Cheetah as a potato vine desiccant.
- **DO NOT** plant treated areas to crops other than those listed in this use precautions section until 120 or more days after an application of Cheetah as a potato vine desiccant.
- **DO NOT** split this application or apply more than one application per harvest (per year).

### FALLOW FIELDS OR POST HARVEST

Cheetah may be used as a substitute for tillage to control or suppress weeds in the **WEED CONTROL FOR ROW CROPS** section of this label. Applications may be made in fallow fields, post harvest, prior to planting or emergence of any crop listed on this label.

Apply Cheetah at 22.0 or 29.0 fl oz/A (0.2 to 0.53 lb ai/A) to fallow fields to control specific weeds. Cheetah must be applied with ammonium sulfate. Tank mixes with 2,4-D, glyphosate or atrazine are specified with Cheetah to enhance total weed control. When using

Cheetah in tank mix combinations, follow the precautions and directions of use of the most restrictive label. See **APPLICATION AND MIXING PROCEDURES** section of this label for additional information on how to apply Cheetah. See the **PRODUCT INFORMATION** section of this label for rotational crop restrictions.

**RESTRICTIONS TO THE DIRECTIONS FOR USE ON FALLOW FIELDS OR POST HARVEST**

**DO NOT** apply more than 29 fl oz/A (0.53 lbs ai/A) in a single application.

**DO NOT** make more than 3 applications per year.

**DO NOT** make sequential applications sooner than 14 days apart.

**DO NOT** apply more than 87 fl oz/A (1.59 lbs ai/A) per year.

**APPLICATION DIRECTIONS FOR USE ON NONCROPLAND USE SITES**

When applied as directed, Cheetah controls undesirable plant vegetation in public, military, and non-crop areas including private, public and military lands in the following areas: airfields, airports, alleys, lanes, paths, trails, access roads, around commercial or industrial structures or outbuildings, around farm and ranch structures and outbuildings, site preparation areas for conifer and hardwood, bare ground, barrier strips, beaches\*, campgrounds, construction sites, ditch banks, drive-in theaters, driveways and ramps, dry ditches and canals, fences and fencerows, firebreaks, gravel yards, Conservation Reserve Program (CRP)\*, habitat restoration and management areas, highways and roadsides (including aprons, medians, guardrails and right of ways), industrial plant sites, industrial areas, lumber yards, natural areas, parking areas, parks, paved areas, petroleum and other tank farms, pumping installations, pipeline, power, telephone and utility rights-of-way, sewage disposal areas, fuel storage areas, power stations, railroad rights-of way, recreation areas, refineries, resorts, schools, sidewalks, sports areas, storage areas, substations, tennis courts, shelter belts, uncropped farmstead areas, vacant lots, walkways, wastelands, wildlife openings, wildlife habitat areas, wildlife food plots\*.

\*Not for use in California

**Industrial:** This product may be used to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way, and wayside structures. This product may be tank mixed with other herbicides for these use sites unless specifically prohibited by the product label.

**Conservation Reserve Program (CRP)\*:** This product can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres. For selective applications with broadcast spray equipment, apply 48 to 56 fl. oz./A (0.88 to 1.0 lb. ai/A) of this product in early spring before desirable CRP grasses, including crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if applications are made when plants are not dormant.

**Wildlife Food Plots\*:** This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling.

**Site Preparation for Conifer and Hardwood Production Areas:** This product may be used as a site preparation treatment prior to planting conifer and hardwood species. **DO NOT** apply Cheetah as an over-the-top broadcast spray to desirable conifer or hardwood plantings. Seedling conifer and hardwood trees may be planted into the treated area after the restricted entry interval (REI) of 12 hours has elapsed.

\*Not for use in California

**Side Trimming:** To control only a portion of the plant, direct the spray solution to thoroughly cover (spray to wet) only the portion of the plant to be controlled.

**Site Preparation for Conifer and Hardwood Production Areas:** When applied in site preparation prior to planting conifer and hardwood species, this product will provide control of undesirable vegetation. Seedling conifer and hardwood trees may be planted into the treated area after the restricted entry interval (REI) of 12 hours has elapsed.

**Brush Control:** This product will provide control or suppression of the perennial woody species (brush) listed below. Use this product at rates from 1 to 3 quarts of this product per acre to impact the growth of woody plants and not to exceed 1.9 gallons per acre per year (4.5 lb ai/A/yr). Non-ionic surfactants (NIS) or methylated seed oils (MSO) may be used when making foliar applications. Follow any instructions on the surfactant manufacturer's label.

For hard-to-control woody plants including elm, certain oaks or when plant leaf surfaces have hardened off, use the higher rate of this product or tank mix this product with other herbicides registered for control of these woody plants. Higher specified rates per acre of this product must be used when conditions are not optimum for spray coverage, including when weed growth is heavy or dense. Lower specified rates must be used when the target species is conifer and when vegetation growth conditions allow for uniform spray coverage.

\* Not for use in California

**USE RESTRICTIONS for NONCROP:**

- **DO NOT** apply more than 246 fl. oz. of this product per acre per year (4.5 lb. ai/A/year).
- **DO NOT** apply more than 82 fl. oz. of this product per acre per single application (1.5 lb. ai/A/application).
- **DO NOT** apply more than a total of 3 broadcast applications (excluding spot treatments) per year.
- **DO NOT** exceed a maximum total of 4.5 lb. ai/A/year.
- Minimum re-treatment interval is 5 days.



## APPLICATION RATES

Mix 0.5 to 2.0 fl. oz. (0.009 to 0.036 lb. ai) of this product per gallon of spray solution (24 to 82 fl. oz./A (0.44 to 1.5 lb. ai/A) and apply 1 gallon of spray solution to 1,000 square feet to actively growing weeds. Adjust application rate as needed when using spray volumes delivering greater or less than 1 gallon per 1,000 square feet. Determine proper use rate based on weed size in Table 1. Larger weeds will require a higher use rate and see Table 1 for details.

**Table 1: USE RATE FOR THIS HERBICIDE**

Apply this product at the rates listed below for broadcast applications based on weed size and stage of growth.

Weed Size and Stage	Rate of this product (Per Gallon of Water)	Rate of this product (Per 1,000 sq. ft.)	Rate of this product (Per Acre)	Spot Spray % Solution
Easily Controlled Weeds < 3 in height*	0.5 fl. oz. (0.009 lb. ai)	0.5 fl. oz. (0.009 lb. ai)	24 fl. oz./A (0.44 lb. ai)	0.5
Weeds < 3 in height	1.0 fl. oz. (0.018 lb. ai)	1.0 fl. oz. (0.018 lb. ai)	48 fl. oz./A (0.88 lb. ai)	0.5-0.75
Weeds < 6 in height pre-tiller grasses	1.25 fl. oz. (0.023 lb. ai)	1.25 fl. oz. (0.023 lb. ai)	56 fl. oz./A (1.0 lb. ai)	0.75-1.25
Weeds > 6 in height and/or grasses that have tillered	1.25 to 2.0 fl. oz. (0.023 to 0.036 lb. ai)	1.25 to 2.0 fl. oz. (0.023 to 0.036 lb. ai)	56-82 fl. oz./A (1.0 to 1.5 lb. ai)	1.25-1.5

\*See Weeds Controlled Table below for details.

For spot or directed spray applications by backpack sprayers, mix this product at 0.5 to 2.0 fl. oz. of product (0.009 to 0.036 lb. ai) per gallon of water. Larger and more difficult to control weeds require a higher use rate. When using the per gallon rate, calibrate sprayers to deliver 1 gallon of spray solution per 1,000 square feet. Adjust application rate as needed when using spray volumes delivering greater or less than 1 gallon per 1,000 square feet. Thorough spray coverage of weeds is necessary to maximize weed control. Spray coverage needs to be uniform, but **DO NOT** spray to the point of runoff. Thoroughly clean the sprayer following use. **DO NOT** make spot or directed spray applications to desired plant foliage or stems as injury may occur.

### Use Restrictions:

- **DO NOT** apply this product within any enclosed structure in residential or commercial landscapes.
- **DO NOT** apply this product over-the-top as a broadcast application to ornamentals, conifers or hardwood plantings.
- **DO NOT** apply this product over-the-top of ornamental plants, and **DO NOT** allow spray of this product to contact or drift onto the foliage, green stems, exposed roots or fruit of desirable plants. Avoid application of this product under conditions that favor drift of sprays onto desired ornamentals or residential lawns.

This product offers postemergence control of susceptible grasses, sedges and broadleaf weeds (See WEEDS CONTROLLED Table), as well as additional mode of action to assist in the control of resistant weeds.

**IMPORTANT:** Contact with spray or spray drift of this product may cause severe injury or destruction of certain desirable plants, especially herbaceous species including bedding plants or direct seeded annual and perennial flowers. The use of spray shields that limit the plant exposure to this product is highly advised when applying this product near desirable plants.

### HOW TO APPLY

#### Spot or Directed Applications

This product may be used as a spot- or directed-spray application. Prepare the desired volume of spray solution by mixing this product in water with the amounts indicated in the following table:

**Table 2.** Amount of this product added to water to make 1, 25, or 100 gallons of spray solution at dosages of ½ to 3%. See Table 1 for % solution to use based on target vegetation.

% SOLUTION	VOLUME OF SPRAY SOLUTION		
	1 GALLON	25 GALLONS	100 GALLONS
	THIS PRODUCT		
0.5 %	0.75 fl. oz.	1 pint	1/2 gallon
1%	1.5 fl. oz.	1 quart	1 gallon
2%	3 fl. oz.	2 quarts	2 gallons
3%	4.5 fl. oz.	3 quarts	3 gallons

Select appropriate solution concentration and spray undesirable vegetation foliage on a spray-to-wet basis. **DO NOT** apply beyond runoff. Ensure uniform and complete coverage. Use a coarse spray. To minimize drift, avoid spraying during windy conditions. Backpack, pump-up, and hydraulic sprayers may be used. Thoroughly clean the sprayer following use.

**Use of Spray Adjuvants:** The addition of a nonionic antifoaming agent may reduce foaming, especially when using soft water. The use of Methylated seed oil (MSO) at 1% v/v (1 gal. per 100 gal. of spray solution) or non-ionic surfactant (NIS) at a minimum rate of 0.25% v/v (1 qt. per 100 gal. of spray solution) may be used for foliar applications. The addition of 8.5 to 17.0 lbs. of ammonium sulfate (spray grade) per 100 gal. of water (1 to 2% by weight) or 2 to 4 lbs. of ammonium sulfate per acre may result in better weed control.

This herbicide can be tank mixed with other non-selective herbicides including glyphosate and preemergence residual herbicides including flumioxazin. Follow the most restrictive label restrictions and precautions for each product. A combination with a residual herbicide including flumioxazin provides effective control of existing weeds as well as lasting residual weed control in areas including landscape beds and xeriscapes.

**Aerial Applications (Helicopter Application Only):** Use a drift control device including a "Microfoil," or "Thru Valve-Boom" or equivalent drift control system when applying as a foliar treatment to utility rights-of-way, tree production areas, ditch banks or other approved sites that may be near susceptible crops. The application volume required will vary with the height and density of the vegetation and the application equipment used. Generally, aerial applications will require a minimum of 15 gallons per acre to ensure thorough coverage. Drift control additives may be used. If a drift control additive is used, observe and follow all directions and precautions as specified on the additive label.

**Foliar Treatments with Ground Equipment**

- **High Volume Applications:** Use high volume applications for optimum performance when spraying medium to high density vegetation. Use equipment calibrated to deliver 50 to 100 gallons of finished spray per acre. For best results, make sure that the targeted plant foliage is thoroughly covered.
- **Low Volume Applications:** Use low volume applications when brush height is less than 6 feet and brush cover is less than 60% of the area. Use equipment calibrated to deliver 10 to 50 gallons of finished spray per acre.

**Broadcast Applications with Ground Equipment:** Use equipment calibrated to deliver 20 to 100 gallons of finished spray per acre. The amount of spray solution to use will depend on the height and density of the brush. Use spray nozzles and equipment that will provide thorough coverage of the targeted brush species.

BRUSH* SUPPRESSED OR CONTROLLED	
Blackberry	Poison ivy/oak
Deer brush	Pine
Douglas fir	Roundleaf
Gallberry	Greenbrier
Hazel	Salmonberry
Honeysuckle	Sweetgum
Huckleberry	Sumac
Maple	Thimbleberry
Multiflora rose	Trumpetcreeper
Oak	Vine Maple
	Western Red Cedar
*not for use in California	

## WEEDS CONTROLLED

Alfalfa+	Cupgrass, woolly	Marshelder, annual+	Rocket, yellow
Alkali sida	Cutleaf eveningprimrose	Mayweed	Rose, wild
Amaranth, Palmer+	Dallisgrass	Milkweed, common***+	Rubus spp.
Ammannia, purple	Dandelion	Milkweed, honeyvine***+	Rice, red+
Anoda, spurred*^	Devil's claw*^	Millet, wild proso+	Rice, volunteer+
Arrowhead, California	Dock, curly	Millet, proso volunteer+	Rush, toad***
Artichoke, Jerusalem+	Dock, smooth+	Morningglory, entireleaf	Ryegrass, annual**
Aster, white heath	Dodder	Morningglory, ivyleaf	Sandbur, field
Bahiagrass	Dogbane (hemp)	Morningglory, pitted	Senna coffee+
Barley, volunteer*^	Eclipta	Morningglory, sharpod*^	Shattercane
Barnyardgrass*	Fescue	Morningglory, smallflower+	Shepherd's Purse
Beggarweed, Florida+	Fleabane, annual	Morningglory, tall+	Sicklepod (java bean)+
Bermudagrass+	Fiddleneck	Mugwort	Sida, prickly+
Bindweed, field	Filaree	Muhly, wirestem***+	Signalgrass, broadleaf*^
Bindweed, hedge	Filaree, redstem	Mullein, common	Smartweed, Pennsylvania
Black medic+	Foxtail, bristly+	Mullein, turkey	Smellmelon+
Bluegrass, annual	Foxtail, giant	Mustard, tansy	Sowthistle, annual
Bluegrass, Kentucky	Foxtail, green	Mustard, wild	Sowthistle, perennial+
Blueweed, Texas+	Foxtail, robust purple+	Nettle	Soybean, volunteer+
Brome, ripgut	Foxtail, yellow*^	Nightshade, black	Sprangletop
Bromegrass, downy	Gallinsoga, hairy+	Nightshade, eastern black	Spurge, prostrate*^
Bromegrass, smooth	Gallinsoga, small flower+	Nightshade, hairy	Spurge, leafy
Buckwheat, wild	Geranium, cutleaf+	Nightshade, silverleaf+	Spurge, spotted*^
Buffalobur	Goosefoot	Nutsedge, purple	Starbur, bristly+
Burhulm***	Goosegrass*^	Nutsedge, yellow	Starthistle, yellow
Burclover, California	Goldenrod, gray	Oat, wild*^	Stinkgrass
Burcucumber+	Gromwell, field	Onion, wild	Sunflower, common
Burdock	Groundcherry, cutleaf	Orchardgrass	Sunflower, prairie*^
Bursage, woolyleaf+	Groundsill, common	Panicum, fall*^	Sunflower, volunteer
Canarygrass	Guineagrass	Panicum, Texas	Swinecress
Carpetgrass	Hempnettle+	Paragrass	Thistle, bull
Carpetweed	Henbit	Pennygrass	Thistle, Canada
Catchweed bedstraw (cleavers) *^	Horsenettle, Carolina*^	Pigweed, redroot*^	Thistle, musk
Chess, soft	Horsetail	Pigweed, prostrate*^	Thistle, Russian
Chickweed, common	Johnsongrass, rhizome+	Pigweed, spiny*^	Timothy+
Chickweed, mouse-ear+	Johnsongrass, seedling*^	Pigweed, smooth*^	Torpedograss
Chinese thornapple	Jimsonweed	Pigweed, tumble*^	Turnip, wild
Clover, Alsike	Junglerice*^	Pineapple weed	Vaseygrass
Clover, red	Knotted*^	Plantain	Velvetleaf*^
Clover, white	Kochia	Pointsettia, wild+	Vervain
Cocklebur, common	Ladysthumb+	Poison ivy/oak	Vetch
Copperleaf, hophornbeam+	Lambquarters, common	Pokeweed+	Waterhemp, common+
Copperleaf, Virginia	Lettuce, miners	Puncturevine	Waterhemp, tall+
Corn, volunteer+	Lettuce, prickly	Purslane, common*^	Wheat, volunteer
Cotton, volunteer+	London rocket	Pusley, Florida+	Willowherb, panicle
Crabgrass, large*^	Lovegrass	Quackgrass	Windgrass
Crabgrass, smooth*^	Mallow, common	Radish, wild	Witchgrass
Croton, tropic*^	Mallow, Venice+	Ragweed, common	Woodsorrel
Croton, woolly*^	Malva (little mallow)	Ragweed, giant	Wormwood, biennial+
Cudweed	Marestail	Redmaids	Yarrow, common

+Not for use in California

^Use rate in California 24 fl. oz./A (0.44 lb. ai)

\*easily controlled species

\*\*apply to annual ryegrass prior to 3 inches in height

\*\*\*indicates suppression only

# STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** **DO NOT** use or store near heat or open flame. Keep container tightly closed and dry in a cool, well ventilated place. Storage temperature must not exceed 125° F. If storage temperature of this product is below 32° F, the material must not be pumped until its temperature exceeds 32° F. Protect against direct sunlight.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

## **CONTAINER HANDLING:**

**NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "No refillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size.

**Non-refillable Containers 5 Gallons or Less:** Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Non-refillable Containers Larger than 5 Gallons:** Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Containers Larger than 5 Gallons:** Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable Container:** Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

**SEED DISPOSAL:** To dispose of out-of-date or otherwise unmarketable seed from plants, which have been treated with Cheetah, broadcast and lightly incorporate seed into field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternatively, seed may be destroyed by deep burial, incineration or landfill disposal.

## WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

## LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you DO NOT agree with or DO NOT accept any of directions for use, the warranty disclaimers, or limitations on liability, DO NOT use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

RV011520

Cheetah is a registered trademark of Nufarm Americas Inc.