DOC ID:584322



Classified for "RESTRICTED USE" in New York State under 6NYCRR Part 326

ACCEPTED VIA NOTIFICATION LABEL NOT REVIEWED

Nov. 22, 2022

New York State Department of Environmental Conservation

Pesticide Product Registration

SPECIALTY FUNGICIDE

Eagle[®] 20EW

A systemic, protectant and curative fungicide for disease control in established turfgrass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruits and grapes

Group 3 FUNGICIDE

Active Ingredient

Keep Out of Reach of Children CAUTION

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Using

EPA Reg. No. 62719-463 EPA Est. 5905-GA-01

CD02-195-020

^{TM®}Trademarks of Corteva Agriscience and its affiliated companies Produced for Corteva Agriscience LLC

9330 Zionsville Road Indianapolis, IN 46268

NET CONTENTS 1 PT

Precautionary Statements

Hazards to Humans and Domestic Animals

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Note to Physician: This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- · Chemical-resistant gloves made from barrier laminate
- · Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants
- · Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE)and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- · Chemical-resistant gloves made from any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in a cool, dry area above freezing. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continues to drain exposed. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

General Information

Shake Well Before Using

Eagle® 20EW specialty fungicide is a systemic, protectant and curative fungicide for the control of the diseases listed on this label in established turfgrass (including residential and commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), landscape ornamentals, greenhouse and nursery ornamentals, and non-commercial tree fruits and vines, specifically apples, stone fruits and grapes. Optimum disease control is achieved when this product is applied in a regularly scheduled preventive program.

General Use Precautions

Fungicide Resistance Management

Eagle 20EW belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Eagle 20EW should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. After two consecutive applications of Eagle 20EW, another myclobutanil product, or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from Eagle 20EW. Apply the alternate products within the intervals specified on the label for Eagle 20EW. Do not apply Eagle 20EW at rates below those specified on the label. If tank mixing, use the full label rate of Eagle 20EW with the full label rates of other products effective on the target pest. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

The following practices can delay development of fungicide/bactericide resistance:

- Avoid the consecutive use of Eagle 20EW or other target site of action Group 3 fungicides/bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local company representative or by calling 800-258-3033.

Mixing Directions

Be sure sprayer is clean and not contaminated with other materials prior to use. Fill the spray tank 1/4 to 1/2 of the total amount of water required for the load. Start agitation and maintain agitation throughout mixing and application. Add the required amount of Eagle 20EW directly into the spray tank. Complete filling the tank. Always add Eagle 20EW to the spray tank before adding other materials.

Compatibility

Eagle 20EW is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialist prior to use. When an adjuvant is to be used with this product, Corteva Agriscience recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Carefully read, understand and follow label use rates and restrictions. For proper application, determine the size of the area to be treated, the specified label use rate and the gallonage to be applied to the area. Under low disease conditions, minimum label use rates per application can be used. Use maximum label rates and shortened spray schedules for severe or threatening disease conditions. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is recommended prior to use.

Ground Application

Thorough coverage sprays generally result in optimum disease control. Application equipment must be properly calibrated and provide uniform spray coverage.

Handgun or Pressurized Sprayers: For best results when applying this product on a protectant schedule, ensure thorough coverage of all plant parts.

Chemigation Application

Eagle 20EW must be applied on a regular protectant fungicide schedule, not an irrigation schedule. If irrigation cycles are less frequent than the application intervals for Eagle 20EW, ground or handgun applications must supplement chemigation applications to achieve adequate disease control. Directions for Sprinkler Chemigation: Apply this product only through solid set or hand-move sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Chemigation Equipment Preparation: The following use directions are to be followed when this product is applied through irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injection system with scoap or a cleaning agent and water. Determine the amount of Eagle 20EW needed to cover the desired area. Mix according to instructions in the Mixing Directions section. Continually agitate the mixture during mixing and application.

Chemigation Equipment Calibration: In order to calibrate the irrigation system and injector to apply the mixture containing Eagle 20EW, determine the following: 1) Determine area covered by sprinkler; 2) Fill injector solution tank with water and adjust flow rate to use the contents over a 10- to 30-minute interval; (3) Determine the amount of Eagle 20EW required for treatment area; 4) Add the required amount of Eagle 20EW into the same quantity of water used to calibrate the injection equipment. Maintain constant solution tank agitation during the injection period. Operate system at normal pressures specified by the manufacturer of the injection equipment and used for the time interval established during calibration. Inject Eagle 20EW at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention. Stop injection equipment after treatment is completed. Continue to operate the system until the solution with Eagle 20EW has cleared the last sprinkler head.

Chemigation Equipment Requirements:

- The system must contain an air gap, an approved backflow prevention device, a functional check valve, vacuum relief valve (including inspection port), and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information or state specific regulations.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- To insure uniform mixing of the fungicide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all backflow prevention devices on the water line.
- The tank holding the fungicide mixture should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector point.

Chemigation Precautions:

- Crop injury, lack of fungicidal effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.
- Public water system means a system for the provision to the public of piped water for human consumption if such system that has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection.

Chemigation Restrictions:

- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quickclosing check valve to prevent the flow of fluid back toward the injection.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the pesticide labelprescribed safety devices for public water systems are in place with current certification. Specific local regulations may apply and must be followed.
- Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate nontarget areas.
- Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops.

- Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required PPE is worn.
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

Uses

Established Turfgrass

Use Eagle 20EW in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Use diagnostic kits, extension experts, or other identification methods when developing disease control strategies.

In non-residential turfgrass (including commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), optimum disease control is achieved when Eagle 20EW is applied in a preventative disease control program at a rate of 1 to 2.4 fl oz per 1000 sq ft. In residential turfgrass, optimum disease control is achieved when Eagle 20EW is applied in a preventative disease control program at a rate of 1.2 fl oz per 1000 sq ft. See the tables below for specific application rates for various diseases.

Apply Eagle 20EW in sufficient water to ensure thorough coverage. For foliar diseases, use approximately 1 gallon of water per 1000 sq ft. Use 2 to 3 gallons of spray solution per 1000 sq ft to control diseases causing root and crown rots. Under conditions favorable for high disease development, reduce the spray interval between applications of Eagle 20EW. Under light to moderate disease pressure, apply Eagle 20EW at the low use rate and/or longer treatment interval. When disease pressure is high or when used as a curative treatment, use higher rates of Eagle 20EW and/or shorter treatment interval unless otherwise specified. Non-Residential Turfgrass¹

Diseases	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
anthracnose red thread septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft
brown patch		14	Begin applications when conditions are favorable for disease development, but before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore [®] T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.	per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).

Non-Residential Turfgrass¹ (Cont.)

Diseases	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
copper spot zonate leaf spot	1.2	14	Apply when conditions are favorable	Do not apply more than 13.8 fl oz of
crown rot leaf spot melting-out			for disease development.	Eagle 20EW per 1000 sq ft per year. • For Nassau
dollar spot	0.5	7	Apply when conditions are favorable for disease development.	and Suffolk Counties in New York State, do not apply more
		14	Tank mix with a low label rate of chlorothalonil.	than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year
	1	21 – 28	Tank mix with the label rate of chlorothalonil.	(1.95 lb myclobutanil per acre).
	1 – 2.4	14 - 28	If using this rate without tank mixing, make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	

Non-Residential Turfgrass¹ (Cont.)

Diseases	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
fusarium blight	1.2 - 2.4	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.	 per year. For Nassau and Suffolk Counties in New York State, do not
gray leaf spot	1.2 - 2.4	14	Apply when conditions are favorable for disease development. If using the lower rate, tank mix with a registered contact fungicide at its specified rate.	apply more than 3.43 floz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
leaf smuts	1.2		Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	

Non-Residential Turfgrass¹ (Cont.)

Diseases	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
necrotic ring spot	1.2 - 2.4	spring: 28	Make applications on a preventative basis in early to mid-spring.	Do not apply more than 13.8 fl oz of Eagle 20EW
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant. Apply 2.4 fl oz per 1000 sq ft followed by a second application one month later.	per 1000 sq ft per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
powdery mildew rusts	1.2	14 - 28	Apply when conditions are favorable for disease development.	
spring dead spot	2.4	fall: 28	Make 1 to 2 applications in the fall before turfgrass dormancy. Make a second application one month later.	

Diseases	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
summer patch	1.2 - 2.4	14 - 28	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrase sextension experts. Use at least 2 to 3 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	 Do not apply more than 13.8 fl oz of Eagle 20EW per year For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per year (1.55 lb myclobutanil per acre).

Non-Residential Turfgrass¹ (Cont.)

Diseases	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
take-all patch	2.4	spring/ fall: 28	To reduce the severity, make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F, and 1 to 2 spring applications in April and May depending upon local recommendations.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz
zoysia large patch		fall: 28	Make applications in fall before turfgrass dormancy.	of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).

¹Including commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens.

Residential Turfgrass

Disease	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
anthracnose red thread septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft
brown patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore® T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.	per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.55 ib myclobutanil per acre).
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	

Residential Turfgrass (Cont.)

Disease	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
crown rot leaf spot melting-out	1.2	14	Apply when conditions are favorable for disease development.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft
dollar spot			Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	per 1000 sq ft per year. • For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
fusarium blight			Apply when conditions are favorable for disease development.	
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.	

Residential Turfgrass (Cont.)

Disease	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
gray leaf spot	1.2	14	Apply when conditions are favorable for disease development.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft
leaf smuts		14	Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	per year. • For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year
necrotic ring spot		spring: 28	Make applications on a preventative basis in early to mid-spring.	
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	(1.95 lb myclobutanil per acre).
powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.	

Residential Turfgrass (Cont.)

Disease	Eagle 20EW (fl oz/ 1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
summer patch	1.2	14	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Use at least 2 to 3 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).

Landscape, Greenhouse and Nursery Ornamentals

Eagle 20EW is a locally systemic fungicide having protectant and curative properties that will translocate to new growth. For best control of labeled diseases, achieve thorough coverage of all plant parts on a protective application schedule. For dilute application sprays (>100 gallons of spray volume per acre) applied to ornamental plants in greenhouses, field grown plantings or in commercial and residential landscapes, apply Eagle 20EW at the rate of 6 to 12 fl oz per 100 gallons of spray volume on a 10- to 14-day application schedule unless otherwise directed. Use the higher rate under conditions of high disease pressure and/or optimum conditions for infection. For concentrate sprays (<100 gallons of spray volume per acre), apply 8 fl oz per acre on a 10- to 14-day application schedule.

The addition of a non-phytotoxic spray adjuvant will improve spray coverage and fungicidal performance. Maintain treated plants in a vigorous growing condition. Plants under nutritional or water stress will not respond as well to treatment as well-maintained plants. Overdosage of Eagle 20EW can result in observable foliar greening, thickened leaves, and/or shortened internodes. If this condition is observed, reduce the fungicide use rate but do not extend the application schedule.

Crop Tolerance

Plant tolerances are acceptable in the specific plants listed on this label. It is not possible to evaluate all ornamental plant species or varieties for tolerance to Eagle 20EW. The user should test for possible phytotoxic responses by treating a limited number of plants, at specified use rates, prior to initiating large-scale use.

The effects of spraying Eagle 20EW in combination with plant growth regulators are not fully understood at this time. If the use of a plant growth regulator is planned in an area being treated, the user should test for possible enhanced growth regulatory effects by treating a small number of plants, at the specified use rates of all products, prior to initiating large-scale use. Since the effectiveness of such products depends upon not just plant species or cultivar but also weather and seasonable differences (e.g., daylight hours), it is recommended that tests be repeated on previously tested varieties as environmental factors change and that observations for growth regulatory responses be made at regular intervals.

Specific Use Directions for Chrysanthemum

Foliar Sprays: Best control is achieved by thorough coverage sprays applied to point of runoff on a protectant application schedule. Use Eagle 20EW at a rate of 8 fl oz per 100 gallons of spray mixture. Do not apply more than 19 fl oz of Eagle 20EW (0.25 lb myclobutanii) per acre per application. Apply on a 10- to 14-day schedule (not to exceed 21 days).

Prestick Dip Treatment: Chrysanthemum cuttings may be treated by a dip procedure prior to planting as follows: Prepare a dip suspension at a concentration equivalent to 8 fl oz of Eagle 20EW per 100 gallons of water. Fully submerge cuttings in the dip suspension until wet throughout (do not submerge cuttings for more than 2 minutes). If cuttings are dipped, this procedure is the first spray under the quarantine program. Dispose of used dip suspension if it becomes contaminated with soil, plant debris or other foreign matter. Dispose of used dip suspension by spraying it onto registered crops (but not onto previously dipped cuttings) after filtering, or in a manner consistent with local, state, and federal guidelines.

Note: All infected plant material must be destroyed if your state is under quarantine directive.

Specific Use Restrictions:

- Do not apply more than 20 fl oz of Eagle 20EW (0.25 lb myclobutanil) per acre per application. On a total volume per acre basis, do not apply more than 333 gallons of spray per acre at the 6 fl oz per 100 gallons rate or 167 gallons per acre at the 12 fl oz per 100 gallons rate per application.
- Do not apply more than 153 fl oz of Eagle 20EW (2 lb myclobutanil) per acre per year.
- Do not use treated plant materials for food or feed.
- Do not apply to landscape, greenhouse and nursery ornamentals in Nassau and Suffolk Counties in New York State.

Crops	Diseases	Directions	Precautions/ Restrictions
abelia	cercospora		
acalypha (copper-leaf)	leaf spot powdery mildew		
achillea (yarrow)	powdery mildew rust		
African violet	powdery mildew		
ageratum	powdery mildew		
alder	rust		
almond, flowering	blossom blight (<i>Monilinia</i> spp.)	Apply prebloom, 50% bloom and at petal fall.	
amelanchier (juneberry, shadbush)	fabraea leaf spot powdery mildew rust		

Do not apply to carrotwood (Cupaniopsis anacardioides).

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
amorpha (false indigo)	cercospora leaf spot powdery mildew rust		
anemone	rust		
angelica	cercospora leaf spot rust		
ash	rust		
aster	powdery mildew rust		
Australian pine	diplodia tip blight		
azalea	petal blight (<i>Ovulinia</i> spp.) powdery mildew	Begin applications when flowers start to exhibit color.	
barberry	powdery mildew rust		May cause temporary damage to crimson pigmy and other atropurposis varieties.
begonia	powdery mildew		
bellflower	cercospora leaf spot powdery mildew rust		
birch	rust		
bittersweet	powdery mildew		
buckeye			

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
buttonbush	cercospora leaf blight powdery mildew rust		
calendula	cercospora leaf spot		
California poppy	powdery mildew		
canna lily	rust		
carnation	powdery mildew rust		
catalpa	cercospora leaf spot powdery mildew		
cherry, flowering	leaf spot powdery mildew		
chestnut, horse	powdery mildew		
China aster	rust		
chokeberry	rust twig and fruit blight		Fruit may not be used for food or feed.
Christmas trees	rust		
chrysanthemum	ascochyta blight rust white rust		
columbine	rust		
cornflower			
cosmos	powdery mildew		
cottonwood			
crabapple, flowering	powdery mildew rust scab		

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
crepe-myrtle	powdery mildew		
daffodil	rust		
dahlia	powdery mildew		
delphinium	powdery mildew rust		
dogwood	anthracnose powdery mildew septoria leafspot		
Douglas fir	needle rust	Apply 12 to 18 fl oz per acre starting early spring. Continue applications at 2- to 3-week intervals until the threat of infection has passed. Spray adjuvants must be added to spray solutions to obtain good spray coverage and disease control.	
dianthus	rust		
elm	powdery mildew		
euonymus			
fern	rhizoctonia aerial blight		
fleabane	cercospora leaf spot powdery mildew rust		

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
four o'clock	rust		
fuchsia]		
gaillardia	powdery mildew		
gardenia	rust		
geranium			
gerbera daisy	powdery mildew		
gourd, ornamental			
grape leaf ivy]		
hackberry	cercospora leaf spot powdery mildew		
hawthorn	fabraea leaf spot powdery mildew rust scab		
hibiscus	powdery mildew		
holly	powdery mildew		
hollyhock	powdery mildew rust		
honeysuckle	cercospora leaf spot powdery mildew		
hydrangea	cercospora leaf spot		
iris	didymellina leaf spot rust	Apply 12 fl oz per 100 gallons of spray solution.	
juniper	rust		
leucothoe	cercospora leaf spot		

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
leyland cyprus	cercospora leaf spot		
lilac	powdery mildew		
loblolly pine	fusiform rust	Refer to Douglas fir	
locust	powdery mildew		
maple			Do not use treated trees for syrup production. Do not apply to abutilon (flowering maple).
marigold	cercospora leaf spot rust		
mock-orange	powdery mildew rust		
moonflower	rust		
mountain laurel	cercospora leaf spot ovulinia petal blight powdery mildew	Refer to azalea	
nephthytis	cephalosporium leaf spot		
ninebark	rust		
oak	powdery mildew		
pansy	powdery mildew rust		
pear, flowering	powdery mildew rust scab		

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
petunia	powdery mildew rust		
phlox	cercospora leaf spot powdery mildew rust		
photinia	entomosporium leaf spot powdery mildew rust		
poinsettia	poinsettia scab powdery mildew		
poplar	rust		
potentilla			
privet	cercospora leaf spot powdery mildew		
pyracantha (firethorn)	fusicladium scab		
quince, flowering	blossom and twig blight cercospora leaf spot fabraea leaf spot rust		
rhododendron	cercospora leaf spot ovulinia petal blight powdery mildew	Refer to azalea	

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
rose	black spot powdery mildew rust	Apply on a 7- to 10-day protectant schedule. In areas where black spot is not a problem, spray intervals may be increased to a maximum of 14 days. Greenhouse rose varieties vary in their sensitivity to Eagle 20EW. User should evaluate for possible abnormal response by treating a limited number of plants, at specified rates, prior to initiating large-scale use.	
Russian olive	cercospora leaf spot rust		
salvia	powdery mildew rust		
sedum	powdery mildew		
slash pine	fusiform rust	Refer to Douglas fir	
smoke-tree (cotinus)	cercospora leaf spot rust		
snapdragon	powdery mildew rust		
spirea	powdery mildew		

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
sunflower	cercospora leaf spot powdery mildew rust		Seeds from treated plants may not be used for food or feed.
sycamore	powdery mildew		
trumpet creeper	cercospora leaf blight powdery mildew		
viburnum	powdery mildew rust		
walnut	powdery mildew		Do not use nuts from treated trees for food purposes.
willow			
zinnia	cercospora leaf spot powdery mildew		

Home Orchards, Vineyards, or Fruit Trees

Best control of labeled diseases is achieved when Eagle 20EW is applied on a 7- to 10-day protectant schedule. Eagle 20EW is a systemic fungicide and does not redistribute after application. Adjust application equipment spray nozzles to apply a uniform spray throughout the entire tree canopy.

Dilute (thorough coverage) applications are based upon the amount of spray solution required to thoroughly wet plants to the point of run-off. Refer to use directions for specific tree fruits and vines to determine actual use rate per 100 gallons of spray for control of labeled diseases. The following specific use directions are based on a dilute spray volume of 300 gallons per acre.

Apple

Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
powdery mildew (<i>Podosphaera</i> spp.)	4 - 6	Begin application at tight cluster and continue through the second cover spray. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. Use high rate if powdery mildew was present in previous years.	Preharvest Interval: Do not apply within 14 days of harvest. O not apply more than 153 fl oz of Eagle 20EW (2 lb myclobutanil) per acre per season.
rusts (Gymnospor- angium spp.)		Begin applications at pink stage and continue through the second cover spray.	

Apple (Cont.)

Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
scab (<i>Venturia</i> spp.) prebloom	4 - 6	Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply Eagle 20EW alone or tank mixed with a protectant fungicide on a 7- to 10-day schedule.	Preharvest Interval: Do not apply within 14 days of harvest. On ont apply more than 153 floz of Eagle 20EW (2 lb myclobutanil) per acre per season.
bloom, postbloom		Use Eagle 20EW in a tank mixture with the specified rate of a protectant fungicide, registered for use on apples, for improved fruit scab and summer disease control.	
post-infection	6	Eagle 20EW provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

Grape

Thorough spray coverage is essential for good disease control. Apply Eagle 20EW in sufficient spray volume to ensure complete and uniform coverage.

Diseases	Eagle 20EW (fl oz/acre)	Directions	Restrictions
anthracnose (<i>Elsinoe</i> spp.)	6 - 10	Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days.	 Preharvest Interval: Do not apply within 14 days of harvest. Do not apply
black rot (<i>Guignardia</i> spp.)		Preventative Schedule: Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days. Use a higher rate under heavy disease pressure. Post-infection Schedule: Apply within 72 hours after the beginning of an infection period.	more than 46 fl oz of Eagle 20EW (0.6 lb myclobutanil) per acre per year.
powdery mildew (<i>Uncinula</i> spp.)		Begin application at prebloom (12- to 18-inch shoots) and do not extend applications beyond a 21-day interval. Use a higher rate or shorter spray interval on susceptible varieties or under heavy disease pressure.	

Stone Fruits

Crops	Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
apricot	brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.
	brown rot (Monilinia spp.)		Apply 12 fl oz (0.16 lb myclobutanil) per acre on a 7- to 14-day protectant schedule. Apply when environmental conditions favor disease development during the month prior to harvest.	

Stone Fruits (Cont.)

Crops	Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
apricot (Cont.)	powdery mildew (<i>Podosphaera</i> spp.)	2 - 3	Follow brown rot blossom blight schedule. Reapply at 10- to 14-day intervals until terminal growth ceases.	 Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.
	shothole (Stigmina spp.)		Follow brown rot blossom blight schedule. Reapply at 7- to 10-day intervals as long as needed.	
cherries	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at early popcorn stage, before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	 Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	

Crops	Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
cherries (Cont.)	powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)	2 - 3	Refer to apricot	• Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb
	leaf spot (Blumeriella spp.)		Follow the brown rot blossom blight schedule. Reapply at 7- to 10-day intervals. Make additional applications after harvest.	myclobutanil) per acre per season. • Applications may be made up to the day of harvest.
nectarine	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	

Crops	Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
nectarine (Cont.)	powdery mildew (Podosphaera and Sphaerotheca spp.)	2 - 3	Refer to apricot	• Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb
	shothole (Stigmina spp.)		Follow brown rot blossom blight schedule, Reapply at 7- to 10-day intervals as long as needed.	 myclobutanil) per acre per season. Applications may be made up to the day of harvest.
peach	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
	powdery mildew (<i>Podosphaera</i> spp.)			

Crops	Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
peach (Cont.)	rust (<i>Tranzschelia</i> spp.)	2-3	Apply 12 fl oz (0.16 lb myclobutanil) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on a protectant schedule exceeding 21 days.	 Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.

Crops	Diseases	Eagle 20EW (fl oz/ 100 gallons)	Directions	Restrictions
plum prune	brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at green tip before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.
	rust (<i>Tranzschelia</i> spp.)		Refer to peach	

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. In no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

^{TM®}Trademarks of Corteva Agriscience and its affiliated companies EPA accepted 02/09/11

NOTES

NOTES



CORTEVA

Eagle[®] 20EW

SPECIALTY FUNGICIDE

A systemic, protectant and curative fungicide for disease control in established turfgrass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruits and grapes

Group	3	FUNGICIDE				
Active Ingre						
myclobut						
(chloroph triazole-1	enyl)-1E	<u>1</u> -1,2,4, ienitrile 19.7%				
Total						
Contains petroleum distillates						
Contains 1.67 lb of active ingredient						
per gallon	_	(D)				

Keep Out of Reach of Children CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard. Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Using

EPA Reg. No. 62719-463 EPA Est. 5905-GA-01


CD02-195-020


^{™®}Trademarks of Corteva Agriscience and its affiliated companies


Produced for Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268

NET CONTENTS 1 PT













Classified for "RESTRICTED USE" in New York State under 6NYCRR Part 326 ECICIE® 20EW

ACCEPTED VIA NOTIFICATION LABEL NOT REVIEWED



New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

SPECIALTY FUNGICIDE

A systemic, protectant and curative fungicide for disease control in established turfgrass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruits and grapes

Group	3	FUNGICIDE

Active Ingredient

myclobutanil: a-butyl-a-(chlorophenyl)-	
1H-1,2,4,triazole-1-propanenitrile	%
Other Ingredients	%
Total	
Other Ingredients80.39	%

Contains petroleum distillates Contains 1.67 lb of active ingredient per gallon

Keep Out of Reach of Children CAUTION

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Using

EPA Reg. No. 62719-463

EPA Est. 5905-GA-01

CD02-195-020

^{TM®}Trademarks of Corteva Agriscience and its affiliated companies

Produced for Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268

NET CONTENTS 1 GAL

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of

water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **Note to Physician:** This product may pose an

aspiration pneumonia hazard. Contains petroleum distillates.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves made from barrier laminate
- Shoes plus socks

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- · Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE)and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made from any waterproof material
- · Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry area above freezing.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Storage and Disposal (Cont.)

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. or by other procedures allowed by state and local authorities.

General Information

Shake Well Before Using

Eagle® 20EW specialty fungicide is a systemic, protectant and curative fungicide for the control of the disease listed on this label in established turfgrass (including residential and commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), landscape ornamentals, greenhouse and nursery ornamentals, and non-commercial tree fruits and vines, specifically apples, stone fruits and grapes. Optimum disease control is achieved when this product is applied in a regularly scheduled preventive program.

General Use Precautions

Fungicide Resistance Management

Eagle 20EW belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified

as a Group 3 fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Eagle 20EW should be part of a resistance management strategy that includes alternation and/ or tank mixing with fungicides of different modes of action. After two consecutive applications of Eagle 20EW, another myclobutanil product, or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from Eagle 20EW. Apply the alternate products within the intervals specified on the label for Eagle 20EW. Do not apply Eagle 20EW at rates below those specified on the label. If tank mixing, use the full label rate of Eagle 20EW with the full label rates of other products effective on the target pest. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

The following practices can delay development of fungicide/bactericide resistance:

- Avoid the consecutive use of Eagle 20EW or other target site of action Group 3 fungicides/ bactericides that have a similar target site of action on the same pathogens.
- Use tank mixtures or premixes with fungicides/ bactericides from different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern.
- Base fungicide/bactericide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/ bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
- For further information or to report suspected resistance, you may contact your local company representative or by calling 800-258-3033.

Mixing Directions

Be sure sprayer is clean and not contaminated with other materials prior to use. Fill the spray tank 1/4 to 1/2 of the total amount of water required for the load. Start agitation and maintain agitation throughout mixing and application. Add the required amount of Eagle 20EW directly into the spray tank. Complete filling the tank. Always add Eagle 20EW to the spray tank before adding other materials.

Compatibility

Eagle 20EW is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialist prior to use. When an adjuvant is to be used with this product, Corteva Agriscience recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Carefully read, understand and follow label use rates and restrictions. For proper application, determine the size of the area to be treated, the specified label use rate and the gallonage to be applied to the area. Under low disease conditions, minimum label use rates per application can be used. Use maximum label rates and shortened spray schedules for severe or threatening disease conditions. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is recommended prior to use.

Ground Application

Thorough coverage sprays generally result in optimum disease control. Application equipment must be properly calibrated and provide uniform spray coverage.

Handgun or Pressurized Sprayers: For best results when applying this product on a protectant schedule, ensure thorough coverage of all plant parts.

Chemigation Application

Eagle 20EW must be applied on a regular protectant fungicide schedule, not an irrigation schedule. If irrigation cycles are less frequent than the application intervals for Eagle 20EW, ground or handgun applications must supplement chemigation applications to achieve adequate disease control.

Directions for Sprinkler Chemigation: Apply this product only through solid set or hand-move sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

Chemigation Equipment Preparation: The following use directions are to be followed when this product is applied through irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of Eagle 20EW needed to cover the desired area. Mix according to instructions in the Mixing Directions section. Continually agitate the mixture during mixing and application.

Chemigation Equipment Calibration: In order to calibrate the irrigation system and injector to apply the mixture containing Eagle 20EW, determine the following: 1) Determine area covered by sprinkler: 2) Fill injector solution tank with water and adjust flow rate to use the contents over a 10- to 30-minute interval; (3) Determine the amount of Eagle 20EW required for treatment area: 4) Add the required amount of Eagle 20EW into the same quantity of water used to calibrate the injection equipment. Maintain constant solution tank agitation during the injection period. Operate system at normal pressures specified by the manufacturer of the injection equipment and used for the time interval established during calibration. Inject Eagle 20EW at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention. Stop injection equipment after treatment is completed. Continue to operate the system until the solution with Eagle 20EW has cleared the last sprinkler head.

Chemigation Equipment Requirements:

- The system must contain an air gap, an approved backflow prevention device, a functional check valve, vacuum relief valve (including inspection port), and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information or state specific regulations.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- To insure uniform mixing of the fungicide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all backflow prevention devices on the water line.
- The tank holding the fungicide mixture should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector point.

Chemigation Precautions:

- Crop injury, lack of fungicidal effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.
- Public water system means a system for the provision to the public of piped water for human consumption if such system that has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection.

Chemigation Restrictions:

- Chemigation systems connected to public water systems must contain a functional, reducedpressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place with current certification. Specific local regulations may apply and must be followed.
- Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate nontarget areas.
- Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops.
- Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required PPE is worn.
- Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

Uses

Established Turfgrass

Use Eagle 20EW in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Use diagnostic kits, extension experts, or other identification methods when developing disease control strategies.

In non-residential turfgrass (including commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), optimum disease control is achieved when Eagle 20EW is applied in a preventative disease control program at a rate of 1 to 2.4 fl oz per 1000 sq ft. In residential turfgrass, optimum disease control is achieved when Eagle 20EW is applied in a preventative disease control program at a rate of 1.2 fl oz per 1000 sq ft. See the tables below for specific application rates for various diseases.

Apply Eagle 20EW in sufficient water to ensure thorough coverage. For foliar diseases, use approximately 1 gallon of water per 1000 sq ft. Use 2 to 3 gallons of spray solution per 1000 sq ft. Use conditions favorable for high disease development, reduce the spray interval between applications of Eagle 20EW. Under light to moderate disease pressure, apply Eagle 20EW at the low use rate and/ or longer treatment interval. When disease pressure is high or when used as a curative treatment, use higher rates of Eagle 20EW and/or shorter treatment interval unless otherwise specified.

Non-Residential Turfgrass¹

Diseases	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/Timing (Days)	Directions	Restrictions
anthracnose red thread septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	• Do not apply more than 13.8 fl oz of Eagle 20EW per
brown patch		14	Begin applications when conditions are favorable for disease development, but before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore® T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.	1000 sq ft per year. • For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
copper spot zonate leaf spot			Apply when conditions are favorable for disease	
crown rot leaf spot melting-out			development.	
dollar spot	0.5	7	Apply when conditions are favorable for disease development.	
		14	Tank mix with a low label rate of chlorothalonil.	
	1	21 – 28	Tank mix with the label rate of chlorothalonil.	
	1 - 2.4	14 – 28	If using this rate without tank mixing, make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
fusarium blight	1.2 - 2.4	14 - 21	Apply when conditions are favorable for disease development.	

Non-Residential Turfgrass¹ (Cont.)

Diseases	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/Timing (Days)	Directions	Restrictions
fusarium patch (pink snow mold)	1.2 - 2.4	fall - winter	Apply prior to snow cover.	• Do not apply more than 13.8 fl oz of Eagle 20EW per
gray leaf spot	1.2 - 2.4	14	Apply when conditions are favorable for disease development. If using the lower rate, tank mix with a registered contact fungicide at its specified rate.	 1000 sq ft per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft
leaf smuts	1.2		Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	per year (1.95 lb myclobutanil per acre).
necrotic ring spot	1.2 - 2.4	spring: 28	Make applications on a preventative basis in early to mid-spring.	
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant. Apply 2.4 fl oz per 1000 sq ft followed by a second application one month later.	
powdery mildew rusts	1.2	14 - 28	Apply when conditions are favorable for disease development.	
spring dead spot	2.4	fall: 28	Make 1 to 2 applications in the fall before turfgrass dormancy. Make a second application one month later.	

Non-Residential Turfgrass¹ (Cont.)

Diseases	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/Timing (Days)	Directions	Restrictions
summer patch	1.2 - 2.4	14 - 28	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Use at least 2 to 3 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	 Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
take-all patch	2.4	spring/fall: 28	To reduce the severity, make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F, and 1 to 2 spring applications in April and May depending upon local recommendations.	
zoysia large patch		fall: 28	Make applications in fall before turfgrass dormancy.	

¹Including commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens.

Residential Turfgrass

Disease	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/Timing (Days)	Directions	Restrictions
anthracnose red thread septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 13.8 fl oz of
brown patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore® T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.	Eagle 20EW per 1000 sq ft per year. • For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	
crown rot leaf spot melting-out				
dollar spot			Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
fusarium blight			Apply when conditions are favorable for disease development.	
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.	
gray leaf spot		14	Apply when conditions are favorable for disease development.	
leaf smuts		14	Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	

Residential Turfgrass (Cont.)

Disease	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/Timing (Days)	Directions	Restrictions
necrotic ring spot	1.2	spring: 28	Make applications on a preventative basis in early to mid-spring.	 Do not apply more than 13.8 fl oz of
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	Eagle 20EW per 1000 sq ft per year. • For Nassau and Suffolk
powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.	Counties in New York State, do not apply more
summer patch		14	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Use at least 2 to 3 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).

Landscape, Greenhouse and Nursery Ornamentals

Eagle 20EW is a locally systemic fungicide having protectant and curative properties that will translocate to new growth. For best control of labeled diseases, achieve thorough coverage of all plant parts on a protective application schedule. For dilute application sprays (≥100 gallons of spray volume per acre) applied to ornamental plants in greenhouses, field grown plantings or in commercial and residential landscapes, apply Eagle 20EW at the rate of 6 to 12 fl oz per 100 gallons of spray volume on a 10- to 14-day application schedule unless otherwise directed. Use the higher rate under conditions for infection.

For concentrate sprays (<100 gallons of spray volume per acre), apply 8 fl oz per acre on a 10- to 14-day application schedule.

The addition of a non-phytotoxic spray adjuvant will improve spray coverage and fungicidal performance. Maintain treated plants in a vigorous growing condition. Plants under nutritional or water stress will not respond as well to treatment as wellmaintained plants. Overdosage of Eagle 20EW can result in observable foliar greening, thickened leaves, and/or shortened internodes. If this condition is observed, reduce the fungicide use rate but do not extend the application schedule.

Crop Tolerance

Plant tolerances are acceptable in the specific plants listed on this label. It is not possible to evaluate all ornamental plant species or varieties for tolerance to Eagle 20EW. The user should test for possible phytotoxic responses by treating a limited number of plants, at specified use rates, prior to initiating large-scale use.

The effects of spraying Eagle 20EW in combination with plant growth regulators are not fully understood at this time. If the use of a plant growth regulator

is planned in an area being treated, the user should test for possible enhanced growth regulatory effects by treating a small number of plants, at the specified use rates of all products, prior to initiating large-scale use. Since the effectiveness of such products depends upon not just plant species or cultivar but also weather and seasonable differences (e.g., daylight hours), it is recommended that tests be repeated on previously tested varieties as environmental factors change and that observations for growth regulatory responses be made at regular intervals.

Specific Use Directions for Chrysanthemum

Foliar Sprays: Best control is achieved by thorough coverage sprays applied to point of runoff on a protectant application schedule. Use Eagle 20EW at a rate of 8 fl oz per 100 gallons of spray mixture. Do not apply more than 19 fl oz of Eagle 20EW (0.25 lb myclobutanil) per acre per application. Apply on a 10- to 14-day schedule (not to exceed 21 days).

Prestick Dip Treatment: Chrysanthemum cuttings may be treated by a dip procedure prior to planting as follows: Prepare a dip suspension at a concentration equivalent to 8 fl oz of Eagle 20EW per 100 gallons of water. Fully submerge cuttings in the dip suspension until wet throughout (do not submerge cuttings for more than 2 minutes). If cuttings are dipped, this procedure is the first spray under the quarantine program. Dispose of used dip suspension if it becomes contaminated with soil, plant debris or other foreign matter. Dispose of used dip suspension by spraying it onto registered crops (but not onto previously dipped cuttings) after filtering, or in a manner consistent with local, state, and federal guidelines.

Note: All infected plant material must be destroyed if your state is under quarantine directive.

Specific Use Restrictions:

- Do not apply more than 20 fl oz of Eagle 20EW (0.25 lb myclobutanil) per acre per application. On a total volume per acre basis, do not apply more than 333 gallons of spray per acre at the 6 fl oz per 100 gallons rate or 167 gallons per acre at the 12 fl oz per 100 gallons rate per application.
- Do not apply more than 153 fl oz of Eagle 20EW (2 lb myclobutanil) per acre per year.
- Do not use treated plant materials for food or feed.
- Do not apply to landscape, greenhouse and nursery ornamentals in Nassau and Suffolk Counties in New York State.
- Do not apply to carrotwood (Cupaniopsis anacardioides).

Crops	Diseases	Directions	Precautions/ Restrictions
abelia	cercospora leaf spot		
acalypha (copper-leaf)	powdery mildew		
achillea (yarrow)	powdery mildew rust		
African violet	powdery mildew		
ageratum	powdery mildew		
alder	rust		
almond, flowering	blossom blight (<i>Monilinia</i> spp.)	Apply prebloom, 50% bloom and at petal fall.	
amelanchier (juneberry, shadbush)	fabraea leaf spot powdery mildew rust		
amorpha (false indigo)	cercospora leaf spot powdery mildew rust		

Crops (Cont.) Diseases		Directions	Precautions/ Restrictions
anemone	rust		
angelica	cercospora leaf spot rust		
ash	rust		
aster	powdery mildew rust		
Australian pine	diplodia tip blight		
azalea	petal blight (<i>Ovulinia</i> spp.) powdery mildew	Begin applications when flowers start to exhibit color.	
barberry	powdery mildew rust		May cause temporary damage to crimson pigmy and other atropurposis varieties.
begonia	powdery mildew		
bellflower	cercospora leaf spot powdery mildew rust		
birch	rust		
bittersweet	powdery mildew		
buckeye			
buttonbush	cercospora leaf blight powdery mildew rust		
calendula	cercospora leaf spot		
California poppy	powdery mildew		
canna lily	rust		
carnation	powdery mildew rust		
catalpa	cercospora leaf spot powdery mildew		
cherry, flowering	leaf spot powdery mildew		
chestnut, horse	powdery mildew		
China aster	rust		
chokeberry	rust twig and fruit blight		Fruit may not be used for food or feed.

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions	
Christmas trees	rust			
chrysanthemum	ascochyta blight rust white rust			
columbine	rust			
cornflower				
cosmos	powdery mildew			
cottonwood				
crabapple, flowering	powdery mildew rust scab			
crepe-myrtle	powdery mildew			
daffodil	rust			
dahlia	powdery mildew			
delphinium	powdery mildew rust			
dogwood	anthracnose powdery mildew septoria leafspot			
Douglas fir	needle rust	Apply 12 to 18 fl oz per acre starting early spring. Continue applications at 2- to 3-week intervals until the threat of infection has passed. Spray adjuvants must be added to spray solutions to obtain good spray coverage and disease control.		
dianthus	rust			
elm	powdery mildew			
euonymus				
fern	rhizoctonia aerial blight			
fleabane	cercospora leaf spot powdery mildew rust			
four o'clock	rust			
fuchsia				

Crops (Cont.) Diseases		Directions	Precautions/ Restrictions
gaillardia	powdery mildew		
gardenia	rust		
geranium			
gerbera daisy	powdery mildew		
gourd, ornamental			
grape leaf ivy			
hackberry	cercospora leaf spot powdery mildew		
hawthorn	fabraea leaf spot powdery mildew rust scab		
hibiscus	powdery mildew		
holly	powdery mildew		
hollyhock	powdery mildew rust		
honeysuckle	cercospora leaf spot powdery mildew		
hydrangea	cercospora leaf spot		
iris	didymellina leaf spot rust	Apply 12 fl oz per 100 gallons of spray solution.	
juniper	rust		
leucothoe	cercospora leaf spot		
leyland cyprus	cercospora leaf spot		
lilac	powdery mildew		
loblolly pine	fusiform rust	Refer to Douglas fir	
locust	powdery mildew		
maple			Do not use treated trees for syrup production. Do not apply to abutilon (flowering maple).
marigold	cercospora leaf spot rust		
mock-orange	powdery mildew rust		

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
moonflower	rust		
mountain laurel	cercospora leaf spot ovulinia petal blight powdery mildew	Refer to azalea	
nephthytis	cephalosporium leaf spot		
ninebark	rust		
oak	powdery mildew		
pansy	powdery mildew rust		
pear, flowering	powdery mildew rust scab		
petunia	powdery mildew rust		
phlox	cercospora leaf spot powdery mildew rust		
photinia	entomosporium leaf spot powdery mildew rust		
poinsettia	poinsettia scab powdery mildew		
poplar	rust		
potentilla			
privet	cercospora leaf spot powdery mildew		
pyracantha (firethorn)	fusicladium scab		
quince, flowering	blossom and twig blight cercospora leaf spot fabraea leaf spot rust		
rhododendron	cercospora leaf spot ovulinia petal blight powdery mildew	Refer to azalea	

Crops (Cont.)	Diseases	Directions	Precautions/ Restrictions
rose	black spot powdery mildew rust	Apply on a 7- to 10-day protectant schedule. In areas where black spot is not a problem, spray intervals may be increased to a maximum of 14 days. Greenhouse rose varieties vary in their sensitivity to Eagle 20EW. User should evaluate for possible abnormal response by treating a limited number of plants, at specified rates, prior to initiating large-scale use.	
Russian olive	cercospora leaf spot rust		
salvia	powdery mildew rust		
sedum	powdery mildew		
slash pine	fusiform rust	Refer to Douglas fir	
smoke-tree (cotinus)	cercospora leaf spot rust		
snapdragon	powdery mildew rust		
spirea	powdery mildew		
sunflower	cercospora leaf spot powdery mildew rust		Seeds from treated plants may not be used for food or feed.
sycamore	powdery mildew		
trumpet creeper	cercospora leaf blight powdery mildew		
viburnum	powdery mildew rust		
walnut	powdery mildew		Do not use nuts from treated trees for food purposes.
willow			
zinnia	cercospora leaf spot powdery mildew		

Home Orchards, Vineyards, or Fruit Trees

Best control of labeled diseases is achieved when Eagle 20EW is applied on a 7 - to 10-day protectant schedule. Eagle 20EW is a systemic fungicide and does not redistribute after application. Adjust application equipment spray nozzles to apply a uniform spray throughout the entire tree canopy.

Apple

Dilute (thorough coverage) applications are based upon the amount of spray solution required to thoroughly wet plants to the point of run-off. Refer to use directions for specific tree fruits and vines to determine actual use rate per 100 gallons of spray for control of labeled diseases. The following specific use directions are based on a dilute spray volume of 300 gallons per acre.

Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
powdery mildew (<i>Podosphaera</i> spp.)	4 - 6	Begin application at tight cluster and continue through the second cover spray. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. Use high rate if powdery mildew was present in previous years.	 Preharvest Interval: Do not apply within 14 days of harvest. Do not apply more than 153 fl oz of Eagle 20EW (2 lb myclobutanil) per
rusts (Gymnospor- angium spp.)		Begin applications at pink stage and continue through the second cover spray.	acre per season.
scab (<i>Venturia</i> spp.) prebloom		Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply Eagle 20EW alone or tank mixed with a protectant fungicide on a 7- to 10-day schedule.	
bloom, postbloom		Use Eagle 20EW in a tank mixture with the specified rate of a protectant fungicide, registered for use on apples, for improved fruit scab and summer disease control.	
post-infection	6	Eagle 20EW provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

Grape

Thorough spray coverage is essential for good disease control. Apply Eagle 20EW in sufficient spray volume to ensure complete and uniform coverage.

Diseases	Eagle 20EW (fl oz/acre)	Directions	Restrictions
anthracnose (<i>Elsinoe</i> spp.)	6 - 10	Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days.	 Preharvest Interval: Do not apply within 14 days of harvest. Do not apply more
black rot (Guignardia spp.)		Preventative Schedule: Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days. Use a higher rate under heavy disease pressure. Post-infection Schedule: Apply within 72 hours after the beginning of an infection period.	than 46 fl oz of Eagle 20EW (0.6 lb myclobutanil) per acre per year.
powdery mildew (Uncinula spp.)		Begin application at prebloom (12- to 18-inch shoots) and do not extend applications beyond a 21-day interval. Use a higher rate or shorter spray interval on susceptible varieties or under heavy disease pressure.	

Stone Fruits

Crops	Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
apricot	brown rot blossom blight (<i>Monilinia</i> spp.)	2 - 3	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	 Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.
	brown rot (<i>Monilinia</i> spp.)		Apply 12 fl oz (0.16 lb myclobutanil) per acre on a 7 to 14-day protectant schedule. Apply when environmental conditions favor disease development during the month prior to harvest.	

Crops	Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
apricot (Cont.)	powdery mildew (Podosphaera spp.)	2 - 3	Follow brown rot blossom blight schedule. Reapply at 10- to 14-day intervals until terminal growth ceases.	• Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season.
	shothole (S <i>tigmina</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 7- to 10-day intervals as long as needed.	 Applications may be made up to the day of harvest.
cherries	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at early popcorn stage, before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	 Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb myclobutanil) per acre per season. Applications may be made up to the day
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	of harvest.
	powdery mildew (Podosphaera and Sphaerotheca spp.)			
	leaf spot (<i>Blumeriella</i> spp.)		Follow the brown rot blossom blight schedule. Reapply at 7- to 10-day intervals. Make additional applications after harvest.	
nectarine	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
	powdery mildew (Podosphaera and Sphaerotheca spp.)			

Crops	Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
nectarine (Cont.)	shothole (Stigmina spp.)	2 - 3	Follow brown rot blossom blight schedule, Reapply at 7- to 10-day intervals as long as needed.	Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb myclobutanil) per acre per season.
peach	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Applications may be made up to the day of harvest.
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
	powdery mildew (<i>Podosphaera</i> spp.)			
	rust (<i>Tranzschelia</i> spp.)		Apply 12 fl oz (0.16 lb myclobutanil) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on a protectant schedule exceeding 21 days.	
plum prune	brown rot blossom blight (<i>Monilinia</i> spp.)		Begin application at green tip before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	 Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made up to the day
	rust (<i>Tranzschelia</i> spp.)		Refer to peach	of harvest.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. In no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

TM®Trademarks of Corteva Agriscience and its affiliated companies EPA accepted 02/09/11 NOTES





Eagle[®] 20EW

SPECIALTY FUNGICIDE

A systemic, protectant and curative fungicide for disease control in established turfgrass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruits and grapes

Group	3	FUNGICIDE

Active Ingredient

nyciobutanii: a-c	outyi-a-	
(able readers)	111101	

(chiorophenyi)-1 <u>H</u> -1,2,4,	
triazole-1-propanenitrile	19.7%
Other Ingredients	
Total	100.0%

Contains petroleum distillates Contains 1.67 lb of active ingredient per gallon

Keep Out of Reach of Children CAUTION

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard. Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Using

EPA Reg. No. 62719-463 EPA Est. 5905-GA-01


CD02-195-020


™®Trademarks of Corteva Agriscience and its affiliated companies

Produced for Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268

NET CONTENTS 1 GAL







CORTEVA



A systemic, protectant and curative fungicide for disease control in established turfgrass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruits and grapes

Group	3	FUNGICIDE				
Active Ingredient myclobutanii: a-butyl-a-(chlorophenyl)- 1 <u>H</u> -1,2,4,triazole-1-propanenitrile						
Other Ingredients						

Total.

Contains petroleum distillates Contains 1.67 lb of active ingredient per gallon

Keep Out of Reach of Children

CAUTION

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by

mouth to an unconscious person. If on skin or on clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Note to Physician: This product may pose an aspiration pneumoni

hazard. Contains petroleum distillates. Have the product container or label with you when calling a poison

ontrol center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE) WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear:

 Long-sleeved shirt and long pants Chemical-resistant gloves made from barrier laminate

Shoes plus socks
 Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

Long-sleeved shirt and long pants

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 Remove PPE immediately after handling this product. Wash the outside
- of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS

ACCEPTED VIA NOTIFICATION LABEL NOT REVIEWED

Nov. 22, 2022

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

Classified for "RESTRICTED USE" in New York State under 6NYCRR Part 326

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

Agricultural Use Requirements Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal. Pesticide Storage: Store in a cool, dry area above freezing. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Container Handling: Nonrefillable contained Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining content into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain or 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refer to label booklet for Directions for Use. Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened In case of emergency endangering health or the environment involving this product, call Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing. Shake Well Before Using EPA Reg. No. 62719-463

EPA Est. 62719-ITA-001

CD02-195-020

TM®Trademarks of Corteva Agriscience and its

affiliated companies Produced for Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268

NET CONTENTS 52.835 GAL/200 L

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (MYCLOBUTÁNIL)

UN3082

NOT REGULATED FOR DOMESTIC SURFACE TRANSPORTATION WITHIN THE U.S.A.



Eagle[®] 20EW

SPECIALTY FUNGICIDE

A systemic, protectant and curative fungicide for disease control in established turfgrass, landscape ornamentals, greenhouse and nurserv ornamentals, apples, stone fruits and grapes

Group 3 FUNGICIDE

Active Ingredient myclobutanil: a-butyl-a-(chlorophenyl)-1H-1.2.4.triazole-1propanenitrile 19.7% Other Ingredients 80.3% 100.0% Contains petroleum distillates

Contains 1.67 lb of active ingredient per gallor

Agricultural Use Requirements

Jse this product only in accordance with its labeling and with the Norker Protection Standard, 40 CFR Part 170. Refer to the label booklet under "Agricultural Use Requirements" in the Directions for Use ection for information about this standard.

Refer to inside of label booklet for additional precautionary information including Directions for Use.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are inacceptable, return at once unopened

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. Agricultural Chemical: Do not ship or store with food, feeds, drugs or

Shake Well Before Using

Agricultural Use Requirements

Jse this product only in accordance with its labeling and with the Norker Protection Standard, 40 CFR Part 170. This Standard contain equirements for the protection of agricultural workers on farms, fore nurseries, and greenhouses, and handlers of agricultural pesticides. t contains requirements for training, decontamination, notification, ind emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE)and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: Coveralls

Chemical-resistant gloves made from any waterproof material Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural s used to produce agricultural plants on farms, forests, nurseries of

Keep unprotected persons out of treated area until sprays have dried

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store in a cool, dry area above freezing. Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility

Nonrefillable containers 5 gallons or less: Container Handling: Nonrefillable container. Do not reuse or refill this ontaine

Friple rinse or pressure rinse container (or equivalent) promptly after amptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after

Storage and Disposal (Cont.) he flow begins to drip. Fill the container 1/4 full with water and ecap. Shake for 10 seconds. Pour rinsate into application equip or a mix tank or store rinsate for later use or disposal. Drain for

In a mix tail to be the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents nto application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and drips at show the down of the seconds. Drain for 10 seconds after the down of the seconds of the seconds of the container, and disposal. Insert pressure rinsing nozzle in the side of the container, and the second seconds of the seconds. inse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities

Refillable containers 5 gallons or larger: Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. leaning the container before final disposal is the responsibility of Creating the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate and, in possible, spray an sides write adding water. In practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local with other sets. uthorities.

ocal authorities

General Information

General Use Precautions Fungicide Resistance Management

Shake Well Before Using

Nonrefillable containers 5 gallons or larger: Container Handling: Nonrefillable container. Do not reuse or refill this

Friple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with application equipment of a mix tank. Fin the container of 4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this rocedure two more times. Pressure rinse as follows: Empty the

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

First Aid

f in eyes: Hold eye open and rinse slowly and gently with water for 5-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. Call a poison control center or loctor for treatment advice.

If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person. If on skin or on clothing: Take off contaminated clothing. Rinse skin

Eagle[®] 20EW specialty fungicide is a systemic, protectant and curative fungicide for the control of the diseases listed on this label in established turfgrass (including residential and commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), landscape immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **Note to Physician:** This product may pose an aspiration pneumonia hazard. Contains petroleum distillates.

Have the product container or label with you when calling a poison

control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Causes Moderate Eye Irritation • Harmful If Swallowed Or Absorbed Through Skin

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE) WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) muter uncorr 170) must wear:

- Long-sleeved shirt and long pants Chemical-resistant gloves made from barrier laminate
 Shoes plus socks
- Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Storage and Disposal (Cont.) emaining contents into application equipment or a mix tank and ontinue to drain for 10 seconds after the flow begins to drip. Hold sontainer upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle n the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary andfill, or by incineration, or by other procedures allowed by state and
 - - either automatically or manually shut down.
 - The system must contain functional interlocking controls to nutomatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is
 - Systems must use a metering pump, such as a positive displacement
 - pesticide label-prescribed safety devices for public water systems are in place with current certification. Specific local regulations may apply

 - livestock, wells, or adjoining crops. Do not enter treated area during the reentry interval specified in the Agricultural Use Requirements section of this label unless required F
 - Do not apply through sprinkler systems that deliver a low coefficient of

Non-Residential Turfgrass

Diseases	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
anthracnose red thread septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft per year.
brown patch		14	Begin applications when conditions are favorable for disease development, but before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore® T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.	 For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.	
crown rot leaf spot melting-out				
dollar spot	0.5	7	Apply when conditions are favorable for disease development.	
		14	Tank mix with a low label rate of chlorothalonil.	
	1	21 – 28	Tank mix with the label rate of chlorothalonil.	
	1 – 2.4	14 – 28	If using this rate without tank mixing, make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.	
fusarium blight	1.2 - 2.4	14 - 21	Apply when conditions are favorable for disease development.	
fusarium patch (pink snow mold)		fall - winter	Apply prior to snow cover.	

Diseases	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
gray leaf spot	1.2 - 2.4	14	14 Apply when conditions are favorable for disease development. If using the lower rate, tank mix with a registered contact fungicide at its specified rate. • Do not apply 13.8 floz of 1000 sq ft p	
leaf smuts	1.2		Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.	Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sg ft per year
necrotic ring spot	1.2 - 2.4	spring: 28	Make applications on a preventative basis in early to mid-spring.	(1.95 lb myclobutanil per acre).
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant. Apply 2.4 fl oz per 1000 sq ft followed by a second application one month later.	
powdery mildew rusts	1.2	14 - 28	Apply when conditions are favorable for disease development.	
spring dead spot	2.4	fall: 28	Make 1 to 2 applications in the fall before turfgrass dormancy. Make a second application one month later.	
summer patch	1.2 - 2.4	14 - 28	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Use at least 2 to 3 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	
take-all patch	2.4	spring/fall: 28	To reduce the severity, make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F, and 1 to 2 spring applications in April and May depending upon local recommendations.	1
zoysia large patch] [fall: 28	Make applications in fall before turfgrass dormancy.	

¹Including commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee

Residential Turforass

Disease	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/ Timing (Days)	Directions Restriction				
anthracnose red thread septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 13.8 fl oz of Eagle 20EW pe 1000 sq ft per year.			
brown patch		14	Begin applications when conditions are favorable for disease development and before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore® T/O fungicide. Under conditions of high temperature and humidity, use the shorter spray interval.	 For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acreed) 			
copper spot zonate leaf spot			Apply when conditions are favorable for disease development.				
crown rot leaf spot melting-out							
dollar spot			Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for dollar spot control before rotating to a registered fungicide with a different mode of action.				
fusarium blight			Apply when conditions are favorable for disease development.				
fusarium patch (pink snow mold)] [fall - winter	Apply prior to snow cover.				
gray leaf spot		14	Apply when conditions are favorable for disease development.				
leaf smuts		14	Apply in the fall after turfgrass enters dormancy and/or in the spring prior to the initiation of growth.				

Ground Application

Chemigation Application

spray coverage

plant parts.

Thorough coverage sprays generally result in optimum disease control.

Application equipment must be properly calibrated and provide uniform

Handgun or Pressurized Sprayers: For best results when applying this product on a protectant schedule, ensure thorough coverage of all

Eagle 20EW must be applied on a regular protectant fungicide schedule. not an *irrigation* schedule. If irrigation cycles are less frequent than the application intervals for Eagle 20EW, ground or handgun applications must supplement chemigation applications to achieve adequate disease

Directions for Sprinkler Chemigation: Apply this product only through

Chemigation Equipment Preparation: The following use directions are to be followed when this product is applied through irrigation systems. Thoroughly clean the chemigation system and tank of any fertilizer or

federal laws. Flush the injection system with soap or a cleaning agent and water. Determine the amount of Eagle 20EW needed to cover the desired

Chemigation Equipment Calibration: In order to calibrate the irrigation

solution tank with water and adjust flow rate to use the contents over a

constant solution tank agitation during the injection period. Operate

system and injector to apply the mixture containing Eagle 20EW, determine the following: 1) Determine area covered by sprinkler; 2) Fill injector

10- to 30-minute interval; (3) Determine the amount of Eagle 20EW required for treatment area; 4) Add the required amount of Eagle 20EW into the same quantity of water used to calibrate the injection equipment. Maintain

system at normal pressures specified by the manufacturer of the injection equipment and used for the time interval established during calibration

Inject Eagle 20EW at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention. Stop injection equipment after treatment is completed. Continue to operate the system until the solution with Eagle 20EW has cleared the last sprinkler head.

The system must contain an air gap, an approved backflow prevention device, a functional check valve, vacuum relief valve (including

nspection port), and low-pressure drain appropriately located on

the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the

The pesticide injection pipeline must also contain a functional, normally

closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to

automatically shut off the pesticide injection pump when the water

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

To insure uniform mixing of the fungicide in the water line, inject the mixture in the center of the pipe diameter or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. The injection point must be located after all backflow prevention devices on the water line.

The tank holding the fungicide mixture should be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer

Crop injury, lack of fungicidal effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service

piped water system means a system for the provision to the public to piped water for human consumption if such system that has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall operate the system and make necessary adjustments should the need arise and continuously monitor the injection.

situated between the tank and the injector point.

specialists, equipment manufacturers or other experts. Public water system means a system for the provision to the public of

or state spe

Chemigation Equipment Requirements:

njection pump.

pump motor stops.

Chemigation Precautions:

Engineering Practice 409 for more information

chemical residues, and dispose of the residues according to state and

area. Mix according to instructions in the Mixing Directions section. Continually agitate the mixture during mixing and application.

solid set or hand-move sprinkler irrigation systems. Do not apply this

product through any other type of irrigation system.

- **Chemigation Restrictions:** Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow of the reservoir tank of at least twice the inside diameter of the
- fill pipe. The pesticide injection pipeline must contain a functional, automatic, guick-closing check valve to prevent the flow of fluid back toward the
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is
- adversely affected.
- Systems must use a mereining pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
 Do not connect an irrigation system used for pesticide application (including greenhouse systems) to a public water system unless the capable belows the displacement bulk as the capable of the system value of the sys
- and must be followed.
- Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they
- rrigate nontarget areas. Do not allow irrigation water to collect or run off and pose a hazard to

™®Trademarks of Corteva Agriscience and its affiliated companies Produced for Corteva Agriscience LLC 9330 Zionsville Road apolis, IN 46268

Keep Out

Children

of Reach of

CAUTION

EPA Est. 62719-ITA-001

CD02-195-020

EPA Reg. No. 62719-463

Long-sleeved shirt and long pants

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations Jsers should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the
- outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent vith its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. The following practices can delay development of fungicide/bactericide

ornamentals, greenhouse and nursery ornamentals, and non-commercial Optimum disease control is achieved when this product is applied in a regularly scheduled preventive program.

Fungicide resistance management Eagle 20EW belongs to the steroid demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Eagle 20EW should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action.

After two consecutive applications of Eagle 20EW, another myclobutanil product, or another DMI, rotate to a product that is effective on the target pathogen and has a mode of action different from Eagle 20EW. Apply the alternate products within the intervals specified on the label for Eagle 20EW. Do not apply Eagle 20EW at rates below those specified on the label. If tank mixing, use the full label rate of Eagle 20EW with the full label rates of drar products effective on the target peet.

rates of other products effective on the target pest. Consult your local or

state agricultural authorities for resistance management strategies that are

- sistance: Avoid the consecutive use of Eagle 20EW or other target site of action Group 3 fungicides/bactericides that have a similar target site of action on the same pathogens. Use tank mixtures or premixes with fungicides/bactericides from
- different target site of action groups as long as the involved products are all registered for the same use and are both effective at the tank mix or prepack rate on the pathogen(s) of concern. Base fungicide/bactericide use on a comprehensive Integrated Pest
- Management (IPM) program.
- Monitor treated fungal/bacterial populations for loss of field efficacy Monitor treater unique valuerral populations for nois of neid entracty.
 Contact your local extension specialist, certified crop advisors, and/or manufacturer for fungicide/bactericide resistance management and/or IPM recommendations for specific crops and resistant pathogens.
 For further information or to report suspected resistance, you may
- contact your local company representative or by calling 800-258-3033.

Mixing Directions

Be sure sprayer is clean and not contaminated with other materials prior to use. Fill the spray tank 1/4 to 1/2 of the total amount of water required for the load. Start agitation and maintain agitation throughout mixing and application. Add the required amount of Eagle 20EW directly into the spray tank. Complete filling the tank. Always add Eagle 20EW to the spray tank before adding other materials.

Compatibility Eagle 20EW is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, consult spray compatibility charts or State Cooperative Extension Service Specialist prior to use. When an adjuvant is to be used with this product, Corteva Agriscience recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Application Directions

Carefully read, understand and follow label use rates and restrictions. For proper application, determine the size of the area to be treated, the specified label use rate and the gallonage to be applied to the area. Under low disease conditions, minimum label use rates per application can be used. Use maximum label rates and shortened spray schedules for severe or threatening disease conditions. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is recommended prior to use.

Uses

Established Turfgrass

Use Eagle 20EW in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Use diagnostic kits, extension experts, or other identification methods when developing disease control strategies.

In non-residential turfgrass (including commercial lawns, ornamental turfgrass, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), optimum disease control is achieved when Fagle 20FW is applied in a preventative disease control program at a rate of 1 to 2.4 fl oz per 1000 sq ft. In residential turfgrass, optimum disease control is achieved when Eagle 20EW is applied in a preventative disease control program at a rate of 1.2 fl oz per 1000 sq ft. See the tables below for specific application rates for various

Apply Eagle 20EW in sufficient water to ensure thorough coverage. For Foliar diseases, use approximately 1 gallon of water per 1000 sq ft. Use 2 to 3 gallons of spray solution per 1000 sq ft to control diseases causing root and crown rots. Under conditions favorable for high disease development, reduce the spray interval between applications of Eagle 20EW. Under light to moderate disease pressure, apply Eagle 20EW at the low use rate and/or longer treatment interval. When disease pressure is high or when used as a curative treatment, use higher rates of Eagle 20EW and/or shorter treatment interval unless otherwise specified.

Disease	Eagle 20EW (fl oz/1000 sq ft)	Application Interval/ Timing (Days)	Directions	Restrictions
necrotic ring spot		spring: 28	Make applications on a preventative basis in early to mid-spring.	Do not apply more than 13.8 fl oz of Eagle 20EW per
		fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	1000 sq ft per year.For Nassau and Suffolk Counties in New York State,
powdery mildew rusts		14 - 28	Apply when conditions are favorable for disease development.	do not apply more than 3.43 fl oz of Eagle 20EW
summer patch		14	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending upon recommendations from local turfgrass extension experts. Use at least 2 to 3 gallons of water per 1000 sq ft to increase spray penetration to crown and roots.	per 1000 sq ft per year (1.95 lb myclobutanil per acre).

Landscape, Greenhouse and Nursery Ornamentals

Eagle 20EW is a locally systemic fungicide having protectant and curative properties that will translocate to new growth. For best control of labeled diseases, achieve thorough coverage of all plant parts on a protective application schedule. For dilute application sprays (\geq 100 gallons of spray volume per acre) applied to ornamental plants in greenhouses, field grown plantings or in commercial and residential landscapes, apply Eagle 20EW at the rate of 6 to 12 f oz per 100 gallons of spray volume on a 10- to 14-day application schedule unless otherwise directed. Use the higher rate under conditions for infection.

For concentrate sprays (<100 gallons of spray volume per acre), apply 8 fl oz per acre on a 10- to 14-day application schedule

The addition of a non-phytotoxic spray adjuvant will improve spray coverage and fungicidal performance. Maintain treated plants in a vigorous growing condition. Plants under nutritional or water stress will not respond as well to treatment as well-maintained plants. Overdosage of Eagle 20EW can result in observable foliar greening, thickened leaves, and/or shortened internodes. If this condition is observed, reduce the fungicide use rate but do not extend the application schedule.

Crop Tolerance

Plant tolerances are acceptable in the specific plants listed on this label. It is not possible to evaluate all ornamental plant species or varieties for tolerance to Eagle 20EW. The user should test for possible phytotoxic responses by treating a limited number of plants, at specified use rates, prior to initiating large-scale use

The effects of spraying Eagle 20EW in combination with plant growth regulators are not fully understood at this time. If the use of a plant growth regulator is planned in an area being treated, the user should test for possible enhanced growth regulatory effects by treating a small number of p at the specified use rates of all products, prior to initiating large-scale use. Since the effectiveness of such products depends upon not just plant ber of plants, species or cultivar but also weather and seasonable differences (e.g., daylight hours), it is recommended that tests be repeated on previously tested varieties as environmental factors change and that observations for growth regulatory responses be made at regular intervals.

Specific Use Directions for Chrysanthemum Foliar Sprays: Best control is achieved by thorough coverage sprays applied to point of runoff on a protectant application schedule. Use Eagle 20EW at a rate of 8 fl oz per 100 gallons of spray mixture. Do not apply more than 19 fl oz of Eagle 20EW (0.25 lb myclobutanil) per acre per application. Apply on a 10- to 14-day schedule (not to exceed 21 days).

Prestick Dip Treatment: Chrysanthemum cuttings may be treated by a dip procedure prior to planting as follows: Prepare a dip suspension at a concentration equivalent to 8 fl oz of Eagle 20EW per 100 gallons of water. Fully submerge cuttings in the dip suspension until wet throughout (do not submerge cuttings for more than 2 minutes). If cuttings are dipped, this procedure is the first spray under the quarantine program. Dispose of used dip suspension if it becomes contaminated with soil, plant debris or other foreign matter. Dispose of used dip suspension by spraying it onto registered crops (but not onto previously dipped cuttings) after filtering, or in a manner consistent with local, state, and federal guidelines. $\label{eq:Note: All infected plant material must be destroyed if your state is under quarantine directive.$

Specific Use Restrictions:
Do not apply more than 20 fl oz of Eagle 20EW (0.25 lb myclobutanil) per acre per application. On a total volume per acre basis, do not apply more than 333 gallons of spray per acre at the 6 fl oz per 100 gallons rate or 167 gallons per acre at the 12 fl oz per 100 gallons rate per application.
Do not apply more than 153 fl oz of Eagle 20EW (2 lb myclobutanil) per acre per year.
Do not use treated plant materials for food or feed.

Do not apply to landscape, greenhouse and nursery ornamentals in Nassau and Suffolk Counties in New York State.
Do not apply to carrotwood (*Cupaniopsis anacardioides*).

Crops	Diseases	Directions	Precautions/Restrictions
abelia	cercospora leaf spot		
acalypha (copper-leaf)	powdery mildew		
achillea (yarrow)	powdery mildew rust		
African violet	powdery mildew		
ageratum	powdery mildew		
alder	rust		
almond, flowering	blossom blight (Monilinia spp.)	Apply prebloom, 50% bloom and at petal fall.	
amelanchier (juneberry, shadbush)	fabraea leaf spot powdery mildew rust		
amorpha (false indigo)	cercospora leaf spot powdery mildew rust		
anemone	rust		
angelica	cercospora leaf spot rust		

Crops (Cont.)	Diseases	Directions	Precautions/Restrictions				Eagle		
marigold	cercospora leaf spot rust				rops	Diseases	20EW (fl oz/100 gallons)	Directions	Restriction
mock-orange	powdery mildew rust				erries	brown rot blossom	2 - 3	Begin application at early popcorn	Do not app more than
moonflower	rust			1		blight		stage, before	100 fl oz
mountain laurel	cercospora leaf spot ovulinia petal blight powdery mildew	Refer to azalea				(Monilinia spp.)		infection occurs. If conditions are favorable for disease	of Eagle 20EW (1.3 I myclobutar per acre pe
nephthytis	cephalosporium leaf spot			1				development,	season.
ninebark	rust			1				reapply at full bloom and	 Applications may be made
oak	powdery mildew]				petal fall.	up to the da
pansy	powdery mildew rust					brown rot (Monilinia		Refer to apricot	of harvest.
pear, flowering	powdery mildew rust scab					spp.) powdery mildew			
petunia	powdery mildew rust					(Podosphaera and Sphaerotheca			
phlox	cercospora leaf spot powdery mildew rust					spp.) leaf spot (Blumeriella		Follow the brown rot blossom	_
photinia	entomosporium leaf spot powdery mildew rust					spp.)		blight schedule. Reapply at 7- to 10-day intervals. Make additional	
poinsettia	poinsettia scab powdery mildew]				applications after harvest.	
poplar	rust			nec	nectarine			Begin application	
potentilla	1			1		blossom blight		at early pink bud stage before	
privet	cercospora leaf spot powdery mildew					(<i>Monilinia</i> spp.)		infection occurs.	
pyracantha (firethorn)	fusicladium scab			1		5	CPP-7		are favorable
quince, flowering	blossom and twig blight cercospora leaf spot fabraea leaf spot rust							for disease development, reapply at full bloom and petal fall.	

Precautions/Restrictions

Seeds from treated plants may not be used for food or feed.

Crops (Cont.)	Diseases	Directions	Precautions/Restrictions		Crops (Cont.)	Diseases	Directions
ash	rust			rh	ododendron	cercospora leaf spot	Refer to azalea
aster	powdery mildew rust			1 L		ovulinia petal blight powdery mildew	
Australian pine	diplodia tip blight			ro	rose	black spot powdery mildew	Apply on a 7- to 10-day protectant schedule. In areas where black
azalea	petal blight (Ovulinia spp.) powdery mildew	Begin applications when flowers start to exhibit color.				rust	spot is not a problem, spray intervals may be increased to a
barberry	powdery mildew rust		May cause temporary damage to crimson pigmy and other atropurposis varieties.				maximum of 14 days. Greenhouse rose varieties vary in their sensitivity to Eagle 20EW. User should evaluate for possible
begonia	powdery mildew						abnormal response by treating
bellflower	cercospora leaf spot powdery mildew rust						a limited number of plants, at specified rates, prior to initiating large-scale use.
birch	rust			Ru	Russian olive salvia	cercospora leaf spot	
bittersweet	powdery mildew					rust	
buckeye				sa		powdery mildew rust	
buttonbush	cercospora leaf blight powdery mildew			se	dum	powdery mildew	
	rust			sla	ash pine	fusiform rust	Refer to Douglas fir
calendula	cercospora leaf spot			sn	noke-tree (cotinus)	cercospora leaf spot	
California poppy	powdery mildew					rust	
canna lily	rust			sn	apdragon	powdery mildew rust	
carnation	powdery mildew rust			sp	irea	powdery mildew	
catalpa	cercospora leaf spot powdery mildew			su	nflower	cercospora leaf spot powdery mildew rust	
cherry, flowering	leaf spot powdery mildew			sy	sycamore	powdery mildew	
chestnut, horse	powdery mildew			- tru	Impet creeper	cercospora leaf blight	
China aster	rust			\dashv		powdery mildew	
chokeberry	rust		Fruit may not be used for food or	- vit	viburnum	powdery mildew	
CHORODOTTY	twig and fruit blight		feed.			rust	
Christmas trees	rust			7			

Crops	Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
nectarine	brown rot (<i>Monilinia</i> spp.)	2 - 3	Refer to apricot	 Do not apply more than 100 fl oz
	powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)			of Eagle 20EW (1.3 lb myclobutanil) per acre per season. • Applications may be made
	shothole (Stigmina spp.)		Follow brown rot blossom blight schedule, Reapply at 7- to 10-day intervals as long as needed.	up to the day of harvest.
peach	brown rot blossom blight <i>(Monilinia</i> spp.)		Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	
	brown rot (<i>Monilinia</i> spp.)		Refer to apricot	
	powdery mildew (<i>Podosphaera</i> spp.)			
	rust (<i>Tranzschelia</i> spp.)		Apply 12 fl oz (0.16 lb myclobutanil) per acre.	

Crops (Cont.)	Diseases	Directions	Precautions/Restrictions
chrysanthemum	ascochyta blight rust white rust		
columbine	rust		
cornflower			
cosmos	powdery mildew		
cottonwood			
crabapple, flowering	powdery mildew rust scab		
crepe-myrtle	powdery mildew		
daffodil	rust		
dahlia	powdery mildew		
delphinium	powdery mildew rust		
dogwood	anthracnose powdery mildew septoria leafspot		
Douglas fir	needle rust	Apply 12 to 18 fl oz per acre starting early spring. Continue applications at 2- to 3-week intervals until the threat of infection has passed. Spray adjuvants must be added to spray solutions to obtain good spray coverage and disease control.	
dianthus	rust		
elm	powdery mildew		
euonymus			
fern	rhizoctonia aerial blight		
fleabane	cercospora leaf spot powdery mildew		

Crops (Cont.)	Diseases	Directions	Precautions/Restrictions
walnut	powdery mildew		Do not use nuts from treated trees for food purposes.
willow			
zinnia	cercospora leaf spot powdery mildew		

Iome Orchards, Vineyards, or Fruit Trees

test control of labeled diseases is achieved when Eagle 20EW is applied on a 7- to 10-day protectant schedule. Eagle 20EW is a systemic fungicide nd does not redistribute after application. Adjust application equipment spray nozzles to apply a uniform spray throughout the entire tree canopy. vilute (thorough coverage) applications are based upon the amount of spray solution required to thoroughly wet plants to the point of run-off. Refer o use directions for specific tree fruits and vines to determine actual use rate per 100 gallons of spray for control of labeled diseases. The following pecific use directions are based on a dilute spray volume of 300 gallons per acre.

Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
powdery mildew (<i>Podosphaera</i> spp.)	4 - 6	Begin application at tight cluster and continue through the second cover spray. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. Use high rate if powdery mildew was present in previous years.	 Preharvest Interval: Do not apply within 14 days of harvest. Do not apply more than 153 fl oz of Eagle 20EW (2 lb
rusts (Gymnospor- angium spp.)		Begin applications at pink stage and continue through the second cover spray.	myclobutanil) per acre per season.
scab (<i>Venturia</i> spp.) prebloom		Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply Eagle 20EW alone or tank mixed with a protectant fungicide on a 7- to 10-day schedule.	
bloom, postbloom		Use Eagle 20EW in a tank mixture with the specified rate of a protectant fungicide, registered for use on apples, for improved fruit scab and summer disease control.	
post-infection	6	Eagle 20EW provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

Crops	Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
peach	rust (Tranzschelia spp.)	2-3	Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not apply on a protectant schedule exceeding 21 days.	 Do not apply more than 100 fl oz of Eagle 20EW (1.3 lb myclobutanil) per acre per season. Applications may be made up to the day of harvest.
plum prune	brown rot blossom blight (Monilinia spp.)	2 - 3	Begin application at green tip before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made
	rust (<i>Tranzschelia</i> spp.)		Refer to peach	up to the day of harvest.

Eaglo

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

st	•			

Crops (Cont.)	Diseases	Directions	Precautions/Restrictions	Grape				
our o'clock	rust			Thorough spray	coverage is ess	ential for good disease co to ensure complete and u	ntrol. Apply Eagl	
iuchsia				20EW IN SUIIICIER	it spray volume	to ensure complete and t	iniiorm coverage.	
jaillardia	powdery mildew				Eagle 20EW			
ardenia	rust			Diseases	(fl oz/acre)	Directions	Restrictions	
geranium				anthracnose (Elsinoe	6 - 10	Begin application when new shoots are 1 to	Do not apply within 4 days of	
erbera daisy	powdery mildew			spp.)		3 inches in length. Reapply on a protectant schedule		
ourd, ornamental								
rape leaf ivy						that does not exceed		
ackberry	cercospora leaf spot powdery mildew			black rot		14 days. • Do not ap more than Preventative more than Schedule: Begin application when new shorts are of Eagle		
nawthorn	fabraea leaf spot powdery mildew rust scab			(Guignardia spp.)			of Eagle 20EW (0.6 lb myclobutanil)	
hibiscus	powdery mildew							
nolly	powdery mildew						-	
ollyhock	powdery mildew rust							
oneysuckle	cercospora leaf spot powdery mildew						pressure.	
nydrangea	cercospora leaf spot							
is	didymellina leaf spot rust	Apply 12 fl oz per 100 gallons of spray solution.						
uniper	rust			powdery		Begin application at	-	
eucothoe	cercospora leaf spot			mildew		prebloom (12- to		
eyland cyprus	cercospora leaf spot			(Uncinula		18-inch shoots)	nd or	
lac	powdery mildew			spp.)	a	and do not extend applications beyond a 21-day interval.		
oblolly pine	fusiform rust	Refer to Douglas fir						
ocust	powdery mildew					Use a higher rate or shorter spray interval		
naple			Do not use treated trees for syrup production. Do not apply to abutilon (flowering maple).			on susceptible varieties or under heavy disease pressure.		

	Stone Fru	its			
Eagle age.	Crops	Diseases	Eagle 20EW (fl oz/100 gallons)	Directions	Restrictions
hin of oply n	apricot	brown rot blossom blight (Monilinia spp.)	2-3	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	 Do not apply more than 84 fl oz of Eagle 20EW (1.1 lb myclobutanil) per acre per season. Applications may be made up to the day
6 lb tanil) per		brown rot (Monilinia spp.)		Apply 12 fl oz (0.16 lb myclobutanii) per acre on a 7- to 14-day protectant schedule. Apply when environmental conditions favor disease development during the month prior to harvest.	of harvest.
		powdery mildew (<i>Podosphaera</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 10- to 14-day intervals until terminal growth ceases.	
		shothole (Stigmina spp.)		Follow brown rot blossom blight schedule. Reapply at 7- to 10-day intervals as long as needed.	

Warranty Disclaimer

Corteva Agriscience warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Corteva Agriscience MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Corteva Agriscience or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Corteva Agriscience's election, one of the following:

Refund of purchase price paid by buyer or user for product bought, or
 Replacement of amount of product used

To the extent permitted by law, Corteva Agriscience shall not be liable for losses or damages resulting from handling or use of this product unless Corteva Agriscience is promptly notified of such loss or damage in writing. In no case shall Corteva Agriscience be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Corteva Agriscience or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.

 ${}^{\rm TM}{}^{\rm TM}$ Trademarks of Corteva Agriscience and its affiliated companies EPA accepted 02/09/11