

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Eagle[®] 20EW

EPA Reg. No. 62719-463

Special 2(ee) Recommendation[†] (For Distribution and Use Only in New York State)

Control of Brown Patch/Large Patch (*Rhizoctonia solani*) in Established Turfgrass

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Eagle[®] 20EW specialty fungicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Eagle 20EW according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Eagle 20EW.

Directions for Use

Refer to product label for Eagle 20EW for Use Precaution, Mixing Directions and Application Directions.

Use Eagle 20EW to control brown patch/large patch (*Rhizoctonia solani*) in established residential and non-residential turfgrass.

In established residential and non-residential turfgrass, apply 1.2 fl oz per 1000 sq ft. Begin applications when conditions are favorable for disease development and before disease symptoms are apparent. If disease is present, mix Eagle 20EW with an EPA registered contact fungicide, such as Fore[®] T/O fungicide. The application interval is 14 days. Do not apply more than 13.8 fl oz of Eagle 20EW per 1000 sq ft per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl oz of Eagle 20EW per 1000 sq ft per year (1.95 lb myclobutanil per acre).

Read the label affixed to the container for Eagle 20EW and follow all use directions, precautions, and restrictions.

[®]Trademark of Dow AgroSciences LLC

R195-003

Issued: 08/17/11

Initial printing.