


ACCEPTED  
FOR REGISTRATION

July 17, 2014

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

DOC ID: 539377


**EVERGREEN®**  
 PYRETHRUM DUST
**ACTIVE INGREDIENTS:**
 \*Pyrethrins ..... 1.00%  
 OTHER INGREDIENTS ..... 99.00%  
 100.00%

 Evergreen® - Registered trademark of McLaughlin  
 Gormley King Company

 Pyrethrins are a botanical insecticide derived from  
 chrysanthemum flowers.

 0.01 lbs. active ingredient per lbs. of Evergreen  
 Pyrethrum Dust

**KEEP OUT OF  
 REACH OF CHILDREN  
 CAUTION**
**FIRST AID**
**IF ON SKIN OR CLOTHING:** Take off contaminated  
 clothing. Rinse skin immediately with plenty of water for  
 15-20 minutes. Call a poison control center or doctor for  
 treatment advice. Have the product container or label  
 with you when calling a poison control center or doc-  
 tor, or going for treatment. For additional information  
 on this pesticide product (including health concerns,  
 medical emergencies or pesticide incidents), you may  
 call 1-800-745-8712.
**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**
 Harmful if absorbed through skin. Avoid contact with  
 skin, eyes or clothing. Remove and wash contaminated  
 clothing before reuse.
**PERSONAL PROTECTIVE EQUIPMENT (PPE)**
 Mixers, Loaders, Applicators, and other handlers  
 must wear the following:

- long-sleeved shirt
- long pants
- shoes and socks
- chemical-resistant gloves made of any waterproof  
 material

**USER SAFETY REQUIREMENTS**
 Follow manufacturer's instructions for cleaning/  
 maintaining PPE. If no such instructions for washables  
 exist, use detergent and hot water. Keep and wash PPE

separately from other laundry.

 Discard clothing and other absorbent material that  
 have been drenched or heavily contaminated with the  
 product's concentrate. Do not reuse them.
**USER SAFETY RECOMMENDATIONS:**
**Users should:** Wash hands before eating, drinking,  
 chewing gum, using tobacco or using the toilet. Remove  
 clothing/PPE immediately if pesticide gets inside. Then  
 wash thoroughly and put on clean clothing. Remove PPE  
 immediately after handling this product. Wash the out-  
 side of gloves before removing. As soon as possible,  
 wash thoroughly and change into clean clothing.
**ENVIRONMENTAL HAZARDS**
 This product is toxic to aquatic organisms, including fish  
 and invertebrates. To protect the environment, do not  
 allow pesticide to enter or runoff into storm drains,  
 drainage ditches, gutters or surface waters. Applying this  
 product in calm weather when rain is not predicted for  
 the next 24 hours will help to ensure that wind or rain  
 does not blow or wash pesticide off the treatment area.  
 This product is highly toxic to bees exposed to direct  
 treatment on blooming crops or weeds. Do not apply this  
 product or allow it to drift to blooming crops or weeds  
 while bees are actively visiting the treatment area.
**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**
 It is a violation of Federal law to use this product in a  
 manner inconsistent with its labeling.

See inside booklet label for full directions for use.

EPA Reg. No. 1021-1871 F7461-0264

EPA Est. Nos. 1021-MN-24, Rev 0214-0414

395378-TX-1<sup>†</sup>

Superscript is the first character of the lot number.

**NET WEIGHT**  
**10 Ounces**

Manufactured by:


 8810 Tenth Avenue North  
 Minneapolis, MN 55427

PULL HERE TO OPEN / PRESS TO RESEAL

PF-48708-3

### FIRST AID

**IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

## **PRECAUTIONARY STATEMENTS** **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

### **CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Mixers, Loaders, Applicators, and other handlers must wear the following:

- long-sleeved shirt
- long pants
- shoes and socks
- chemical-resistant gloves made of any waterproof material

### **USER SAFETY REQUIREMENTS**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

### **USER SAFETY RECOMMENDATIONS:**

**Users should:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

*(continued)*

## **PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL BEFORE EACH USE**

## **USE RESTRICTIONS**

- Apply this product only as specified on the label.
- Aerial applications are prohibited.
- Do not allow contamination of food and feedstuffs.
- In indoor residential areas or other areas where children may come into contact, only apply as a crack-and-crevice or as a spot treatment to areas no greater than 3 feet by 3 feet per room.
- Applications with power duster equipment are prohibited.
- Do not enter or allow others to enter until dusts have settled.
- Do not apply this product in a way that will contact workers or other persons, either directly or indirectly through drift.
- Only protected handlers may enter treated area during application.
- Do not allow adults, children or pets to enter the treated area until dusts have settled.
- For use outdoors, do not apply directly to or near water, storm drains or drainage ditches. Do not apply when windy.
- Do not water the treated area to the point of runoff.
- Do not make applications during rain.
- Do not apply this product in a way that will contact adults, children or pets, either directly or indirectly through drift.
- Remove or cover exposed food and drinking water before application.
- Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use. Do not use in serving areas while food is exposed.
- In Food Areas of Food Handling Establishments: Do not contaminate food or feedstuffs. During applications, cover or remove food. Cover food processing surfaces and equipment or clean after treatment with a suitable detergent and rinse with potable water before use. Do not use in serving areas (dining areas where prepared foods are served) while food is exposed or facility is in operation.
- In Schools, do not apply to classrooms when in use.
- Do not use in aircraft cabins.
- Remove pets, birds and cover fish aquariums before application.
- When used in dairy barns or facilities; close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Except when applying to sewers or drains as instructed on this label, do not apply directly in to sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product any drain during or after application.
- Not for use on commercial sod farms.

- USE RESTRICTIONS IN FOOD HANDLING AREAS:
  - Do not apply when food or feed processing facility is in operation or foods are exposed.
  - APPLICATIONS OF THIS PRODUCT IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS ARE NOT PERMITTED, *OTHER THAN AS A CRACK AND CREVICE TREATMENT.*

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE required for early entry to treated areas (that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Long-sleeved shirt and long pants;
- Shoes plus socks.
- chemical-resistant gloves made of any waterproof material

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until dusts have settled.

### Kills Exposed listed insects:

Adelgids	Alfalfa Weevil	Almond Moth	American Black Flour Beetle
Angoumois Grain Moths	Ants	Aphids	Apple Maggot
Armyworms	Asian Cerambycid Beetles	Asian Lady Beetle	Asian Pine Tip Moths
Bagworm	Barn Flies	Bat Bugs	Beet Beetles
Bean Weevils	Bed Bugs	Bees	Beet Armyworms
Beetles	Birch Leafminer	Black Carpet Beetles	Black Fungus Beetles
Black Turpentine Beetles	Black Vine Weevil	Blister Beetles	Blow Flies
Booklice	Bottle Flies	Box Elder Bugs	Broadhorned Flour Beetles
Broadnosed Grain Weevils	Blacklegged Tick Nymphs	Brown Spider Beetles	Brown Spruce Longhorn Beetle
Budworms	Cadelles	California Oakworm	Cankermolds
Carpet Beetles	Carion Beetles	Carrot Weevil	Caterpillars
Catorama Beetles	Cheese Skippers	Centipedes	Cereal Beetles
Cheese Mites	Chrysanthemum Leaf Miners	Chicken Mites	Chocolate Moths
Cicadas	Cigarette Beetle	Clearwing Borer	Clothes Moths
Clover Weevil	Clover Mites	Cluster Flies	Cocoa Bean Moths
Coffee Bean Weevils	Cone Beetles	Cone Worms	Conifer Spider Mite (Adult)

Confused Flour Beetle	Corn Sap Beetles	Cottonwood Borer	Cottonwood Leaf Beetles
Cranberry Girdlers	Crane Flies	Crickets	Cross-striped Cabbageworm
Cucumber Beetles	Cutworms (exposed)	Dark Mealworms	Darkling Beetles
Deer Tick	Depressed Flour Beetles	Dermeid Beetles	(lesser meal worm)
Diamondback Larvae	Dingy Cutworm Adult	Douglas Fir Midge	Douglas Fir Tussock Moth
Drain Flies	Dried Fruit Beetles	Drugstore Beetles	Earwigs
Eastern Tent Caterpillar	Elmleaf Beetle	Emerald Ash Borer	European Grain Moths
European Pine Shoot Moth	European Pine Tip Moth	Face Flies	Fall Cankerworms
Fall Webworm	Fannia Flies	Field Crickets	Firebrats
Fireworms	Flat Grain Beetles	Flea Beetles	Fleas
Flies	Foreign Grain Beetles	Forest Tent Caterpillar	Flying Moths
Fruit Flies	Fruittree Leafroller	Fungus Beetles	Fungus Gnats
German Cockroaches	Giant Root Aphid	Glassy Winged Sharpshooter	Grain Mites
Granary Insects	Granary Weevils	Grape Leaf Skeletonizer	Grape Leafhopper
Grasshoppers	Green Peach Aphid	Greenbugs	Green Fruit Worm
Greenhouse Thrips	Ground Beetles	Gypsy Moth (adults & larvae)	Hairy Fungus Beetles
Hairy Spider Beetles	Harlequin Bug	Heliothis spp.	Hessian Fly (Adult)
Hide Beetles	Hornworm	Inchworm	Indian Meal Moth
Iris Borer	Japanese Beetle	Katydid	Khapra Beetles
Lace Bugs	Larger Cabinet Beetles	Large Black Flour Beetles	Larger Grain Borers
Leafhoppers	Leafminers	Leafrollers	Leaftiers
Lesser Grain Beetles	Lesser Grain Borers	Lesser Meal Worm	Lesser Peach Tree Borer
Longheaded Flour Beetles	Loopers	Lygus Bugs	Maize Weevils
Mealy Bugs	Mediterranean Flour Moths	Merchant Grain Beetles	Mexican Bean Beetle
Mexican Grain Beetles	Millers	Millipedes	Mimosa Webworm
Mole Crickets	Moth Flies	Mushroom Flies	Murmidius Beetles
Nantucket Pine Tip Moth	Northern Fowl Mites	Onion Maggot	Palmetto Bug
Peach Twig Borer	Pear Psylla	Pillbugs	Pine Bark Beetles
Pine Engraver	Pink Scavenger Caterpillars	Pinworms	Plant Bugs
Plum Curculio	Ponderosa Pine Needle Miner	Poultry Lice	Powder Post Beetles
Psocids	Psyllids	Red Flour Beetles	Red Gum Lerp Psyllid
Red Horned Grain Beetles	Red Lily Beetles	Red Oak Borer	Redbanded Leafroller
Rice Flour Beetles	Rice Moths	Rice Weevils	Root Weevils
Rose Chafer	Royal Palm Bugs	Rusty Grain Beetles	Saw Flies
Saw-Toothed Grain Beetle	Scale (Crawlers)	Seedbugs	Silverfish
Skipper Flies	Skippers	Small Flying Moths	Small Fruit Flies
Smalleyed Flour Beetles	Slenderhorned Flour Beetles	Southern Red Mite (Adult)	Southern Pine Beetles
Sowbugs	Spider Beetles	Spider Mites	Spruce Mite (Adult)
Squarenecked Grain Beetles	Stable Flies	Stalk Borers	Stink Bugs
Stored Product Pests	Sycamore Leaf Beetle	Subtropical Pine Tip Moth	Tarnished Plant Bug
Trogoderma Beetles	Tent Caterpillar	Thrips	Tobacco Moths
Two-Banded Beetles	Warehouse Beetles	Wasps (Yellow Jackets, Hornets, and Paper Wasps)	Waterbugs
Webworms	Weevils	Whiteflies	Western Pine Tip Moth
White Apple Leafroller	White Marked Spider Beetles	Yellow Mealworms	Wireworms
Vegetable Leaf Miners	Vinegar Flies		

**Application Methods:**

Apply Evergreen Pyrethrum Dust using a non-power dust applicator, a shaker can, dust gun, puffer or other bulbous duster, hand pump duster, hand operated rotary duster, static charge duster, or other appropriate hand operated duster.

**INDOOR AREAS:** For use in and around indoor areas:

Animal Areas	Animal Quarters	Apartments
Asian Markets	Attics	Bakeries
Barns	Basements	Beef Barns
Beverage Plants	Bins	Bottling Plants
Brandy Storage Warehouses	Breweries	Brewery Warehouses
Buildings Housing Animals	Buildings Housing Poultry	Cabinets
Cafeterias	Calving Barns	Canneries
Cat Sleeping Quarters	Cattle Barns	Certified Organic Facilities
Cheese Caves	Citrus Packing Houses	Coffee Houses
Coffee Shops	Commercial Buildings	Condominiums
Convenience Stores	Crawl Spaces	Dairy and Livestock Barns
Dairy Food Processing	Dairies	Day Care Centers
Delicatessens	Department Stores	Diners
Distilleries	Dog Houses	Dog Sleeping Quarters
Drains & Sewers	Dried Flower Warehouses	Dried Food Products in Processing
Dried Food Products in Storage	Dried Fruit Processing Plants	Dried Fruit Warehouses
Dumpsters	Dwellings	Egg Processing Plants
Egg Storage Facilities	Factories	False Ceilings
Farm Structures	Feed Mills	Flour Mills
Food Handling Establishments	Food Preparation Facilities	Food Processing Facilities
Food Quarantine Areas	Food Service Facilities	Food Storage Facilities
Freight Containers	Fruit Packing Sheds	Fruit Processing Plants
Fruit Storage Warehouses	Fruit Quarantine Areas	Garbage Bins
Garbage Cans	Garbage Compactors	Garbage Trucks
Goat Holding Areas	Goat Houses	Grain Elevators
Grain Handling Equipment	Grain Harvesting Equipment	Grain Mills
Grain Storage	Granaries	Greenhouses
Grocery Stores	Homes	Hog Barns
Horse Stables	Horse Stables and Barns	Horse Trailers
Hospitals	Hotels	Houses
Indoor Eating Facilities	Industrial Installations	Industrial Plants
Institution Dining Areas	Institutions	Kennels
Liveries	Livestock Barns	Livestock Hauling Equipment
Livestock Holding Areas	Loafing Sheds	Meat Markets
Meat Packaging Plants	Meat Packing Plants	Mess Halls
Milk Houses	Milk Processing Facilities	Milk Rooms
Milk Sheds	Milking Rooms	Mills
Motels	Mushroom Caves	Mushroom Processing Plants
Office Buildings	Other Public Buildings	Packaged Food Storage
Pantries	Peanut Processing Plants	Peanut Warehouses

Pet Bedding	Pet Carriers	Pet Food Plants
Pet Sleeping Areas	Poultry Houses	Poultry Processing Plants
Rabbit Houses	Rabbit Processing Plants	Railroad Cars
Research Animal Areas	Restaurants	Rice Mills
Schools	Seed Storage	Shade Houses
Sheep Holding Areas	Ship Holds	Ships
Spice Markets	Stable Bins	Stables
Storage Areas	Storage Bins (areas)	Storages
Stored Product Areas	Stored Products	Stored Seed Warehouses
Stores	Supermarkets	Swine Houses
Taverns	Tea Houses	Textile Mills
Tobacco Factories	Tobacco Warehouses	Tool Sheds
Trains	Transportation Equipment (except Aircraft)	Tree Nuts Storage Areas
Truck Trailers	Vegetable Packing Facilities	Vegetable Processing Facilities
Vegetable Storage Facilities	Veterinary Clinics	Veterinary Hospitals
Voids in Attics	Wall Voids	Warehouses
Wheat Mills	Wholesale Stores	Wine Cellars
Wine Storage Warehouses	Wine Tasting Rooms	Wineries
Zoos		

#### APPLICATIONS WITHIN FOOD HANDLING ESTABLISHMENTS:

**Food and/or Feed areas** include areas for receiving, storage (dry, cold, frozen or raw), packaging (canning, bottling, wrapping, boxing and grinding), preparing, edible waste storage and enclosed processing systems (mills, dairies, and systems for processing edible oils and syrups). Serving areas are places where prepared foods are served.

FOOD HANDLING ESTABLISHMENTS	
Application Type	Application Rate
Crack and Crevice Food/Feed Areas of Food/Feed Handling Establishments	1.5 oz. / 50 sq. ft.
Crack and Crevice Stored product pests in Non-food/Non-Feed Areas of Food Handling Establishments	1 oz. / 50 sq. ft.
Crack and Crevice treatments in Storage Sites	1.0 oz/ 50 sq. ft.

Using a non-power duster, apply directly into cracks and crevices, in expansion joints between different elements of construction, under or between equipment bases and the floor, beneath appliances, into wall voids, motor housing, junction boxes or switch boxes, conduits, behind and under cabinets or drawers, in and around drains, around drain pipes, plumbing and utility entrances, hollow equipment legs and other sites where listed insects can enter or hide.

#### Stored Product Pests in Non-Food/Non-Feed areas of Food handling Establishments:

Non-food/feed areas of food/feed handling establishments are garbage rooms, lavatories, floor drains (to sewers, entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

To aid in the kill of stored product pests in listed areas. Apply Evergreen Pyrethrum Dust at the specified rate. Clean up waste materials, infested materials, dust, dirt, and any other debris. Thoroughly treat floors, walls, shelves, drawers, cabinets and other surfaces in bins, storage and handling areas, around cartons or containers and other areas where these insects tend to

congregate. Apply into cracks and crevices and other difficult to reach areas where insects may hide. Apply in voids or cracks and crevices in pantries and other infested areas within the structure.

**For Use in Storage Sites:**

This product can be used in warehouse bins, trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment and other listed sites as a cleanup prior to using them for storage. Remove and dispose of all of the infested stored commodity accumulations and debris, with special attention to material lodged in the cracks and crevices. After sanitation measures have been employed, dust areas prior to use for storage, paying particular attention to cracks and crevices.

For packaged food warehouses, treat potential points of insect entry, as well as harborage sites in, on and around wooden and plastic pallets, pallet-racking systems, loading docks, scales, around cartons.

**INDOOR TREATMENTS:**

IN INDOOR RESIDENTIAL AREAS OR OTHER AREAS WHERE CHILDREN MAY COME INTO CONTACT, ONLY APPLY AS A CRACK-AND-CREVICE OR AS A SPOT TREATMENT TO AREAS NO GREATER THAN 3 FEET BY 3 FEET PER ROOM.

**APPLICATION RATES FOR INDOOR TREATMENTS**

<b>Application Type</b>	<b>Application rates for Residential Areas such as homes, condominiums, apartments, hotels, motels and pet areas</b>	<b>Application Rates for Non-Residential Areas such as offices, warehouses, hospitals, and other commercial buildings and institutions.</b>
Crawling Insects—Crack and Crevice	Apply as a crack-and-crevice treatment to areas no greater than 3 x 3 feet per room	1.5 oz. / 50 sq. ft.
Bed Bugs—Crack and Crevice	Apply as a crack-and-crevice treatment to areas no greater than 3 x 3 feet per room	2.0 oz. / 50 sq. ft.
Ants—Crack and Crevice	Apply as a crack and crevice treatment to areas no greater than 3 x 3 feet per room	1.0 oz. / 50 sq. ft.
Crawling Insects for Severe infestations—Spot treatments	Apply as a crack and crevice or as a spot treatment to areas no greater than 3 x 3 feet per room	16 to 20 oz. / 1,000 sq. ft. [8 to 10 oz/ 500 sq. ft.]
Fleas, ticks and Mites in Areas where Pets frequent—Spot treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room	1 oz. / 70 sq. ft.
Powder Post Beetles –Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room	16 to 20 oz./ 1,000 sq. ft. [8 to 10 oz/ 500 sq. ft.]
Asian Lady Bird Beetles and Box Elder Bugs in Crawl Spaces and Attics—Crack and Crevice	Apply as a spot treatment to areas no greater than 3 x 3 feet per room	1 oz/ 50 sq. ft.

*(continued)*



Application Type	Application rates for Residential Areas such as homes, condominiums, apartments, hotels, motels and pet areas	Application Rates for Non-Residential Areas such as offices, warehouses, hospitals, and other commercial buildings and institutions.
Asian Lady Bird Beetles and Box Elder Bugs in Crawl Spaces and Attics—Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room	16 to 24 oz / 1,000 sq. ft. [4 to 6 oz/ 250 sq. ft.]
Ornamental Plants	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust

**Crawling Insects:**

Apply Evergreen Pyrethrum Dust into voids, cracks and crevices or other places where insects are traveling or harboring. Apply a continuous visible film underneath and behind stairs, behind and beneath sinks, stoves, refrigerators and other appliances, around garbage cans, cabinets. Treat on, around and behind moldings, shelving, baseboards, doorsills, doorframes, floors, around drains, pipes and pipe openings, plumbing, bookcases, storage, utility installations and other areas where insects hide or travel. For severe infestations apply behind cabinets and walls, in crawl spaces and attics at the specified rate.

**Fleas, Ticks, and Mites:**

DO NOT TREAT DOGS OR CATS WITH THIS PRODUCT.

For Fleas, apply Evergreen Pyrethrum Dust only as a direct treatment. Treat on and around animal bedding, sleeping quarters, kennel, furniture, floor and floor coverings and all areas where these pests inhabit. Treat cracks and crevices behind and around window and door frames, wall cracks, moldings, shelving, baseboards, pipe openings and all other cracks. Treat the entire inside surface of doghouses. For Ticks, apply Evergreen Pyrethrum Dust in places where the insects are traveling or harboring.

**Bed bugs:**

Thoroughly inspect area of suspected bed bug infestation, and note potential harborage sites or actual infestations. Remove and/or pick up clothing and other obstacles that may prevent thorough inspection or treatment. Remove and sanitize infested bedding (pillows and linens), clothing, or other items.

Take bed apart. Apply Evergreen Pyrethrum Dust into joints and channels if hollow, such as square or round tubing, and into cracks & crevices on the interior of bed frame, headboards, and footboards. If hollow, ensure that the interior framework is well dusted. Wall-mounted headboards and their hardware should be removed and treated. Mattresses should be dusted, especially seams, edges, and tufts. Mattress treatments must be limited to seams, folds and edges only.

Bed bugs may also harbor in areas of the room away from the bed. Apply Evergreen Pyrethrum Dust as a crack & crevice application where evidence of bed bugs occurs. All furniture and furniture accessories in the room must be treated, including inside empty drawers and shelving in clothes closets, night stands and other furniture in the room, carpet edges, high and low wall moldings (such as floor molding or window casings or ledges/sills), door jambs and hinges, wall voids, and behind wall hangings and mirrors. Pull carpeting from tack strip and treat under the baseboard, sill plate, and under the carpet tack strips. Remove all clothes and other articles from dressers or

closets before application. Treat all cracks & crevices associated with light switches, wall plates, faceplates, outlet covers, mounting boxes and housings. Apply Evergreen Pyrethrum Dust as a continuous film to other potential harborage sites, such as table and wall-mounted lamps, wall crevices, drapery rods, curtains, blinds/mounting hardware, and other crack & crevices in the room.

**Ants:**

To aid in the kill of listed ants, apply Evergreen Pyrethrum Dust at the specified rate. Treat door frames, around window frames, behind, on and around moldings, cabinets, pipe openings, utility openings, shelves, ant trails, hills, wall voids, cracks and crevices and other points of entry into premises. Combine interior treatments with exterior perimeter application of Evergreen Pyrethrum Dust.

**Powder Post Beetles\*:**

To aid in protecting against insect entry. Apply to distribute Evergreen Pyrethrum Dust through insect entry points, crawl holes into attics and crawl spaces of buildings. Close all vents and blowers so that all target surfaces can be treated. Contact all wood surfaces. In case of localized infestations, probe to locate galleries and drill holes to inject Evergreen Pyrethrum Dust into the galleries and void areas. If a heavy infestation is suspected, a conventional, industry accepted techniques and treatments should be employed.

\*The purpose of such applications of Evergreen Pyrethrum Dust is to kill workers or winged reproductive forms that may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. This product is not for use against termites. For active termite infestations, use conventional, industry accepted techniques and treatments.

**Asian Lady Beetles and Box Elder Bugs:**

To kill or repel these insects, as well as Clover Mites, Cluster Flies and Elmleaf Beetles, inject Evergreen Pyrethrum Dust into cracks and crevices and/or voids where these insects may be entering the premise, harboring or hibernating. Apply in crawl spaces and attics. On building exterior, treat underneath wood fascia, behind siding, cracks around windows, doors, siding, wall voids, utility pipes and entry points, behind chimneys, around foundations and foundation cracks, patios and deck joints. When treating for Asian Lady Beetles apply Evergreen Pyrethrum Dust in, on and around areas with abrupt color contrasts on buildings (shutters, drainpipes, etc), southwest-facing or sun-exposed sides of buildings, and vertical corners of buildings.

**Ornamental Trees, Shrubs, Flowers and Foliage Plants:** Including: African Violet, Ageratum, Andromeda, Arborvitae, Ash, Aster, Azalea, Beech, Begonia, Birch, Boxwood, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Chrysanthemum, Cineraria, Coleum, Cotoneaster, Crabapple, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Euonymus, Fern, Ficus, Fir, Firethorn, Foliage Plants, Forsythia, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Hemlock, Hickory, Holly, Honey Locust, Horse Chestnut, Hyacinth, Hydrangea, Iris, Juniper, Larch, Laurel, Lilac, Lily, Linden, Maidenhair Fern, Marigold, Mimosa, Myrtle, Narcissus, Oak, Palm, Pansy, Peony, Pelargonium, Petunia, Philodendron, Phlox, Pine, Privet, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Sweet Pea, Tulip Tree, Tulips, Viburnum, Wandering Jew, Willow, Yew, Zinnia.

Evergreen Pyrethrum Dust must come into contact with insects in order to kill them. Coverage of the target insect is vital to achieving good performance. Apply Evergreen Pyrethrum Dust to cover upper and lower surfaces of leaves stems and flowers with a thin, even film of dust. Apply when insects or insect damage first appears.

When treating trees, shrubs and similar plants, treat the trunk, the base of the trunk and the critical root zone. Treat the tree crotch, knotholes, wounds and other areas where insects may harbor. Allow dust to settle in treated areas before reentering. Do not water or irrigate immediately following treatment. Do not eat or consume treated plants for 1 year after treatment.

**Phytotoxicity:** Plant safety is an important consideration when using insecticides. It is not possible to evaluate the phytotoxicity of Evergreen Pyrethrum Dust on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of Evergreen Pyrethrum Dust treat a limited number of plants and observe for phytotoxicity over a 10 day period.

**Greenhouses:** This product can be used in greenhouses to kill listed insects. Apply to cover upper and lower surfaces of leaves stems and flowers with a thin even film of dust. Apply Evergreen Pyrethrum Dust to ornamental plants or only to fruit, vegetable and herb crops in the plug stage prior to transplant. Do not apply to fruit, vegetable, or herb crops that are ready for consumption.

**OUTDOOR AREAS:** For use in and around areas:

Ant Mounds	Arboretums	Back Yards
Balconies	Beneath Decks	Beneath Gazebos
Calf Hutches	Compost Sites	Courtyards
Decks	Dog Runs	Drive-In Restaurants
Drive-In Theaters	Dumpsters	Farmers Markets
Flea Markets	Front Yards	Gardens
Gazebos	Home Gardens	Kennels
Lanais	Landscapes	Lath Houses
Loading Docks	Lumber Storage Areas	Nurseries
Outdoor Eating Facilities	Patios	Porches
Sanitary Sewers (Manholes)	Side Yards	Swine Yards
Wood Piles		

OUTDOOR APPLICATIONS	
Application Type	Application Rate
Asian Lady Beetles and Box Elder Bugs –Perimeter Building Exterior	16 to 24 oz./ 1,000 sq. ft.
Ornamental Plants	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust
Insects Harboring in Lawns, Landscaping and Backyards	Apply 8 to 24 oz. / 1,000 sq. ft.
Building perimeter treatments	Apply 16 to 24 oz/ 1,000 sq. ft.

**Building Perimeter Treatments:**

To aid in preventing entry and infestations of buildings by listed insects, apply Evergreen Pyrethrum Dust to building exterior as a perimeter treatment in a band extending 12 - 18" along and adjacent to foundation, and to a height of 2 to 3 feet up the side of the building where these pests are active and may find entrance.

**To Kill Insects Harboring in Lawns, Landscaping, and Backyards:**

To aid in the kill of listed insects, apply Evergreen Pyrethrum Dust at the specified rate. For best results, mow lawn one to two (1 – 2) days before applying, and remove clippings. Be sure to cover the entire lawn surface where insects are present, or normally frequent. Apply around the base of ornamental plants, shrubs and trees. Do not water or irrigate immediately following application. Keep pets and children out of the treated area until after the dust has settled.

**LIVESTOCK, DAIRY, SWINE, POULTRY AND OTHER ANIMAL BUILDINGS AND PREMISES:**

<b>LIVESTOCK PREMISES</b>	
<b>Application Type</b>	<b>Application Rate</b>
Crack and Crevice	1 oz. / 50 sq. ft.
Spot treatments and surface treatments	16 to 24 oz / 1, 000 sq. ft. [4 to 6 oz / 250 sq ft.]
Building perimeter treatments	16 to 24 oz/ 1,000 sq. ft. [4 to 6 oz / 250 sq ft.]
Spot Treatments for ticks	16 to 24 oz/ 1,000 sq. ft. [4 to 6 oz / 250 sq ft.]

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken mites, Deer Tick, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice, and Stable Flies

To aid in the kill of listed insects in listed indoor and outdoor animal areas, apply Evergreen Pyrethrum Dust to cracks, crevices, voids, structural joints, floors, walls, wall joints, support beams, trusses, rafters, ceilings, around scales, chutes, gates, and other areas where insects harbor, alight, hide or travel. For Ticks, apply Evergreen Pyrethrum Dust, in areas where they like to harbor such as cracks and crevices.

Apply Evergreen Pyrethrum Dust to exterior of livestock buildings in an 18" band along and adjacent to building foundations, and to a height of 2 to 3 feet up the side of the building where insects are active and may find entrance.

**LIVESTOCK ANIMAL USE:**

<b>LIVESTOCK ANIMALS</b>	
<b>Application Type</b>	<b>Application Rate</b>
Beef, Dairy Cattle, horses	2 oz. dust per animal
Swine	1 oz. per head as a uniform coat to head, shoulders and back.
Poultry	1 lb. / 100 birds. [10 oz / 60 birds]

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken mites, Deer Tick, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice, and Stable Flies

**Beef, Dairy, Cattle, Horses and other Livestock:**

Can be used in dust bags, shaker can and mechanical dust applicator to aid in the kill or repellency of listed livestock insects.

Direct Application: Apply 2 oz. of dust per animal by shaker can, or suitable mechanical dust applicator over the head, neck, shoulders, back and legs.

Application through Dust Bags: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be

suspended in the exit through which the cows leave the milking barn. Hang the bags 4 to 6 inches below the back line of the cattle. For reduction of *Face Flies*, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

**Swine:**

USE RESTRICTIONS FOR APPLICATION TO SWINE:

- Do not apply more often than once every 10 days
- Do not ship animals for slaughter within 5 days of treatment.

**Direct Application:** Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. In severe infestation, both animals and the bedding may be treated.

**Poultry:**

To kill listed livestock insects, apply at a rate of 1 lb. per 100 birds [10 oz per 60 birds].

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep this product in its tightly closed original container, when not in use. Do not store near heat or open flame. Store in a cool, dry, well-ventilated (preferably locked) area that is inaccessible to children and animals, and away from food and pet food.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty container by shaking and tapping sides and bottom to loosen clinging particles promptly after emptying. Then offer for recycling if available or reconditioning, if appropriate or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.



**ACTIVE INGREDIENTS:**

*Pyrethrins	1.00%
<b>OTHER INGREDIENTS</b>	99.00%
	100.00%

Evergreen® - Registered trademark of McLaughlin Gormley King Company

\* Pyrethrins are a botanical insecticide derived from chrysanthemum flowers.

0.01 lbs. active ingredient per lbs. of Evergreen Pyrethrum Dust

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

**SEE BOOKLET FOR FIRST AID STATEMENT  
AND DIRECTIONS FOR USE**

EPA Reg. No. 1021-1871 EPA Est. Nos. 1021-MN-2A,  
F7461-026A 39578-TX-1<sup>B</sup>

Superscript is the first character of the lot number.

Rev 0214-0414

**NET WEIGHT: 10 Ounces**

Manufactured by:

8810 Tenth Avenue North  
Minneapolis, MN 55427

