



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

34704-1031

Date of Issuance:

8/16/2019

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Fitness Fungicide

Name and Address of Registrant (include ZIP Code):

Chris Mason, Senior Manger Registration  
 Loveland Products, Inc.  
 3005 Rocky Moutain Avenue  
 Loveland, CO 80538


**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above Propiconazole pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on her/his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted by email on 8/15/2019. EPA grants this request under the authority of section 3(c)(5) of FIFRA, as amended. With this accepted labeling, all requirements set forth in the Reregistration Eligibility Decision for Propiconazole have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of FIFRA, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after 12 months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

Signature of Approving Official:

  
 Shaja B. Joyner, Product Manager 20  
 Fungicide-Herbicide Branch  
 Registration Division 7505P

Date:

8/16/2019

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 04/28/2016
- Alternate CSF 1 dated 10/03/2017
- Alternate CSF 2 dated 10/03/2017
- Alternate CSF 3 dated 10/03/2017
- Alternate CSF 4 dated 10/03/2017
- Alternate CSF 5 dated 10/03/2017
- Alternate CSF 6 dated 10/03/2017
- Alternate CSF 7 dated 10/03/2017
- Alternate CSF 8 dated 10/03/2017
- Alternate CSF 9 dated 10/03/2017
- Alternate CSF 10 dated 10/03/2017

If you have any questions, please contact Shaja B. Joyner by phone at 703-308-3194, or via email at [joyner.shaja@epa.gov](mailto:joyner.shaja@epa.gov).

Enclosure



# FITNESS<sup>®</sup>

## FUNGICIDE

For control of certain diseases in almonds, berries, carrot, celery, cereals, citrus (non-bearing), corn, cranberry, filbert, mint, onions and bulb vegetables, peanut, pecan, pineapple, rice (including wild rice), sorghum, soybeans, stone fruits, sugarcane, tree nuts, grasses grown for seed, turfgrass and ornamentals.

(Not for Turf and Ornamental use in California)

**ACTIVE INGREDIENT:**

Propiconazole:.....41.8%

**OTHER INGREDIENTS:**.....58.2%

**TOTAL                    100.0%**

Contains petroleum distillates.

Contains 3.6 pounds active ingredient per gallon.


Emulsifiable Concentrate

### KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

For Additional Precautionary Statements, Directions for Use, Storage and Disposal and  
Other Use Information, See Inside This Label Booklet.

<b>FIRST</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p> <p><b>FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.</b></p> <p><b>Note to Physician:</b> Contains petroleum distillate — vomiting may cause aspiration pneumonia.</p>	



EPA REG. NO. 34704-  
1031 EPA EST. NO.  
34704-MS-002  
**NET CONTENTS 1.0 GAL (3.78 L)**

EXP 0619

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**WARNING**

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using the toilet.

**Personal Protective Equipment (PPE):**

**Applicators and all other handlers must wear:**

- Protective eyewear.
- Long-sleeved shirt and long pants,
- Chemical-resistant gloves, such as barrier laminate or viton,
- Shoes plus socks, and
- Protective eyewear

**All handlers using propiconazole as a seed piece treatment must wear:**

- Protective eyewear (goggles, face shield, or safety glasses)
- Chemical-resistant gloves made of barrier laminate or viton
- Chemical-resistant apron

**User Safety Requirements**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)1], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and shrimp. Do not apply directly to water, or to areas where surface water is present, or to inter-tidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

**PHYSICAL OR CHEMICAL HAZARDS**

Combustible. Do not use, pour, spill or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval.

The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Protective eyewear
- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses: Do not enter into treated areas until sprays have dried.**

**CROPS**

**Product Information**

Fitness Fungicide is a broad-spectrum fungicide for the control of certain diseases in almonds, berries, carrots, celery, cereals (wheat, barley, rye, oats, and rice), citrus (non-bearing), corn, cranberries, filberts, mint, onions and other bulb vegetables, sorghum, soybeans, stone fruit (apricots, nectarines, peaches, plums, and prunes), peanuts, pecans, pineapple, sugarcane, tree nuts, grasses grown for seed, and turfgrass and ornamentals.

**Restriction:** Do not use in greenhouses or as a tree injection.

Failure to follow directions and precautions on this label may result in crop injury, poor disease control, or illegal residues.

**Integrated Pest Management**

Fitness Fungicide must be integrated into an overall disease and pest management (IPM) strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development

**FITNESS® FUNGICIDE EPA REG. NO**  
**34704-1031**

must be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. Fitness Fungicide may be used in state agricultural extension advisory (disease forecasting) programs using the recommend application timing based upon environmental factors favorable for disease development.

**Resistance Management**

Fitness Fungicide belongs to the sterol demethylation inhibitor (DM1) class of fungicides and is classified as Group 3 Fungicide. Any fungal population may contain individuals naturally resistant to Fitness Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of (name of product) or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Loveland Products, Inc. at 888-574-2878 or at [www.lovelandproducts.com](http://www.lovelandproducts.com). You can also contact your pesticide distributor or university extension specialist to report resistance.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**Rotational Crops:**

<b>Rotational Crops</b>	<b>Planting Time from Last Fitness Application</b>
Beans, Dry and Succulent Brassica, Leafy greens (Subgroup 5B) Bulb Vegetables Carrots Celery and Leaf Petioles, Subgroup 4B Cereals (wheat, barley, rye, triticale, oats) Cilantro (Coriander) Leaves Corn (field, seed, pop, and sweet) Dill Garden Beets Grasses grown for seed (Cool season grasses only) Mint Parsley, Fresh and Dried leaves Peanuts Quinoa Radish Rapeseed (Canola), Subgroup 20A Rice (including Wild Rice) Sorghum Soybeans Strawberry and other Low-Growing Berry Subgroup 13-07G Sugar Beets Sugarcane Watercress	0 days
Buckwheat Millet	12 Months
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/year)	75 days
All Other Crops Intended for Food, Grazing, or Feed	105 days

Any crop listed and approved for use on this label has a 0 day plant back interval.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**MIXING INSTRUCTIONS**

**Application Rate Reference Table**

<b>Application Rate of Fitness Fungicide (Fl Oz/A)</b>	<b>Active Ingredient Equivalent (Lb AI/A)</b>	<b>A/Gal of Fitness Fungicide</b>
2.0	0.056	64
4.0	0.1125	32
6.0	0.169	21.3
8.0	0.225	16
10.0	0.28	12.8
12.0	0.34	10.7
16.0	0.45	8
20.0	0.56	6.4
24.0	0.67	5.3
30.0	0.84	4.3
32.0	0.90	4

Prepare no more spray mixture than is required for the immediate operation. Thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application. Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

**Fitness Fungicide - Alone:** Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add Fitness Fungicide to the tank. Continue agitation while adding the remainder of the water. Begin application of the spray solution after Fitness Fungicide has completely dispersed into the mix water. Maintain agitation until all of the mixture has been sprayed.

**Fitness Fungicide - Tank Mixes:** Fitness Fungicide is usually compatible with all tank mix partners listed on this label. To determine the physical compatibility of Fitness Fungicide with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1.0 quart of water. Add wettable powders and water dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank. Add 1/2 to 2/3 of the required amount of water to the spray or mixing tank. With the agitator running, add the tank mix partner into the tank. Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Fitness Fungicide to the spray tank. Allow Fitness Fungicide to completely disperse. Spray the mixture with the agitator running.

**Precaution:** Do not apply this product in a tank mix with a dodine fungicide or crop injury may occur.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.



**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

If using Fitness Fungicide in a tank mix, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations required by the label of the tank mix product. No label dosage rate can be exceeded and the most restrictive label precautions and limitations must be followed. Do not tank mix this product with any product that prohibits such mixing. Tank mixes or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

### **Application Instructions**

Fitness Fungicide is most effective when applied and allowed to dry before a rainfall. Avoid applying Fitness Fungicide under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner which results in exposure to humans or animals.

**Ground Application:** For tree crops, apply Fitness Fungicide in a minimum of 50.0 gallons of water per acre unless otherwise specified. For all other crops, apply Fitness Fungicide in a minimum of 10.0 gallons of water per acre unless otherwise specified.

**Aerial Application:** For tree crops, apply Fitness Fungicide in a minimum of 10.0 gallons of water per acre unless otherwise specified. For all other crops, apply Fitness Fungicide in a minimum of 2.0 gallons of water per acre unless otherwise specified.

**Chemigation Application:** This product may be applied through properly equipped chemigation systems for disease control in the labeled crops. Refer to crop specific use directions for application rates, timing and frequency of application. Do not apply Fitness Fungicide by chemigation to other labeled crops except as specified in Loveland Products supplemental labeling or product bulletins. When applying this product by chemigation, do not exceed labeled rates or apply more frequently than specified for conventional application methods. Fitness Fungicide, alone or in combination with other pesticides that are registered for application through irrigation systems, may be applied through irrigation systems. For chemigation application to labeled crops, apply in 0.1 to 0.25 inches of water unless otherwise specified. Chemigation with excessive water may lead to a decrease in efficacy.

### **Spray Drift Management**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

- **Wind Speed:** Do not apply at wind speeds greater than 15 mph.
- **Droplet Size:** Apply as a medium or coarser spray (ASAE Standard 572).
- **Temperature Inversions:** If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.
- **Other State and Local Requirements:** Applicators must follow all state and local pesticide drift requirements regarding application of propiconazole. Where states have more stringent regulations, they must be observed.
- **Equipment:** All application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.
- **Aerial application:** The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. When applications are made with a crosswind, the swath must be

**FITNESS® FUNGICIDE EPA REG. NO**  
**34704-1031**

displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft

- **Ground boom application:** Do not apply with a nozzle height greater than 4 feet above the crop canopy.

### **Spray Equipment**

Thorough coverage is necessary to provide good disease control. Avoid spray overlap as crop injury may occur.

Air assisted or air blast sprayers move spray droplets into the canopy using a forced air stream. Set up the fan to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use.

Use a pump with sufficient capacity to maintain 35 to 40 psi at nozzles and provide sufficient agitation in tank to keep mixture in suspension (this requires recirculation of 10% of tank volume per minute). Use a jet agitator or liquid sparger tube for agitation. Do not use air sparging.

Although Fitness Fungicide is an emulsifiable concentrate, it is suggested that screens be used to protect the pump and to prevent nozzles from clogging. Screens placed on suction side of pump must be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom and, where required, at the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

### **CHEMIGATION INSTRUCTIONS**

- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.
- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

**Note:** Do not inject Fitness Fungicide at full strength or deterioration of valves and seals may

## FITNESS® FUNGICIDE EPA REG. NO

34704-1031

occur. Use a dilution ratio of at least 10 parts water to 1 part Fitness Fungicide. Fitness Fungicide is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but replace once a year. Do not use Viton, Buna-N, Neoprene, or PVC seals.

### Specific Equipment Requirements

- The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline, must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being with-drawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

### Center Pivot Irrigation Equipment

- Use only with drive systems that provide uniform water distribution.
- Do not use end guns when applying Fitness Fungicide through center pivot systems because of non-uniform application.
- Determine size of area to be treated.
- Determine the time required to apply 1/8 to 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying Fitness Fungicide through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80 to 95% of the manufacturer's rated capacity.
- Using only water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of Fitness Fungicide required to treat the area covered by the irrigation system.
- Add the required amount of Fitness Fungicide and sufficient water to meet the injection time requirements of the solution tank.
- Make sure the system is fully charged with water before starting injection of Fitness Fungicide. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Stop injection equipment after treatment is completed. Continue to operate the system until the solution of Fitness Fungicide has cleared the sprinkler head.

### Solid-Set, Hand Move, and Moving Wheel Irrigation Equipment

- Determine the acreage covered by the sprinkler.

## FITNESS® FUNGICIDE EPA REG. NO

34704-1031

- Fill the injector solution tank with water and adjust flow rate to use the contents over a 20- to 30-minute interval. When applying Fitness Fungicide through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of Fitness Fungicide required to treat the area covered by the irrigation system.
- Add the required amount of Fitness Fungicide into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate the system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during the calibration.
- Inject Fitness Fungicide at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until the solution of Fitness Fungicide has cleared the last sprinkler head.

### CROP SPECIFIC INSTRUCTIONS

#### ALMONDS

##### Restrictions

- **DO NOT** apply more than 32.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (8 fl oz/A) or 8 applications per year when applying at the lowest rate (4 fl oz/A).
- **DO NOT** apply more than 0.90 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** graze livestock in treated areas or cut treated cover crop for feed.
- Preharvest Interval (PHI) is 60 days.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 15.0 gallons of water per acre). Fitness Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume must be used to provide thorough coverage.

Target Disease	Application Rate (FI Oz/A)	Application Instructions
Brown rot blossom blight ( <i>Monilinia laxa</i> , <i>M. fructicola</i> )	4.0 to 8.0	Apply Fitness Fungicide at 5 to 10% bloom and 50 to 100% bloom. Under severe disease conditions, use the highest rate. Minimum retreatment interval is 7 days.
Anthracnose ( <i>Collectotrichum acutatum</i> )	8.0	Apply Fitness Fungicide beginning at bud break on a 7- to 14- day interval.

#### BUSHBERRIES (Subgroup 13-07B) AND CANEBERRIES (Subgroup 13-07A)

Bushberries (subgroup 13-07B) including aronia berry, black currant, buffalo currant, Chilean guava, edible honeysuckle, elderberry, European barberry, gooseberry, highbush blueberry, highbush cranberry, huckleberry, jostaberry, Juneberry, lingonberry, lowbush blueberry, native currant, red currant, salal, sea buckthorn, cultivars, varieties, and/or hybrids of these.

Caneberries (subgroup 13-07A) including bingleberry, blackberry, black raspberry, boysenberry, dewberry, Loganberry, lowberry, Marionberry, olallieberry, red raspberry, wild raspberry, youngberry, cultivars, varieties and/ or hybrids of these.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**Restrictions**

- **DO NOT** apply more than 30.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 6.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 5 applications per year.
- Preharvest Interval (PHI) is 30 days.
- **DO NOT** apply more than 0.84 pound active ingredient propiconazole-containing product per acre per year.

Apply Fitness either ground (a minimum of 15.0 gallons per acre) or aerial application (a minimum of 5.0 gallons per acre). Fitness is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume must be used to provide thorough coverage.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Mummyberry disease ( <i>Monilinia vaccinocorymbosi</i> )	6.0	Make first application of Fitness Fungicide beginning at green tip and repeat in 7 to 10 days. If conditions are favorable for disease development, additional applications may need to be made at pink bud and repeating every 7 to 10 days through petal fall.
Leaf spot and Stem canker ( <i>Septoria albopuncatata</i> ) Rust ( <i>Pucciniastrumvaccinii</i> )	6.0	Apply when conditions favor disease development. Repeat applications on a 4-week spray interval.
Leaf and Cane spot ( <i>Septoria rubi</i> )	6.0	Apply as a delayed dormant spray after training in the spring. Repeat this application in the late spring, again at bud break, and again once flowering has begun.
Powdery mildew ( <i>Microsphaera vaccinii</i> )	6.0	Apply Fitness Fungicide at 5 to 10% bloom. Repeat this application at full bloom and on a 14-day interval while conditions are favorable for disease.
Leaf spot ( <i>Septoria</i> spp.)	6.0	Make first application any time prior to bloom and again after petal fall. If needed, repeat application just after harvest.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**CARROTS**

**Restrictions**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (4 fl oz/A) or 8 applications per year when applying at the lowest rate (2 fl oz/A).
- Preharvest Interval (PHI) is 14 days.

For best results, sufficient water volume must be used to provide thorough coverage. Apply Fitness Fungicide by ground (a minimum of 15.0 gallons per acre) or aerial equipment (a minimum of 5.0 gallons per acre). Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Leaf blights ( <i>Cercospora carotae</i> ) (suppression of <i>Alternaria dauci</i> )	4.0	Apply 4.0 oz of Fitness Fungicide when conditions favor disease development. Continue applications on a 7- to 10-day interval, using the shorter interval when disease conditions are severe. If desired, a spreader-sticker may be used.
Powdery mildew ( <i>Erysiphe polygoni</i> )	Fitness Fungicide plus chlorothalonil 2.0	Apply 2.0 oz of Fitness Fungicide with 0.75 lb AI of chlorothalonil/A (1.0 pt Initiate® 720 Flowable Fungicide). Begin applications when conditions favor disease development. Continue applications on a 7- to 10-day interval.

**CELERY AND LEAF PETIOLES (Subgroup 4B)**

Leaf petioles (subgroup 4B) including cardoon, celery, Chinese celery, celtuce, Florence fennel, rhubarb, Swiss chard.

**Restrictions**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 4 applications per year.
- Preharvest Interval (PHI) is 14 days.

Fitness Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume must be used to provide thorough coverage. Fitness Fungicide may be applied by either ground (a minimum of 10.0 gallons per acre) or aerial application (a minimum of 5.0



**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

gallons per acre).

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Early blight ( <i>Cercospora apii</i> ) Late blight ( <i>Septoria apicola</i> )	4.0	Apply Fitness Fungicide on a 7-day schedule, either by ground or aerial application. If desired, Fitness Fungicide may be mixed with appropriate spreader-sticker.

**CEREALS**

Barley, buckwheat, corn (see specific directions elsewhere on this label), Pearl millet, proso millet, oats, rye, rice (see specific directions elsewhere on this label), teosinte, triticale, wheat, wild rice (see specific directions elsewhere on this label) and other commodities included in the cereal grains crop grouping.

**Restrictions**

- **DO NOT** apply more than 8.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces of Fitness Fungicide (0.11 pound active ingredient) per acre per year if forage or hay will be harvested.
- **DO NOT** apply more than 4.0 fluid ounces of Fitness Fungicide (0.11 pound active ingredient) per application.
- **DO NOT** apply more than 0.22 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 2 applications per year when applying at the highest rate (4 fl oz/A) or 4 applications per year when applying at the lowest rate (2 fl oz/A)
- **DO NOT** apply after Feekes growth stage 10.5.4.
- Preharvest Interval (PHI) is 7 days for forage or hay.
- Cereals other than wheat: Preharvest Interval (PHI) is 45 days for grain and straw.

Apply Fitness Fungicide by ground, aerial or chemigation equipment. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
<p><b>Early season suppression of:</b>  Glume blotch (<i>Stagonospora nordorum</i>) Leaf blight (<i>Septoria tritici</i>)  Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.)  Tan spot (<i>Pyrenophora tritici-repentis</i>)</p>	<p>2.0 to 4.0</p>	<p>Apply Fitness Fungicide in the spring. Make a second application up to Feekes growth stage 10.5 for season long control. The minimum retreatment interval is 14 days.</p>
<p><b>Control of leaf diseases:</b>  Barley scald (<i>Rhynchosporium secalis</i>) Barley stripe (<i>Pyrenophora graminea</i>) Glume blotch (<i>Stagonospora nordorum</i>)  Helminthosporium leaf blight (<i>Drechslera tritici-repentis</i>)  Leaf blight (<i>Septoria tritici</i>)  Net blotch (<i>Pyrenophora teres</i>)  Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.)  Rust (<i>Puccinia</i> spp.)  Spot blotch (<i>Bipolaris sorokiniana</i>)  Tan spot (<i>Pyrenophora tritici-repentis</i>)</p>	<p>4.0</p>	<p>Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Fitness Fungicide is applied when the flag leaf is 50% to fully emerged. The minimum retreatment interval is 14 days. Using an oil base adjuvant may improve the spray coverage and canopy penetration. In wheat: Fitness Fungicide can be applied at full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues.</p>
<p>Foot rot (<i>Pseudocercospora</i> spp.)</p>	<p>4.0</p>	<p>Apply Fitness Fungicide plus half rates of fungicides such as thiophanate-methyl. Apply at tillering, but before elongation has occurred.</p>
<p>Fusarium head blight Suppression</p>	<p>4.0</p>	<p>Apply Fitness Fungicide at approximately 50% flowering. Addition of a penetrating type of adjuvant may increase Fusarium head blight suppression.</p>



**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**CITRUS (non-bearing)**

(Calamondin, citron, citrus hybrids, grapefruit, kumquat, lemon, lime, Mandarin, orange (sour and sweet), pummelo, satsuma mandarin, tangerine, including all cultivars and/or hybrids of these.)

**Restrictions**

- **DO NOT** apply more than 24.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.67 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 3 applications per year when applying at the highest rate (8 fl oz/A) or 4 applications per year when applying at the lowest rate (6 fl oz/A).
- **DO NOT** apply to citrus that will bear harvestable fruit within 12 months..

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 15.0 gallons per acre). Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

<b>Target Disease</b>	<b>Application Rate (Fl Oz/A)</b>	<b>Application Instructions</b>
Greasy spot ( <i>Mycosphaerella citri</i> )	6.0 to 8.0	Begin applications in June. Apply on 30-day interval through August.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**CORN**

**Restrictions:**

**For Field corn, Field corn grown for seed, and Popcorn:**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces of Fitness Fungicide per acre per year on field corn harvested for forage.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (4 fl oz/A) or 8 applications per year when applying at the lowest rate (2 fl oz/A).
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.
- Preharvest Interval (PHI) is 30 days for forage, grain and stover.

**For Sweet corn:**

- **DO NOT** apply within 14 days of harvest for ears (PHI).
- **DO NOT** apply within 14 days of harvest for forage (PHI).

Apply Fitness Fungicide by ground, aerial or chemigation equipment. For best results, sufficient water volume must be used to provide thorough coverage. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

Target Disease	Application Rate (FI Oz/A)	Application Instructions
Helminthosporium leaf blights ( <i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i> ) Northern corn leaf blight ( <i>Setosphaeria turcica</i> ) Northern corn leaf spot ( <i>Cochliobolus carbonum</i> ) Southern corn leaf blight ( <i>Cochliobolus heterostrophus</i> )	2.0 to 4.0	Apply Fitness Fungicide when disease first appears and reapply on a 7- to 14-day schedule. Use the low rate when disease pressure is low and the high rate under heavy disease pressure or if conditions are favorable for disease development.
Rusts ( <i>Puccinia</i> spp.) Gray leaf spot ( <i>Cercospora zeae-maydis</i> ) Eye spot ( <i>Aureobasidium zeae</i> )	4.0	Apply Fitness Fungicide when disease first appears. If conditions favorable for disease persist, continue to apply on a 7- to 14-day schedule. For best disease control, early applications (at initial disease onset) perform better.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**CRANBERRY  
Restrictions**

- **For use only in Oregon, Washington and Wisconsin.**
- **DO NOT** apply more than 24.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 6.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.67 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (6 fl oz/A) or 6 applications per year when applying at the lowest rate (4 fl oz/A).
- Preharvest Interval (PHI) is 45 days.
- **DO NOT** allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.
- **DO NOT** use cranberry fields used for aquaculture of fish and crustaceans.
- **DO NOT** apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Applicators must use care in making applications near non-target aquatic habitats.
- **DO NOT** apply to flooded crop.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 20.0 gallons per acre). Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

<b>Target Disease</b>	<b>Application Rate (FI)</b>	<b>Application Instructions</b>
Cottonball ( <i>Monilinia oxycocci</i> )	4.0 to 6.0	Make the first application at leaf bud break and repeat in 14 days. Make the third application at early bloom and repeat in 14 days. Under severe pressure, use the higher rate for control.

**FITNESS® FUNGICIDE EPA REG. NO**

**34704-1031**

**FILBERTS (Hazelnuts)**

**Restrictions**

- **DO NOT** apply more than 32.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide
- **DO NOT** apply more than 0.90 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (8 fl oz/A) or 6 applications per year when applying at the lowest rate (5 fl oz/A).
- **DO NOT** graze livestock in treated areas or cut treated cover crop for feed.
- Preharvest Interval (PHI) is 60 days.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 15.0 gallons per acre). Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

<b>Target Disease</b>	<b>Application Rate (FI Oz/A)</b>	<b>Application Instructions</b>
Eastern filbert blight ( <i>Anisogramma anomala</i> )	5.0 to 8.0	Begin applications when green leaf tissue becomes visible and continue on 14- to 21-day interval. Under severe disease conditions, use the higher rate and shorter interval. <b>Note:</b> On certain varieties, Fitness Fungicide applications may cause smaller and/or greener leaves. Yields of filberts displaying these characteristics have not been reduced due to Fitness Fungicide treatments.

**MINT (Peppermint, Spearmint) Restrictions**

- **DO NOT** apply more than 12.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.34 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 3 application per year.
- **DO NOT** apply within 7 days of harvest (7-day PHI).

Apply Fitness Fungicide by ground in a minimum of 20.0 gallons per acre, or aerial equipment in a minimum of 2.0 gallons per acre. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, use sufficient water to provide thorough coverage.

<b>Target Disease</b>	<b>Application Rate (FI Oz/A)</b>	<b>Application Instructions</b>
Rust ( <i>Puccinia menthae</i> )	4.0	Begin applications when the plants are 2 to 4 inches high or when conditions become favorable for disease development. Continue on a 14-day interval as conditions warrant.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**ONIONS (Subgroups 3-07A) and other BULB VEGETABLES (Subgroups and 3-07B)**

Dry bulb (garlic, dry bulb onion, dry bulb shallots)

Green (green onions, green shallots, green eschalots, Japanese bunching onions, leeks, spring onions, scallions) and/or cultivars or hybrids of these.

**Restrictions**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 2 applications per year when applying at the highest rate (8 fl oz/A) or 8 applications per year when applying at the lowest rate (2 fl oz/A). When applying 4 fl oz/A, do not exceed 4 applications per year.
- Preharvest Interval (PHI) for dry bulb onions is 14 days.
- Preharvest Interval (PHI) for green onions is 0 days.

For best results, sufficient water volume must be used to provide thorough coverage. Apply Fitness Fungicide by ground (a minimum of 15.0 gallons per acre) or aerial equipment (a minimum of 5.0 gallons per acre). Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Purple blotch ( <i>Alternaria porri</i> ) Suppression of Botrytis leaf blight ( <i>Botrytis squamosa</i> )	Fitness Fungicide alone 4.0 to 8.0	Begin applications when conditions favor disease development and continue on a 7- to 10-day interval. Use the higher rate and shorter interval when disease conditions are severe.
	Fitness Fungicide plus tank mix partner 2.0 to 4.0	Apply 2.0 to 4.0 oz of Fitness Fungicide in combination with another fungicide registered for control of Botrytis leaf blight or Purple blotch. Begin applications when conditions favor disease development and continue on 7-day interval or according to the tank mix partner label. Use the higher rate when disease conditions are severe. To achieve optimum disease control use a wetting agent or spreader-sticker.

**PEANUTS**

**Restrictions**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** exceed 2 applications per year when applying at the highest rate (8 fl oz/A) or 6 applications per year when applying at the lowest rate (2.5 fl oz/A). When applying 4 fl oz/A do not exceed 4 applications per year.

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

- **DO NOT** feed hay from treated fields to livestock if the high rate is used (8.0 fluid ounces per acre).
- Preharvest Interval (PHI) is 14 days with the low use rate (4.0 fluid ounces per acre).
- Preharvest Interval (PHI) is 21 days with the high use rate (8.0 fluid ounces per acre).

Apply Fitness Fungicide by ground, aerial or chemigation equipment. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

<b>Target Disease</b>	<b>Application Rate (Fl Oz/A)</b>	<b>Application Instructions</b>
Early leaf spot ( <i>Cercospora arachidicola</i> )	2.5 to 4.0	Apply beginning 35 to 40 days after planting or at the first appearance of disease and reapply on a 10- to 14-day schedule. Under heavy disease pressure, use the higher rate. Fitness Fungicide also may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based upon environmental factors favorable for disease development.
Late leaf spot ( <i>Cercosporidium personatum</i> ) Rust ( <i>Puccinia arachidis</i> ) Web blotch ( <i>Phoma arachidicola</i> )	4.0	
Southern stem rot ( <i>Scierotium rolfsii</i> )	4.0-8.0	

**FITNESS® FUNGICIDE EPA REG. NO  
34704-1031**

**PECANS**

**Restrictions**

- **DO NOT** apply more than 32.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** apply more than 0.90 pound active ingredient propiconazole-containing products per acre per year.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (8 fl oz/A) or 8 applications per year when applying at the lowest rate (4 fl oz/A).
- **DO NOT** apply after shuck split or within 45 days of harvest, whichever comes first..
- **DO NOT** graze livestock in treated areas or cut treated cover crop for feed.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 20.0 gallons per acre). For best results, use sufficient water to provide thorough coverage. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

The active ingredient in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your county has a bulletin consult <http://www.epa.gov/espp/bulletins.htm>. Bulletins may also be available from local pesticide dealers, extension offices, or state pesticide agencies.

<b>Target Disease</b>	<b>Application Rate (Fl Oz/A)</b>	<b>Application Instructions</b>
Downy spot ( <i>Mycosphaerella caryigena</i> ) Liver spot ( <i>Gnomonia caryae pv pecanae</i> ) Pecan scab ( <i>Cladosporium caryigenum</i> ) Powdery mildew ( <i>Microsphaera penicillata</i> ) Vein spot ( <i>Gnomonia nerviseda</i> ) Zonate leaf spot ( <i>Cristulariella moricola</i> )	4.0 to 8.0	<b>Pecan scab:</b> Apply on a 14-day schedule during bud break and pre-pollination sprays. Apply 6.0 to 8.0 fl oz/A during nut formation and cover sprays. Use higher rates when disease pressure is heavier. <b>Other listed foliar diseases:</b> Apply 4.0 fl oz/A with other products registered for pecans and labeled for these mid to late-season foliar diseases. Observe all directions, precautions, and limitations for the other products.

FITNESS® FUNGICIDE EPA REG. NO  
34704-1031

**PINEAPPLE (Hawaii  
Only) Restrictions**

- **DO NOT** use treated crowns for food or feed.
- **DO NOT** graze while plant is growing.
- **DO NOT** graze tops until fruit is harvested.
- Dispose of used dip solution according to local, state and Federal regulations.

Target Disease	Application Rate (Fl Oz/100 gal water)	Application Instructions
Butt rot ( <i>Ceratocystis paradoxa</i> )	0.75 (1:17,000)	Treatments can be made in either a cold or hot water dip. <b>Cold Water Dip:</b> Immerse crowns to give thorough wetting, remove, and allow to drain. <b>Hot Water Dip:</b> Maintain water temperature at 125 °F (52 °C). Soak crown for 20 to 30 minutes, remove, and allow to drain.



**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**RICE**

**Restrictions**

- **DO NOT** apply to stubble or ratoon crop rice.
- **DO NOT** use in rice fields where commercial farming of crayfish will be practiced.
- **DO NOT** drain water from treated rice fields into ponds used for commercial fish farming.
- **DO NOT** use water drained from treated fields to irrigate other crops.
- **DO NOT** release flood water within 7 days of an application.
- **DO NOT** apply more than 12.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 10.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 1 application per year when applying at the highest rate (10 fl oz/A) or 2 applications per year when applying at the lowest rate (6 fl oz/A).
- **DO NOT** apply more than 0.34 pound active ingredient propiconazole-containing product per acre per year.
- Preharvest Interval (PHI) is 35 days.

Apply Fitness Fungicide by aerial equipment only. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your County has a Bulletin, consult [www.epa.gov/espp/bulletins.htm](http://www.epa.gov/espp/bulletins.htm). Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Aggregate sheath spot <i>(Rhizoctonia oryzae-sativa)</i> Black sheath rot <i>(Gaeumannomyces graminis)</i> Brown leaf spot <i>(Helminthosporium oryzae)</i> False smut <i>(Ustilaginoidea virens)</i> (suppression only) Kernel smut <i>(Tilletia barclayana)</i> Leaf smut <i>(Entyloma oryzae)</i> Narrow brown leaf spot <i>(Cercospora oryzae)</i> Sheath blight <i>(Rhizoctoria solani)</i> Sheath spot <i>(Rhizoctonia oryzae)</i> Stem rot <i>(Sclerotium oryzae)</i> (suppression only)	6.0 to 10.0	Timing of Fitness Fungicide application depends upon disease severity, disease complex and rice variety/growth stage. Consult local extension experts for local economic thresholds established for various rice varieties and diseases. Apply at first internode elongation (up to 2-inch panicle) using one of the following schedules:  <b>A.</b> Apply 6.0 fl oz/A and repeat at swollen boot. Make the second application 14 days later but before the boot splits and head emerges. Sheath blight is best controlled when the first application is applied at disease appearance in the field. For first application, apply when 5% or fewer of the tillers are infected.  <b>B.</b> Apply 10.0 fl oz/A. This rate is specified if greater than 10% of the tillers are infected with Sheath blight. If disease reappears, use another registered fungicide for the second application.  Tank mix option: Apply 6.0 fl oz of Fitness Fungicide in a tank mix with registered fungicides for control of rice diseases.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**WILD RICE (Minnesota only) Restrictions**

- **DO NOT** apply to stubble or ratoon crop rice.
- **DO NOT** use in rice fields where commercial farming of crayfish will be practiced.
- **DO NOT** drain water from treated rice fields into ponds used for commercial fish farming.
- **DO NOT** use water drained from treated fields to irrigate other crops.
- **DO NOT** release flood water within 7 days of an application.
- **DO NOT** apply more than 12.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 10.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 1 application per year when applying at the highest rate (10.0 FI Oz/A) or 2 applications per year when applying at the lowest rate (6.0 FI Oz/A)
- **DO NOT** apply more than 0.34 pound active ingredient propiconazole-containing product per acre per year.
- Preharvest Interval (PHI) for forage is 35 days.

Apply Fitness Fungicide by aerial equipment only. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your County has a Bulletin, consult [www.epa.gov/espp/bulletins.htm](http://www.epa.gov/espp/bulletins.htm). Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

Target Disease	Application Rate (FI)	Application Instructions
Brown spot ( <i>Bipolaris</i> spp.) Helminthosporium leaf blight ( <i>Helminthosporium</i> spp.)	6.0 to 8.0	Apply 6.0 fl oz/A of Fitness Fungicide at both booting and heading, or make a single application of 8.0 fl oz/A at booting. Minimum re-application interval is 10 days.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**SORGHUM**

**Restrictions**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per year of Fitness Fungicide. (0.22 pound active ingredient propiconazole) on sorghum to be harvested for forage.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide
- **DO NOT** exceed 4 applications per year.
- Preharvest Interval (PHI) for forage is 30 days
- Preharvest Interval (PHI) for grain and stover is 21 days.
- **DO NOT** graze livestock or cut for green chop or silage within 30 days of application.
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.

Apply Fitness Fungicide by ground (a minimum of 15.0 gallons per acre) or aerial equipment (a minimum of 10.0 gallons per acre). For best results, use sufficient water to provide thorough coverage. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Ergot ( <i>Claviceps sorghi</i> )	3.0 to 4.0	Make the first application at or just prior to flowering. Repeat on a 5- to 7-day interval. Apply up to 4 times.

**SOYBEANS**

**Restrictions**

- **DO NOT** apply more than 12.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 6.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 2 applications per year when applying at the highest rate (6 fl oz/A) or 3 applications per year when applying at the lowest rate (4 fl oz/A).
- **DO NOT** apply more than 0.34 pound active ingredient propiconazole-containing product per acre per year.
- Apply up to Stage R6

**Precautions**

On certain varieties, Fitness Fungicide applications may cause crinkled, smaller and/or greener leaves. Yields of dry beans displaying these characteristics have not been reduced due to Fitness Fungicide treatments.

Fitness Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume must be used to provide thorough coverage. Fitness Fungicide may be applied by either ground or aerial application.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Aerial web blight ( <i>Rhizoctonia solani</i> )	5.0 to 6.0	Aerial web blight: Apply at the first appearance of disease and repeat the application 14 to 21 days later. Under severe disease conditions use the higher

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

<p>Anthracnose (<i>Colletotrichum truncatum</i>),          Brown spot (<i>Septaria glycines</i>)          Frogeye leaf spot (<i>Cercospora sojina</i>)</p>	<p>6.0</p>	<p>Apply at growth stage R3 (early pod set when pods are 1/8 to 1/4 inch long) and 14 to 21 days later at growth stage R5 (pod fill).</p>
<p>Soybean Rust (<i>Phakopsora pachyrhizi</i>)</p>	<p>4.0 to 6.0</p>	<p>Soybean rust: Apply at first indication that disease is in the area. For best control preventive applications work best. Repeat on 14 to 21-day interval. Use higher rate and shorter interval when disease is present in field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.</p>

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**STONE FRUITS (Crop Group 12)**

Apricot, Chickasaw plum, Damson plum, Japanese plum, nectarine, peach, plum, plum-cot, prune, sweet cherry, tart cherry, and cultivars and/or hybrids of these included in the stone fruits crop grouping

**Restrictions**

- **DO NOT** apply more than 20.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 5 applications per year.
- **DO NOT** apply more than 0.56 pound active ingredient propiconazole-containing product per acre per year.
- Preharvest Interval (PHI) is 0.

**Precaution**

Applications of Fitness Fungicide during bloom to Stanley plums have occasionally caused fruit to be less oval in shape and smaller in size at harvest. To avoid this, do not apply to Stanley plums earlier than 21 days before harvest.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 15.0 gallons per acre). For best control of stone fruit diseases, apply by ground application. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

Target Disease	Application Rate (Fl Oz/A)	Application Instructions
Brown rot blossom blight ( <i>Monilinia</i> spp.) Cherry leaf spot ( <i>Blumeriella jaapii</i> ) Fruit brown rot ( <i>Monilinia</i> spp.) Powdery mildew ( <i>Podosphaera</i> spp.) Rust ( <i>Tranzschelia discolor</i> )	4.0	Apply at early bloom stage. If disease pressure is low, a second application may be made as needed through petal fall. Under conditions of high disease pressure and/or very susceptible varieties, make a second application from 75 to 100% bloom and a third application at petal fall. Apply a maximum of 2 sprays as needed during the preharvest period up to the day of harvest. If high inoculum and severe disease conditions persist, apply another registered fungicide after applying Fitness Fungicide twice. See Brown rot blossom blight. Make up to 2 additional applications on a 10- to 14-day interval from the end of petal fall to harvest.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**STRAWBERRY AND LOW GROWING BERRIES (Subgroup 13-07G (except Cranberry))**

Low growing berries (subgroup 13-07G (except cranberry)) including bearberry, bilberry, lowbush blueberry, cloudberry, lingonberry, muntries, Partridgeberry, strawberry, cultivars, varieties, and/or hybrids of these

**Specific Use Restrictions:**

- **DO NOT** apply more than 16.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 4.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 4 applications per year.
- Fitness Fungicide may be applied at 0 day PHI (day of harvest).
- **DO NOT** apply more than 0.45 pound active ingredient propiconazole-containing product per acre per year.

Fitness Fungicide is most effective when applied and allowed to dry before a rainfall. For best results, sufficient water volume must be used to provide thorough coverage. Fitness Fungicide may be applied by either ground (a minimum of 20.0 gallons per acre) or aerial application (a minimum of 15.0 gallons per acre).

Target Disease	Application Rate (FI Oz/A)	Application Instructions
Anthracnose ( <i>Colletotrichum acutatum</i> ) Leaf spot ( <i>Cercospora fragariae</i> ) Powdery mildew ( <i>Sphaerotheca macularis</i> ) Leaf rust ( <i>Phragmidium potentillae</i> )	4.0	Begin applications when disease levels are not more than 5%. Apply 4.0 fluid ounces per acre of Fitness Fungicide up to 4 times on a 7-day interval. Make no more than 2 consecutive applications before rotating to another registered fungicide with a different mode of action.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**SUGARCANE**

**Restrictions**

- Dispose of used dip solution according to local state and Federal regulations.
- **DO NOT** use treated seed pieces for food or feed.

Target Disease	Application Rate (FI Oz/100gal water)	Application Instructions
Pineapple disease ( <i>Ceratocystic paradoxa</i> )	0.75 (1:17,000)	Apply to cut seed pieces using one of the following methods: <b>Cold water dip:</b> Immerse seed pieces to give thorough wetting, remove, and allow to drain. <b>Hot water dip:</b> Maintain water temperature at 125 °F (52 °C). Soak seed pieces for 20 to 30 minutes, remove, and allow to drain. <b>Conveyor belt treatment:</b> Treat seed pieces with a solution of Fitness Fungicide and water using in-line

**TREE NUTS (Crop Group 14)**

Almond (see specific directions), beechnut, Brazil nut, butternut, cashew, chestnut, chinquapin, filbert (see specific directions elsewhere on this label), hickory, macadamia, pecan (see specific directions elsewhere on this label), walnut

**Restrictions**

- **DO NOT** apply more than 32.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (8 fl oz/A) or 8 applications per year when applying at the lowest rate (4 fl oz/A).
- **DO NOT** graze livestock in treated areas or cut cover crop for feed.
- **DO NOT** apply more than 0.90 pound active ingredient propiconazole-containing product per acre per year.
- Preharvest Interval (PHI) is 60 days.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 15.0 gallons per acre). For best control of tree nut diseases, apply by ground application. Fitness Fungicide is most effective when applied and allowed to dry before a rainfall.

Target Disease	Application Rate (FI Oz/A)	Application Instructions
Foliar Diseases	4.0 to 8.0	Apply Fitness Fungicide at first sign of disease. Repeat on a 7- to 14-day

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**GRASSES GROWN FOR SEED (For use only in Idaho, Minnesota, Nebraska, Oregon, and Washington) Restrictions**

- **DO NOT** apply more than 32.0 fluid ounces per acre per year of Fitness Fungicide.
- **DO NOT** apply more than 8.0 fluid ounces per acre per application of Fitness Fungicide.
- **DO NOT** exceed 4 applications per year when applying at the highest rate (8 fl oz/A) or 8 applications per year when applying at the lowest rate (4 fl oz/A).
- **DO NOT** graze treated areas within 140 days of the last application.
- **DO NOT** apply within 20 days of harvest of seed (20 day PHI).
- **DO NOT** feed hay cut within 20 days of the last application.
- **DO NOT** apply more than 0.90 pound active ingredient propiconazole-containing product per acre per year.
- **DO NOT** apply to bermuda grass grown for seed.

Apply Fitness Fungicide by ground or aerial equipment (a minimum of 10.0 gallons per acre). Fitness Fungicide may also be applied by chemigation.

<b>Target Disease</b>	<b>Application Rate (FI)</b>	<b>Application Instructions</b>
Ergot stem disease Powdery mildew ( <i>Erysiphe graminis</i> ) Rusts ( <i>Puccinia</i> spp.) Selenophoma stem eyespot ( <i>Selenophoma</i> spp.)	4.0 to 8.0	Apply when Powdery mildew infections, Selenophoma infections, and/or Rust pustules are noticeable and increasing in number in late spring or early summer. Apply on a 14- to 21-day interval. To maximize control of severe Rust pressure, apply 8.0 fl oz/A and reapply at 14-day intervals until the seed is mature. Make the last application at least 20 days before seed matures. For bluegrass, it is important to begin applications early in the growing season.



**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**TURFGRASS AND ORNAMENTAL INSTRUCTIONS**  
**(Not for Turf and Ornamental use in California)**

**Product Information**

Fitness Fungicide is a systemic fungicide for use on turfgrass for the control of Dollar spot (*Sclerotinia homeocarpa*), Brown patch (*Rhizoctonia solani*), Anthracnose (*Colletotrichum graminicola*), Red thread (*Laetisaria fuciformis*), Pink patch (*Limonomyces roseipellis*), Rust (*Puccinia graminis*), Powdery mildew (*Erysiphe graminis*), Stripe smut (*Ustilago striiformis* and *Urocystis agropyri*), Summer patch (*Magnaporthe poae*), Necrotic ring spot (*Leptosphaeria korrae*), Spring dead spot (*Leptosphaeria korrae*, *Leptosphaeria narmari*, *Ophiosphaerella herpotricha*, *Gaeumannomyces graminis*), Take-all patch (*Gaeumannomyces graminis*), Leaf spot (*Bipolaris* spp., *Drechslera* spp.), Gray leaf spot (*Pyricularia grisea*), Pink snowmold (*Microdochium nivale*), Fusarium patch (*Fusarium nivale*), Gray snowmold (*Typhula* spp.), Yellow patch (*Rhizoctonia cerealis*), and Zoysia patch (*Rhizoctonia solani*).

Fitness Fungicide also controls numerous diseases on ornamentals and other landscape and nursery plantings, including Powdery mildews, Rusts, Leaf spots, Scabs, and Blights. Refer to the appropriate section for specified diseases and plants.

**Use Restrictions**

- **DO NOT** apply more than 1.79 lb a.i./A (equivalent to 0.45 gallon of product/A or 1.3 fl oz/1000 sq ft) per application.
- **DO NOT** apply more than 7.2 lb a.i./A (equivalent to 1.8 gallons of product/A or 5.3 fl oz/1000 sq ft) per calendar year.
- **Minimum Application Interval:** 14 days
- **DO NOT** use this product as a tree injection treatment.
- **DO NOT** use Fitness Fungicide in greenhouses.
- **DO NOT** graze animals on treated areas.
- **DO NOT** feed clippings from treated areas to livestock or poultry.
- **DO NOT** exceed 1.44 fluid ounces per 1000 square feet every 30 days on any variety of bermudagrass
- Chemigation: For turf and ornamental uses, **DO NOT** apply this product through any type of irrigation system.
- Bermudagrass can be sensitive to Propiconazole EC. Do not exceed 1.44 fl oz per 1000 sq ft every 30 days on any variety of bermudagrass. In Florida, do not apply Propiconazole EC to bermudagrass golf course greens when temperatures exceed 90°F.

**Use Precautions**

Failure to follow the directions for use and precautions on this label may result in plant injury or poor disease control.

**Tank Mixes of Fitness Fungicide for Turfgrass and Ornamental**

**Applications See Mixing Instructions Section**

Add E-Z Mix compatibility agent (3.0 pints per 100 gallons) to tank mixes that are incompatible. Observe all directions, precautions and limitations on the labels of all products used in the tank mix. Tank mixes or other applications of products referenced on this label are permitted only in those states where referenced products are registered.

**Uses**

**Turfgrass and Dichondra**

- Use Fitness Fungicide in a preventative disease control program.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to dry completely before mowing.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

- For control of foliar diseases, allow sprayed area to dry completely before irrigation.
- For control of soilborne diseases, Fitness Fungicide can be watered in immediately after application. Under conditions that are optimum for high disease pressure, use the higher rate and shorter interval.
- For optimum turfgrass quality and disease control, use Fitness Fungicide in conjunction with turf management practices that promote good plant health and optimum disease control.
- Evaluate spray additives prior to use. Label directions are based upon data obtained with no additives.
- Before using any fungicide, proper diagnosis of the organism causing the disease is important. Using diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

**Use Precautions**

- Important: Bermudagrass can be sensitive to Fitness Fungicide.
- In Florida, do not apply Fitness Fungicide to Bermudagrass golf course greens when temperatures exceed 90 °F.

**Specific Diseases, Rates and Application Timing**

Disease	(Fl Oz/1000 Sq Ft)	(Fl Oz/A)	Application Interval (Days)	Application Instructions
<b>Turfgrass</b>				
<b>Anthracnose</b> ( <i>Colletotricum graminicola</i> )	0.37 to 0.73	16.0 to 32.0	14 to 28	Apply when conditions favor disease development. When disease pressure is high, use a higher rate and shorter interval. For broad spectrum control, tank mix with a registered contact fungicide at the label rate. If disease is present, mix 0.75 fl oz of Fitness Fungicide per 1000 sq ft with the label rate of a registered contact fungicide.
<b>Brown patch</b> ( <i>Rhizoctonia solani</i> )			14 to 21	Begin application in May or June before disease is present. Tank mix with the label rate of a contact fungicide registered for control of Brown patch. Under conditions of high temperature and humidity, use a higher rate and shorter interval.
<b>Dollar spot</b> ( <i>Sclerotinia homeocarpa</i> )	0.18	8.0	14	Apply when conditions favor disease development. Tank mix with low label rate of a contact fungicide containing chlorothalonil that is

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

<b>Disease</b>	<b>(Fl Oz/1000 Sq Ft)</b>	<b>(Fl Oz/A)</b>	<b>Application Interval (Days)</b>	<b>Application Instructions</b>
	0.37	16.0	21 to 28	EPA-registered for use on turfgrass.
	0.37 to 0.75	16.0 to 32.0	14 to 28	If using the 0.37 to 0.75 fl oz/1000 sq ft rate without tank mixing, make no more than 3 consecutive applications for control of Dollar spot before rotating to an alternate EPA registered fungicide having a different mode of action.
<b>Fusarium patch</b> ( <i>Fusarium nivale</i> )	0.75 to 1.45	32.0 to 63.0	fall to early spring	Apply when conditions favor disease development.
<b>Gray leaf spot</b> ( <i>Pyricularia grisea</i> )	0.37 to 0.75	16.0 to 32.0	14	Apply when conditions favor disease development. If using the 0.37 fl oz/1000 sq ft rate, tank mix with a registered contact fungicide at the label rate.
<b>Melting out Leaf spot</b> ( <i>Bipolaris</i> spp.) ( <i>Drechslera</i> spp.)		16.0 to 63.0		Under light to moderate pressure, apply Fitness Fungicide to reduce the severity of Leaf spot and Melting out. For broad spectrum disease control, tank mix the 0.37 fl oz/1000 sq ft rate with a registered contact fungicide at the label rate.
<b>Necrotic ring spot</b> ( <i>Leptosphaeria korrae</i> )	1.45	63.0	Fall or spring	Apply in fall and/or the early spring depending upon local recommendations.
<b>Pink patch</b> ( <i>Limonomyces roseipellis</i> ) <b>Red thread</b> ( <i>Laetisaria fuciformis</i> )	0.37	32.0	14 to 21	Apply when conditions favor disease development.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

<b>Disease</b>	<b>(Fl Oz/1000 Sq Ft)</b>	<b>(Fl Oz/A)</b>	<b>Application Interval (Days)</b>	<b>Application Instructions</b>
<b>Powdery mildew</b> ( <i>Erysiphe graminis</i> ) <b>Rust</b> ( <i>Puccinia graminis</i> )	0.37 to 0.75	16.0 to 32.0	14 to 28	Apply when conditions favor disease development. If disease is present, use 0.75 fl oz of Fitness Fungicide/1000 sq ft.
<b>Snow mold, Gray</b> ( <i>Typhula</i> spp.) <b>Snow mold, Pink</b> ( <i>Microdochium nivale</i> )	0.75 to 1.45	32.0 to 63.0	Late fall	Make 1 application in the late fall before snow cover. Do not apply on top of snow. For optimum disease control, the 0.75 and 1.1 fl oz Fitness Fungicide rates must be tank mixed with chlorothalonil at label rates.
<b>Spring dead spot</b> ( <i>Leptosphaeria korrae</i> ), ( <i>Leptosphaeria narmari</i> ), ( <i>Ophiosphaerella herpotricha</i> ), ( <i>Gaeumannomyces graminis</i> )	1.45	63.0	30	Make 1 to 3 applications. If a single application is made, apply in September or October. For multiple applications, begin sprays in August.
<b>Sripe smut</b> ( <i>Ustilago striiformis</i> ), ( <i>Urocystis agropyri</i> )	0.37 to 0.75	16.0 to 32.0	Fall or spring	Apply once in the fall after turfgrass becomes dormant or in the early spring before turfgrass starts to grow.
<b>Summer patch</b>	0.75	32.0	14	Apply Fitness Fungicide beginning in April. Use the 1.5 fl oz/1000 rate on 28-day schedule and the 0.75 fl oz/1000 sq ft rate on a 14-day schedule.
<b>Poa patch</b> ( <i>Magnaporthe poae</i> )	1.45	63.0	28	

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

<b>Disease</b>	<b>(FI Oz/1000 Sq Ft)</b>	<b>(FI Oz/A)</b>	<b>Application Interval (Days)</b>	<b>Application Instructions</b>
<b>Take-all patch</b> ( <i>Gaeumanomyces graminis</i> )	0.75 to 1.45	32.0 to 63.0	Spring and fall	Apply Fitness Fungicide to reduce the severity of Take-all patch. Make fall applications in September and October or when night temperatures drop below 55 °F, and spring applications in April and May, depending on local recommendations.
<b>Yellow patch</b> ( <i>Rhizoctonia cerealis</i> )	1.10 to 1.45	48.0 to 63.0	Late fall	Make 1 application in the late fall before snow cover. Do not apply on top of snow. If using the 1.1 fl oz/1000 sq ft rate, tank mix with a registered contact fungicide at the label rate.
<b>Zoysia patch, Large patch of zoysia</b> ( <i>Rhizoctonia solani</i> )	1.10 to 1.45	48.0 to 63.0	Early fall	Make 1 application in the early fall (mid September to mid October) prior to development of disease symptoms. Consult local turfgrass extension experts to determine the optimum application timing for your area.

<b>Disease</b>	<b>(FI Oz/1000 Sq Ft)</b>	<b>(FI Oz/A)</b>	<b>Application Interval (Days)</b>	<b>Application Instructions</b>
<b>Dichondra</b>				
<b>Dichondra rust</b> ( <i>Puccinia dichondrae</i> )	0.75	32.0	14 to 21	Apply when conditions favor disease development.

**Establishment of Cool Season Turfgrass**

Fitness Fungicide controls many turfgrass diseases; its primary use is as a fungicide for use against the diseases listed on this label. As an additional benefit, Fitness Fungicide improves the rate of establishment when it is applied to cool season turfgrass seedlings or sod.

**New Seedlings:** Apply 0.35 fluid ounce per 1000 square feet at the 2- to 3-leaf stage of growth for faster root development and top growth.

**Sod:** Apply 0.35 fluid ounce per 1000 square feet 2 to 6 weeks before cutting for increased sod knitting and faster establishment after laying.

**Ornamental Plants**

Use Fitness Fungicide in a preventative disease control program. To determine the use directions for controlling a disease on an ornamental plant species, select the plant species in Table 1. The number(s) in parentheses following the listed plant species refers to the disease(s) controlled in

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

Table 2. Find the disease in Table 2. The letter in brackets following the disease refers to the application regime in Table 3.

Allow spray to dry before applying overhead irrigation.

Optimum benefit of Fitness Fungicide is obtained when used in conjunction with sound disease management practices.

Fitness Fungicide may be used at rates of 0.75 to 8.7 fluid ounces per 100 gallons of water for disease control in ornamentals (see Tables 1, 2 and 3). For best control, begin applications before disease development. For general disease control in landscapes, apply 2.2 to 3.0 fluid ounces per 100 gallons water every 21 days.

**Use Restrictions**

- To avoid possible illegal residues, do not apply to apple, Bartlett pear, cherry, citrus, nectarine, peach, pecan, plum, or walnut trees that will bear harvestable fruit within 12 months.
- For outdoor uses, up to 2.0 gallons of Fitness Fungicide may be applied per acre per calendar year.
- Do not apply Fitness Fungicide to African violets, begonias, Boston fern, or geraniums.

**Use Precautions**

- Plant tolerances to Fitness Fungicide have been found acceptable for the specific genus and species of plants listed under the Directions for Use.
- Other plant species may be sensitive to Fitness Fungicide and diseases other than those listed may not be controlled.
- Before using Fitness Fungicide on other plants, first test Fitness Fungicide on a small scale basis.
- Apply according to listed rates for a particular disease type, i.e. Rust, Powdery mildew, etc., and evaluate for phytotoxicity and disease control prior to widespread use.

**Table 1. Ornamental Plant Species**

**Number in parentheses (-) refer to diseases controlled in**

**Table 2. Herbaceous Ornamentals**

calendula (4a)	English ivy (3e)	marigold (3a)	sweet william (3k)
carnation (5f)	gomphrena (3a)	monarda (4c)	( <i>Dianthus barbatus</i> )
chrysanthemum (2a)	impatiens (3a, 3b, 4a)	phlox (4c)	zinnia (4c)
delphinium (4a)	iris (5d)	snapdragon (5d)	

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**Woody Ornamentals**

amelanchier (4d)	douglas fir (5b)	maple (3e, 4f)	roses (3g, 4e, 5c)
ash (4c)	elm (4c)	oaks (3p)	(outdoor use only)
azalea (2c, 4b)	euonymus (3e, 4c)	pinos (1a, 1c)	shasta fir (5e)
bayberry (3n)	hawthorn (5a)	poplars (5b)	sweetgum (3b, 3c, 3n)
camellia (3e)	holly (3r)	pyracantha (3o)	sycamore (3e)
cotoneaster (3l)	juniper (1a)	red tip photinia (3l)	tulip tree (3e, 4a)
crabapple (3c, 3g, 4c, 5a)	lilac (4c)	raphiolepis (3e, 3i)	wax myrtle (3n)
crape myrtle (4a)	linden (3e, 3b, 4b)	rhododendron (2c, 3n)	
dogwood (3h, 4c)	magnolia (3e, 4b)		

## Table 2. Plant Diseases

Letters in brackets [-] refer to application regimes in Table 3.

1. Conifer blights
  - a. *Phomopsis juniperovora* (Phomopsis blight) [B]
  - b. *Sirococcus strobolinus* (Tip blight) [D]
  - c. *Sphaeropsis sapinea* (Diplodia tip blight) [B]
2. Flower blight
  - a. *Ascochyta chrysanthemi* (Ray blight) [C]
  - b. *Molinia* spp. [A]
  - c. *Ovulinia* spp. [B]
3. Leaf blights/spots
  - a. *Alternaria* spp. [B]
  - b. *Cercospora* spp. (Brown leaf spot) [C]
  - c. *Cladosporium* spp. (Scab) [C]
  - d. *Coccomyces hiemalis* [A]
  - e. *Collectrichum* spp. [B]
  - f. *Cristulariella* spp. (Zonate leaf spot) [C]
  - g. *Diplocarpon rosae* (Blackspot) [B]
  - h. *Discula* spp. (Anthracnose) [A]
  - i. *Fabraea maculata* (syn. *Entomosporium maculata*) [B]
  - j. *Gnomonia leptostyla* (Anthracnose) [C]
  - k. *Heterosporium echinulatum* [B]
  - l. *Mycosphaerella caryigena* (Downy spot) [C]
  - m. *Mycosphaerella fructicola* (Greasy spot) [E]
  - n. *Septoria* spp. (Leaf scorch) [C]
  - o. *Spilocaea pyracanthae* [B]
  - p. *Tubakia dryina* [D]
  - q. *Venturia inaequalis* (Scab) [A]
  - r. *Rhizoctonia web blight* [B]
4. Powdery mildew
  - a. *Erysiphe* spp. [B]
  - b. *Microsphaera* spp. [C]
  - c. *Oidium* spp. [B]
  - d. *Podosphaera* spp. [B]
  - e. *Sphaerotheca pannosa* [B]
  - f. *Phyllactinia* spp. [B]
5. Rust
  - a. *Gymnosporangium juniperi-virginianae* [A]
  - b. *Melampsora occidentalis* [D]
  - c. *Phragmidium* spp. [B]
  - d. *Puccinia* spp. [B]
  - e. *Pucciniastrum goeppertianum* [D]
  - f. *Uromyces dianthi* [B]



**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**Table 3. Application Regimes**

**[A]** Mix 0.75 to 1.5 fluid ounces of Fitness Fungicide in 100 gallons of water and apply as a full coverage spray to the point of drip. Reapply every 14 to 21 days during the period of primary infection. If disease is present, tank mix with an EPA-registered contact fungicide. For Flower blight, apply Fitness Fungicide when there is 5 to 10% bloom and again at 70 to 100% bloom. For dogwoods, apply the 0.75 to 1.5 fluid ounces rate every 14 days, or apply 3.0 fluid ounces of Fitness Fungicide every 28 days.

**[B]** Mix 1.8 to 3.0 fluid ounces of Fitness Fungicide in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply as needed, beginning when conditions favor disease development. For Blackspot, apply in tank mix with a registered contact fungicide labeled for Blackspot. For Calendula, apply every 30 days. For Diplodia tip blight, make a total of 3 applications every 14 days prior to the major period of infection. For Juniper phomopsis blight, make an initial application as soon as junipers start to grow and reapply every 14 to 21 days during the period of active growth.

**[C]** Mix 3.0 to 4.5 fluid ounces of Fitness Fungicide in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 30 days, beginning when conditions favor disease development. For pecans, apply the 12.0 fluid ounces rate. Beginning at bud break, make a total of 3 applications 14 days apart. For walnuts, apply 3.0 fluid ounces every 14 to 21 days. For Ray blight, apply 4.5 fluid ounces every 7 days or 7.5 fluid ounces every 14 days. For impatiens, bayberry, linden, magnolia, sweetgum and wax myrtle, the maximum use rate is 8.0 fluid ounces.

**[D]** Mix 6.0 fluid ounces of Fitness Fungicide in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply every 14 to 28 days beginning when conditions favor disease development. For Douglas fir needle rust, apply once in May. For Tip blight, make an initial application in mid- to late winter, and 3 additional applications at 2-month intervals.

**[E]** Mix 7.5 to 8.7 fluid ounces of Fitness Fungicide in 100 gallons of water and apply as a full coverage spray to the point of drip. Apply within the June to August time period.

**FITNESS® FUNGICIDE**  
**EPA REG. NO. 34704-1031**

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:**

**Nonrefillable container.** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For containers up to 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For containers greater than 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and

**FITNESS® FUNGICIDE**  
**EPAREG. NO. 34704-1031**

local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.**

**FITNESS® FUNGICIDE  
EPA REG. NO. 34704-1031**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND  
LIABILITY**

**BEFORE BUYING OR USING THIS PRODUCT**, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Fitness and Initiate are registered trademarks of Loveland Products, Inc.

**FITNESS® FUNGICIDE  
EPA REG. NO. 34704-1031**

**FORMULATED FOR  
LOVELAND PRODUCTS, INC.  
P.O. BOX 1286, GREELEY, COLORADO 80632-1286**