



DIMENSION™ 0.27G Turf and Ornamental Herbicide

For season-long control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf (including golf course fairways, roughs and tee boxes) and container and field-grown nursery ornamentals.

- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

- For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

In the state of New York, this product may be applied only by commercial applicators at no more than 185 lb (0.5 lb active ingredient) per acre (4.25 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Active Ingredient:

dithiopyr: 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester0.27%
Other Ingredients99.73%
Total100.00%

This product contains 0.135 pounds of the active ingredient, dithiopyr, per 50 pound bag.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back or container/side panel for additional Precautionary Statements

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet

| FIRST AID | |
|---|--|
| If in eyes | <ul style="list-style-type: none">Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency medical treatment information. | |

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff may adversely affect aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Use Directions for Turf

LESCO Dimension® 0.27G Turf and Ornamental Herbicide provides seasonal-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, (including golf course fairways, roughs, and tee boxes), and container and field-grown nursery ornamentals. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

Use Precautions for Turf

- Do not use this product on golf greens.
- Do not apply this product to Colonial bentgrass.
- For best results, cultural practices that disturb the soil, such as core, spike, hydro-aerification, or verticutting, and topdressing should be done before application of this product. This product should not be applied until turfgrasses have recovered from stresses related to these or other mechanical operations. Irrigation or rainfall within 7 days after application will improve the performance of this product.
- Reseeding or Sprigging:** Reseeding, overseeding, or sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 12 weeks from the time of application. If overseeding with perennial ryegrass, delay overseeding at least 8 weeks after application. When reseeded or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact.
- This product may be used on seeded, sodded, or sprigged turfgrass that is well established. Allow the turf to develop a good root system and uniform stand before application. Use of this product on turfgrass not well established, weakened by lack of moisture, chemical injury, insects, disease, or temperature extremes that cause stress will increase the potential for injury. Allow turfgrass weakened by these stresses to recover before application of this product.
- This product may be applied as a single application or as a split application in the spring, summer or fall.
- This product will prevent germination of annual bluegrass.
- This product will not affect established annual bluegrass. If maintenance of annual bluegrass is desired, do not use this product during time of annual bluegrass germination. For AZ, CA, NV, OR, WA, NM, ID, MT and UT, this product may contribute to thinning or stand reduction in established stands of annual bluegrass.
- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow domestic animals to feed or forage on the treated turf or clippings from treated turf.

When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

| Cool-Season Grasses | |
|-----------------------|-----------------------------|
| bentgrass, creeping † | <i>Agrostis stolonifera</i> |
| bluegrass, Kentucky | <i>Poa pratensis</i> |
| fescue, fine †† | <i>Festuca rubra</i> |
| fescue, tall | <i>Festuca arundinacea</i> |
| ryegrass, perennial | <i>Lolium perenne</i> |

| Warm-Season Grasses | |
|---------------------|--------------------------------|
| bahiagrass | <i>Paspalum notatum</i> |
| bermudagrass | <i>Cynodon dactylon</i> |
| buffalograss | <i>Buchloe dactyloides</i> |
| carpetgrass | <i>Axonopus affinis</i> |
| centipede grass | <i>Eremochloa ophiuroides</i> |
| kikuyugrass | <i>Pennisetum clandestinum</i> |
| seashore paspalum | <i>Paspalum vaginatum</i> |
| St. Augustinegrass | <i>Stenotaphrum secundatum</i> |
| zoysiagrass | <i>Zoysia japonica</i> |

† Do not apply this product to Colonial bentgrass. Use of this product on certain varieties of creeping bentgrass, such as 'Carmen', 'Cohansey', or 'Seaside' will result in undesirable turf injury. Not all varieties of creeping bentgrass have been tested.

†† The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Waldorf', and 'Waldina'.

Application Equipment and Instructions

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Calibrate application equipment prior to usage, and apply this product uniformly over the turfgrass area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction of application. Avoid the use of spreaders that apply granular materials in narrow rows or concentrated bands. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

Preemergence Control of Annual Grasses and Annual Broadleaf Weeds

Used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established weeds, except for crabgrass in early stages of development, but only if application is made to crabgrass prior to the 3-leaf stage of growth or first tiller. Use the higher recommended use rates where weed infestations are dense and/or applications are made early and longer residual control is desirable.

| Weeds Controlled | | Weeds Controlled and Use Rate Recommendations | |
|--|---|---|-------------------|
| | | Suggested Use Rate Per Application | |
| | | Pounds/Acre | Pounds/1000 sq ft |
| barley brome crabgrass, large crabgrass, smooth crabgrass, southern dallisgrass (seedling) foxtail, giant foxtail, green foxtail, yellow | | 100 to 150 †† | 2.3 to 3.4 † |
| barnyardgrass chickweed clover, hop crabgrass, tropical crowfootgrass dichondra, Carolina lespedeza, common kikuyugrass Mary's grass marestail medic, black mulberry weed mustard oats, wild oxalis, buttercup oxalis (creeping woodsorrel) oxalis (yellow woodsorrel) pineappleweed ryegrass, annual ryegrass, perennial sandbur smutgrass | | 150 to 185 †† | 3.4 to 4.25 † |
| bluegrass, annual bittercrass carpetweed chickweed geranium, Carolina henbit knotweed, prostrate lespedeza, common marestail medic, black mulberry weed mustard oxalis, buttercup parsley-piert pigweed, redroot pineappleweed purslane, common rocket, London sheepspurge speedwell, com spurge, garden spurge, prostrate spurge, spotted | henbit knotweed, prostrate parsley-piert pigweed, redroot purslane, common rocket, London sheperd's-purse | 185 †† | 4.25 †† |

† Higher use rates are recommended where weed infestations are heavy and/or dense.

†† Use of this product at this rate will provide only partial control and/or suppression of listed weeds.

Poa Annua (annual bluegrass) Control

Apply LESCO Dimension® 0.27G Turf and Ornamental Herbicide for preemergence control of Poa annua (annual bluegrass) at a rate of 0.38 to 0.5 lbs active ingredient per acre. This is equivalent to 150 to 185 pounds per acre. Use the higher recommended use rates where weed infestations are dense and/or applications are made early and longer residual control is desirable.

- Apply 6 to 8 weeks before overseeding bermudagrass with perennial ryegrass. Specific to perennial ryegrass; not for use on Poa trivialis or bentgrass.
- Minimum seeding rate of perennial ryegrass is 400 lbs per acre.
- Use limited to fairways and roughs
- Perennial varieties of Poa annua (var. repens) may not be controlled as well as the true annual.
- Do not apply earlier than 16 weeks after over-seeding unless injury to the ryegrass can be tolerated.
- A follow-up treatment 16 weeks after overseeding offers an early season crabgrass treatment and helps suppress some winter annual broadleaf weeds.

Restrictions:

- Do not** apply more than 185 pounds of this product per acre per application and no more than 3 applications per year (555 pounds) of this product per acre per year.
- In New York State, this product may only be used by Commercial Applicators at no more than 185 pounds per acre (4.25 lb/1000 sq ft), (0.5 lb active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

Ornamentals (Landscape, Field Grown, and Container Grown)

LESCO Dimension® 0.27G Turf and Ornamental Herbicide provides preemergence control of listed annual grass and broadleaf weeds in ornamental plantings. Ornamental plants on this label have shown tolerance to this product when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

Use RatesRefer to the following table for treatment dosage for treating small areas.

| Pounds Product/Acre | Ounces Product/1000 Square Feet | Ounces (Grams) Product/100 Square Feet |
|---------------------|---------------------------------|--|
| 185 | 68 | 6.8 (193 grams) |

Application Timing and Use Recommendations

This product is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. To be effective, LESCO Dimension® 0.27G Turf and Ornamental Herbicidemust be activated by 1/2 inch or more of rainfall or irrigation prior to germination of target weeds. Once the treatment is activated, avoid excessive soil disruption that may break down the herbicide barrier. Minimal surface disruption such as raking should not break down the herbicide barrier.

Weeds Controlled

Used as directed, this product will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established weeds, except for crabgrass in early stages of development, but only if application is made to crabgrass prior to the 3-leaf stage of growth or first tiller. Therefore, the area to be treated should be free of weeds prior to application. Repeat applications may be made during the growing season at 3 to 4 month intervals to maintain weed control.

Weeds controlled or suppressed by LESCO Dimension 0.27G when applied at 185 pounds per acre:

| Annual Grasses | |
|-------------------------------|----------------------------------|
| Common Name | Scientific Name |
| barley | <i>Hordeum spp.</i> |
| barnyardgrass | <i>Echinochloa crus-galli</i> |
| bluegrass, annual | <i>Poa annua</i> |
| brome | <i>Bromus spp.</i> |
| crabgrass, large | <i>Digitaria sanguinalis</i> |
| crabgrass, smooth | <i>Digitaria ischaemum</i> |
| crabgrass, southern | <i>Digitaria ciliaris</i> |
| crowfootgrass | <i>Dactyloctenium aegyptium</i> |
| dallisgrass (seedling) | <i>Paspalum dilatatum</i> |
| foxtail, giant | <i>Setaria faberi</i> |
| foxtail, green | <i>Setaria verdi</i> |
| foxtail, yellow | <i>Pennisetum glaucum</i> |
| goosegrass | <i>Elyusine indica</i> |
| kikuyugrass | <i>Pennisetum clandestinum</i> |
| oats, wild | <i>Avena fatua</i> |
| ryegrass (annual & perennial) | <i>Lolium spp.</i> |
| sandbur | <i>Cenchrus spp.</i> |
| smutgrass | <i>Sporobolus indicus</i> |
| Broadleaf Weeds | |
| Common Name | Scientific Name |
| bittercrass | <i>Cardamine spp.</i> |
| carpetweed | <i>Mollugo verticillata</i> |
| chickweed | <i>Stellaria spp.</i> |
| geranium, Carolina | <i>Geranium carolinianum</i> |
| henbit | <i>Lamium spp.</i> |
| knotweed, prostrate | <i>Polygonum saviculare</i> |
| lespedeza, common | <i>Lespedeza striata</i> |
| marestail | <i>Hippuris vulgaris</i> |
| medic, black | <i>Medicago lupulina</i> |
| mulberry weed | <i>Fatoua villosa</i> |
| mustard | <i>Brassica spp.</i> |
| oxalis, buttercup | <i>Oxalis pes-caprae</i> |
| parsley-piert | <i>Achemilla arvensis</i> |
| pigweed, redroot | <i>Amaranthus retroflexus</i> |
| pineappleweed | <i>Matricaria matricarioides</i> |
| purslane, common | <i>Portulaca oleracea</i> |
| rocket, London | <i>Sisymbrium irio</i> |
| sheepspurge | <i>Capsella bursa-pastoris</i> |
| speedwell, com | <i>Veronica arvensis</i> |
| spurge, garden | <i>Euphorbia hirta</i> |
| spurge, prostrate | <i>Euphorbia humistrata</i> |
| spurge, spotted | <i>Euphorbia maculata</i> |
| woodsorrel, creeping | <i>Oxalis corniculata</i> |
| woodsorrel, yellow | <i>Oxalis stricta</i> |

Application Equipment and Instructions

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction of application. Avoid streaking, skips, or overlaps during application.

Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules. Shortly after application apply overhead irrigation to activate the herbicide and wash LESCO Dimension® 0.27G Turf and Ornamental Herbicide from plant surface onto soil surface.

Use Precautions for Ornamental Plantings:

- Only apply this product to established ornamentals as directed in the Tolerant Ornamental table below.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply around ornamental plants that have been weakened or are under stress (due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or diseases).
- Do not make applications when weather conditions favor drift to non-target areas. This product may injure foliage of non-target plants.
- Do not apply this product on grasses grown for seed.
- Do not graze livestock or feed forage cut from areas treated with this product.
- Do not apply this product in enclosed structures and greenhouses.
- Do not apply another herbicide within 4 weeks of application of this product.
- Do not apply this product to bare roots of ornamental plants as injury may result. Following transplanting, care must be taken that soil or planting mixes have settled firmly through irrigation, rainfall or packing and that there are no cracks that would allow direct contact of this product to the plant roots or plant injury may occur.

Treatment of Ornamental species not listed on this label:

It is impossible to evaluate tolerance to this product on all ornamental species or varieties or under all growing conditions. Users who wish to use LESCO Dimension® 0.27G Turf and Ornamental Herbicide on species not recommended on this label may determine the suitability for use by treating a small number of plants at a recommended rate under your growing conditions. Treated plants should be observed for any symptoms of herbicidal injury such as foliar damage, reduced vigor or stand reduction during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious.

Restrictions:

- Do not** apply more than 185 pounds of this product per acre per application and no more than 3 applications (555 lb) of this product per acre per year.
- In New York State, this product may only be used by Commercial Applicators at no more than 185 pounds per acre (4.25 ounces/1000 sq ft), (0.5 lb active ingredient per acre) per year; and is prohibited from use in Nassau and Suffolk Counties.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.
Pesticide Storage: Store this product only in its original container. In case of spill, sweep up material and dispose of material according to label directions.
Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.
Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities .

Net Weight: 50 lb (22.7 kg)

EPA REG. NO. 62719-429-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 69759-OR-002 (MAS) 69759-OR-004 (BAN), 93243-TX-1.

First letter of lot code indicates manufacturing site.

F1560

Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

**702265-PG1
PP**



DIMENSION™ 0.27G Turf and Ornamental Herbicide

Tolerant Ornamentals

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials.

| Name | Tolerant Cultivars | Acceptable Application Method Noted by a (X) | | Application to Field/Landscape or Container Grown Ornamentals Noted by a (X) | |
|---|---|--|----------|--|------------------------|
| | | Over the Top | Directed | Container Grown | Field/ Landscape Grown |
| ice plant, rosea (<i>Drosanthemum floribundum</i>) | | X | X | X | X |
| ice plant, white trailing (<i>Delosperma alba</i>) | | X | X | X | X |
| impatiens (<i>Impatiens spp.</i>) | | | X | | X |
| iris (Iris spp) | dwarf blue wedgewood | | X | | X |
| ivy, English (<i>Hedera helix</i>) | Bulgaria thornedale | | X | | X |
| jasmine, Asian (<i>Trachelospermum asiaticum</i>) | | | X | | X |
| jasmine, star (<i>Trachelospermum jasminoides</i>) | | | X | | X |
| juniper (<i>Juniperus spp.</i>) | Arcadia Armstrong bar harbor blue chip blue Pacific blue rug blue star broadmoor buffalo calgary carpet emerald sea emerald spreader endora compacta fruitlandi green gold tip hetzi hughes Manhattan blue parsoni pfitzeriana plumosa Prince of Wales procumbens dwarf San Jose sargent blue sargent green scandia scopulorum moonglow spartan tamariscifolia weberi Youngstown Yukon belle | | X | | X |
| juniper (<i>Juniperus chinensis</i>) | gold coast | X | X | X | X |
| juniper (<i>Juniperus procumbens nana</i>) | prostrate Japanese garden | X | X | X | X |
| juniper (<i>Juniperus sabina tamariscifolia</i>) | tamarix | X | X | X | X |
| juniper (<i>Juniperus scopulorum</i>) | scopulorum skyrocket | X | X | X | X |
| king palm (<i>Archontophoenix cunninghamiana</i>) | | X | X | X | X |
| laurel, Carolina cherry (<i>Prunus caroliniana</i>) | bright n' tight | X | X | X | X |
| laurel, mountain (<i>Kalmia latifolia</i>) | | | X | | X |
| leucothoe (<i>Leucothoe fontanesiana</i>) | | | X | | X |
| ligustrum, Japanese (<i>Ligustrum japonicum</i>) | | | X | | X |
| maple, amur (<i>Acer ginnala</i>) | emerald elf | X | X | X | X |
| maple, Japanese (<i>Acer palmatum</i>) | | | X | | X |

| | | | | | |
|--|---|---|---|---|---|
| maple, Norway (<i>Acer platanoides</i>) | | | X | | X |
| maple, red (<i>Acer rubrum</i>) | red sunset | X | X | X | X |
| maple, silver (<i>Acer saccharinum</i>) | | | X | | X |
| maple sugar (<i>Acer saccharum</i>) | | | X | | X |
| marigold (<i>Tagetes patula</i>) | honeycomb variegata wheeleris dwarf | | X | | X |
| mock orange (<i>Philadelphus spp</i>) | golden snowflake double white | | X | | X |
| mondo grass (<i>Ophiopogon japonicus</i>) | | | X | | X |
| moss rose (Portulaca grandiflora) | sunny side | | X | | X |
| mountainash (<i>Sorbus aucuparia</i>) | | | X | | X |
| myrtle, crape (<i>Lagerstroemia indica</i>) | faurei langer muskogee standard pink | | X | | X |
| myrtle, wax (<i>Myrica californica</i>) | | | X | | X |
| narcissus (<i>Narcissus spp.</i>) | | | X | | X |
| oak, laurel (<i>Quercus laurifolia</i>) | | | X | | X |
| oak, pin (<i>Quercus palustris</i>) | | | X | | X |
| oak, red (<i>Quercus rubra</i>) | | | X | | X |
| oak, southern (<i>Quercus virginiana</i>) | | | X | | X |
| oak, willow (<i>Quercus phellos</i>) | | | X | | X |
| oleander (<i>Nerium oleander</i>) | hardy red petite pink sister agnes | | X | | X |
| osteospermum (<i>Osteospermum fruticosum</i>) | whirligig | | X | | X |
| pachysandra (<i>Pachysandra terminalis</i>) | | | X | | X |
| palm, king (<i>Archontophoenix cunninghamiana</i>) | | | X | | X |
| palm, queen (<i>Syagrus romanzoffianum</i>) | | X | X | X | X |
| pampas grass (<i>Cortaderia sellowiana</i>) | | | X | | X |
| pansy (<i>Viola x wittrockiana</i>) | | | X | | X |
| paper flower (<i>Bougainvillea glabra</i>) | Barbara Karst | X | X | X | X |
| peach (<i>Prunus persica</i>) | | | X | | X |
| periwinkle, dwarf (<i>Vinca minor</i>) | | | X | | X |
| petunia (<i>Petunia x hybrida</i>) | picoti | | X | | X |
| photinia, red tip (<i>Photinia x fraseri</i>) | | | X | | X |
| piers (<i>Pieris taiwanensis</i>) | | | X | | X |
| piers, Japanese (<i>Pieris japonica</i>) | mountain fire | | X | | X |

| | | | | | |
|--|--|---|---|---|---|
| pine, Afghan (<i>Pinus eldarica</i>) | | X | X | X | X |
| pine, Austrian black (<i>Pinus nigra</i>) | | X | X | X | X |
| pine, Japanese black (<i>Pinus thunbergii</i>) | | | X | | X |
| pine, loblolly (<i>Pinus taeda</i>) | | | X | | X |
| pine, longleaf (<i>Pinus palustris</i>) | | | X | | X |
| pine, mugo or Swiss Mt. (<i>Pinus mugo</i>) | | | X | | X |
| pine, Scotch (<i>Pinus sylvestris</i>) | | | X | | X |
| pine, slash (<i>Pinus elliotii</i>) | | | X | | X |
| pine, Virginia (<i>Pinus virginiana</i>) | | | X | | X |
| pine, white (<i>Pinus strobus</i>) | | X | X | X | X |
| pineapple, guava (<i>Feijoa sellowiana</i>) | | | X | | X |
| pittosporum, tobira (<i>Pittosporum tobira</i>) | Wheeler's dwarf | X | X | X | X |
| plumbago, cape (<i>Plumbago auriculata</i>) | royal cape | X | X | X | X |
| potentilla (<i>Potentilla fruticosa</i>)) (<i>Potentilla nepalensis</i>) | abbotswood | | X | | X |
| privet (<i>Ligustrum x vicaryi</i>) (<i>Ligustrum japonicum</i>) | golden vicary regal texanum yellow tipped | | X | | X |
| privet, glossy (<i>Ligustrum lucidum</i>) | | | X | | X |
| Pyracantha or firethorn (<i>Pyracantha x 'Gnome'</i>) (<i>Pyracantha coccinea</i>) (<i>Pyracantha koidzumii</i>) | gnome lalandei victory | | X | | X |
| queen palm (<i>Arecastrum rammanzoffianum</i>) | | | X | | X |
| quince, Japanese (<i>Chaenomeles japonica</i>) | | | X | | X |
| Rhododendron (<i>Rhododendron spp</i>) | album Cunningham-white PJM purple gem silvery pink | | X | | X |
| rhododendron, Carolina (<i>Rhododendron carolinianum</i>) | | | X | | X |
| rhododendron, catawba (<i>Rhododendron catawbiense</i>) | | | X | | X |
| rhododendron, rhodie max – rosebay (<i>Rhododendron maximum</i>) | | | X | | X |
| ribbon grass (<i>Phalaris arundinacea</i>) | | | X | | X |
| rockcress (<i>Arabis caucasica</i>) | snowcap | | X | | X |

| | | | | | |
|---|--|---|---|---|---|
| rose (<i>Rosa banksiae</i>) | luta | | X | | X |
| rose, knockout shrub (<i>Rosa spp. hybrid</i>) | knockout | X | X | X | X |
| rosemary (<i>Rosmarinus officinalis</i>) | | | X | | X |
| rosemary, bog (<i>Andromeda polifolia</i>) | nana | | X | | X |
| salvia (<i>Salvia farinacea</i>) | rhea | | X | | X |
| sedum (<i>Sedum spurium</i>) | dragon blood red red carpet yellow | | X | | X |
| smoketree, royal purple (<i>Cotinus coggygria</i>) | royal purple | | X | | X |
| snapdragon (<i>Antirrhinum spp.</i>) | | | X | | X |
| snowball, common (<i>Viburnum opulus</i>) | sterile | X | X | X | X |
| sourwood (<i>Oxydendrum arboreum</i>) | | | X | | X |
| spiraea (<i>Astilbe X arendsii</i>) | fanall | | X | | X |
| spiraea (<i>Spiraea spp.</i>) | Anthony Waterer red dolchica froebeli pink goldenflame red snowmound white Van Houtte white | | X | | X |
| spiraea, garland (<i>Spiraea x arguta</i>) .) | | | X | | X |
| spruce, Black Hills (<i>Picea glauca var densata</i>) | | | X | | X |
| spruce, Colorado blue (<i>Picea pungens</i>) | glauca | X | X | X | X |
| spruce, Norway (<i>Picea abies</i>) | | | X | | X |
| spruce, white (<i>Picea glauca</i>) | conica | | X | | X |
| sweetflag grass (<i>Acorus calamus</i>) | | | X | | X |
| sweetgum (<i>Liquidambar styraciflua</i>) | | | X | | X |

Net Weight: 50 lb (22.7 kg)

EPA REG. NO. 62719-429-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 69759-OR-002 (MAS) 69759-OR-004 (BAN), 93243-TX-1.

First letter of lot code indicates manufacturing site.

F1560
Distributed by: LESCO, Inc.
1385 East 36th Street
Cleveland, OH 44114-4114

702265-PG3
PP



DIMENSION™ 0.27G Turf and Ornamental Herbicide

Tolerant Ornamentals

When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials.

| Name | Tolerant Cultivars | Acceptable Application Method Noted by a (X) | | Application to Field/Landscape or Container Grown Ornamentals Noted by a (X) | |
|--|---|--|-----------------------|--|------------------------|
| | | Over the Top | Directed | Container Grown | Field/ Landscape Grown |
| sycamore, American (<i>Platanus occidentalis</i>) | | | X | | X |
| tea tree, New Zealand (<i>Leptospermum scoparium</i>) | ruby glow | X | X | X | X |
| tree fern (tiki fern) (<i>Asparagus virgatus</i>) | | | X | | X |
| trumpet flower or Carolina Jessamine (<i>Gelsemium sempervirens</i>) | | | X | | X |
| tufted hairgrass (<i>Deschampsia caespitosa</i>) | | | X | | X |
| verbena, shrub (<i>Lantana sellowiana</i>) | | | X | | X |
| Verbena, St. Paul's (<i>Verbena peruviana</i>) | St. Paul | | X | | X |
| viburnum (<i>Viburnum spp.</i>) | American cranberry bush arrowood European cranberry bush linden Mohican wright | | X X X X X | | X X X X |
| vinca (periwinkle) (<i>Vinca minor</i>) | | | X | | X |
| windmill palm (<i>Trachycarpus fortunei</i>) | | | X | | X |
| xylosma (<i>Xylosma congestum</i>) | | | X | | X |
| yarrow (<i>Achillea spp.</i>) | | | X | | X |
| yaupon (<i>Ilex vomitoria</i>) | dwarf | | X | | X |
| yew (<i>Taxus cuspidata</i>) (<i>Taxus x media</i>) | capitata denisiformis | X | X X | X | X X |

†Ornamental species only. Do not apply this product directly to plants that are grown for food.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use, and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

LESCO is a registered trademark and the sweeping design is a trademark of LESCO Inc. DIMENSION is a registered trademark of Dow AgroSciences LLC. SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC. N:\Regulatory\WPI\Regul Private\MstrLbl\ERIEVIEW ADDRESS\Landscape Style Master Lbls

(080316) Rev. 8/4/22 VT

Net Weight: 50 lb (22.7 kg)

EPA REG. NO. 62719-429-10404

EPA EST. NO. 961-PA-01 (L), 961-IL-01 (D), 8378-IN-1 (K), 9404-FL-1 (C), 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (E), 82757-OH-001 (M), 82757-FL-001 (S), 82757-MA-002 (H), 89135-AZ-001 (A), 69759-OR-002 (MAS) 69759-OR-004 (BAN), 93243-TX-1.

First letter of lot code indicates manufacturing site.

F1560

Distributed by: **LESCO, Inc.**
1385 East 36th Street
Cleveland, OH 44114-4114

702265-PG4

PP