FIFRA §2(ee) RECOMMENDATION

FOR USE IN CONNECTICUT, DELAWARE, MARYLAND, MASSACHUSETTS, NEW JERSEY, NEW YORK, PENNSYLVANIA, RHODE ISLAND AND VIRGINIA ONLY

FOR THE SUPPRESSION OF SPOTTED LANTERNFLY ON APPLES, GRAPES AND PEACHES





EPA Reg. No. 10163-324

ACCEPTED

Sept 27, 2019

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

Doc id: 564240

2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product.

Crops	Pest	Rate	COMMENTS
Apples Grapes Peaches	Spotted lanternfly (SLF),	Apply 1- 2% v/v per 100 gallons at the rate of 100 gallons of water per acre.	Do not apply this product to Calmeria or Italia varieties of grapes. Concentrate (air blast) applications are specified with a maximum volume of 175 gallons per acre on peaches. Initiate applications as soon as pest is observed, or as needed to avoid economic injury. Repeat applications at one-week intervals, or as needed, to control infesting or increasing pest populations. Do not make repeat applications at less than 7-day intervals, or make more than 3 sequential applications at 7- to 14-day intervals unless your experience has shown the use pattern to be non-injurious to the type and the development stage of plant(s) to be treated under local growing conditions.

Application Guidelines:

This product may be applied with most types of ground or aerial spray equipment. Do not use with pulse or thermal fogger applicators. Do not apply through any type of irrigation system.

- ☐ This product must come into contact with the targeted soft-bodied pests or powdery mildew to be effective. Complete coverage of pests is essential for maximum control. Direct sprays to contact the pests. In general, adequate coverage requires a minimum of 50 gallons of total spray volume per acre. Higher volumes may be required on larger perennial crops.
- Apply M-Pede solutions to wet (minimize run-off) to decrease the potential for injury on foliage, fruit and flowers of sensitive plants.
- ☐ Apply promptly after mixing with other pesticides to avoid or reduce alkaline hydrolysis of certain pesticides.
- ☐ Avoid applications when wind conditions impair thorough coverage.
- $\hfill \square$ Avoid applications on new transplants and unrooted cuttings.

2(ee):19-09-02

M-Pede® is a registered trademark of Gowan Company,L.L.C.



Produced For: Gowan Company P.O. Box 5569 Yuma, Arizona 85366-5569