Granular Fly Bait

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)	
methyl]-N nitro-2-imidazolidinimine	0.50%
Z-9-Tricosene	0.10%
OTHER INGREDIENTS:	

TOTAL:

100.00%

EPA Reg. No. 101563-94 EPA Est. No. 7319-NLD-001

STOP - READ THE LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. CAUTION

See Panel for First Aid Instructions and Complete Precautionary Statements and Directions for Use.

D00000958 79534728F 221216AV1 Net Weight 5 Lbs (2.268 kg)



	FIRST AID
lf swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
lf on skin:	 Take off contaminated clothing. Rinse skin immediately with plotty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
lf in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contac lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
Have the pro- treatment. Th	duct container or label with you when calling a poison control center or doctor or going for e Environmental Science U.S., LLC Emergency Response Telephone No. is 1-800-424-9300.
NOTE TO PHY	/SICIAN: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates.

This product a might basic equation metabalace. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.



Flies

PRODUCT FACTS • Fast acting • Ready-to-use scatter bait or paint-on application • Contains a specially formulated mixture of attractants • Controls flies resistant to organophosphates and carbamates	
MAXFORCE [®] Granular Fly Bait	 Ready-to-use bait containing 0.5% imidacloprid. Contains the attractant muscalure. Contains Bitrex® bittering agent to help prevent ingestion by animals, children, and pets.
CONTROLS	Flies
WHERE TO APPLY	 Around the outside of structures or partially enclosed or protected areas. In and around agricultural production facilities. Around the outside of commercial facilities.
MIXING INFORMATION	 Ready-to-use scatter bait. Paint-on application: Mix the fly bait with water.
HOW TO APPLY	 Scatter application. Place in any fly bait station. Paint-on application.
COVERAGE	14,000 Square feet (5.7 to 6.3 oz/1,000 sq ft)
RE-APPLY	Re-apply every 7 days as needed.
QUESTIONS ??	For questions or comments, call toll-free 1-800-331-2867.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLY MANAGEMENT

APPLY	 For use only around the outside of commercial facilities like canneries, dairy, meat and poultry processing plants, beverage plants, food processing plants, restaurants, bakeries, supermarkets, warehouses, dumpsters, grease pits, recycling areas, loading docks. For use only around the outside of agricultural production facilities like broiler houses, feedlots, livestock housing structures, and horse stables. For use only around the outside of and in walkways of caged layer houses. Place bait in areas inaccessible to animals. In horse stables, apply bait only in fly bait stations. Place bait stations in areas inaccessible to animals.
-------	---

PRODUCED FOR Environmental Science U.S., LLC 5000 CentreGreen Way. Suite 400 Cary, NC 27513 Product of The Netherlands 79538448F 221216AV1

FLY MANAGEMENT		
USE RATE	 5.7 to 6.3 oz/1,000 sq ft (scatter bait and bait station) 	
HOW TO APPLY	Apply the bait in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. Time applications to begin at the start of the season before fly populations have reached their peak. • Scatter application: Scatter the bait directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles. • Bait stations: Place in any fly bait station. Use 1 bait station to cover approximately 250 sq ft, and add up to 1.6 oz of bait per station. Secure bait stations at least 4 feet above the ground. Bait stations should be inaccessible to food-producing animals, children, and pets. • Paint-on application: Mix 1.5 oz bait with 1 fl oz warm water (3 lbs bait in 1 quart water) and stir thoroughly. Let stand for about 15 minutes until a paste consistency suitable for painting has formed. Apply the paste with a brush to surfaces where flies rest avoiding surfaces that are dusty.	
	 Do not place bait in areas accessible to food producing animals, children, and pets. Do not allow bait to come into contact with food or water. Do not apply to milking equipment. Do not apply inside food processing plants, restaurants or other commercial facilities. 	
MANAGEMEN	r tips	
Management p • Use proper s • Use along wi	JIts: use the bait as part of an overall Integrated Pest program. antifation and manure management. th dilutable sprays like TEMPO®. pht management.	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Storage

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling

USE RA

Completely empty container. Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Offer for reconditioning, if appropriate. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times

Dispose of container by offering for recycling or reconditioning, or at an approved waste disposal facility, or by incineration if allowed by state and local authorities. Do not use container in connection with food, feed, or drinking water. Snills

If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away. You may contact the Environmental Science U.S., LLC Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Environmental Science U.S., LLC Emergency Response Telephone No. is 1-800-424-9300.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability. CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Environmental Science U.S., LLC. All such risks shall be assumed by the user or buyer. DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. ENVIRONMENTAL SCIENCE U.S., LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Environmental Science U.S., LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES. INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT. NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID. OR AT ENVIRONMENTAL SCIENCE U.S., LLC'S ELECTION. THE REPLACEMENT OF PRODUCT.

Envu and the Envu logo are trademarks and MAXFORCE® and Tempo® are registered trademarks owned by Environmental Science U.S., LLC or one of its affiliates