

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
If in Eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If Swallowed:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If Inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
You may contact 1-800-498-5743 for emergency medical treatment information.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	

Parvosol II RTU Disinfectant is designed specifically as a general non-acid, ready-to-use, non-abrasive cleaner and

disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, restaurants, commercial kitchens, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories, and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross-contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food. In addition, Parvosol II RTU Disinfectant deodorizes those areas that generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

Parvosol II RTU Disinfectant contains no phosphates or phosphorous.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.

Fill or refill the 32 oz. product spray bottle with this Parvosol II RTU Disinfectant refill. Pour product into spray bottle over sink or tub basin.

DISINFECTION, DEODORIZING, AND CLEANING: Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for **HIV-1 1 minute** and for HBV and HCV **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop, or sponge. The product will not leave grit or soap scum.

Disinfection of animal quarters and kennels: Before using Parvosol II RTU Disinfectant, remove all animals and feed from premises. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Apply Parvosol II RTU Disinfectant for disinfection of pre-cleaned animal quarters and kennels. Saturate the surfaces with Parvosol II RTU Disinfectant for a period of **3 minutes** (for Bovine Viral Diarrhea Virus (BVD) **5 minutes**) (for Feline Calicivirus and Rabies virus **30 seconds**) (for Canine Parvovirus **10 minutes**). Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a **3-minute** contact time, unless otherwise noted, Parvosol II RTU Disinfectant kills the following organisms on hard, non-porous inanimate surfaces:

BACTERICIDAL ACTIVITY (BACTERIA): (Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, Parvosol II RTU kills the following organisms on hard non-porous inanimate surfaces:)
Pseudomonas aeruginosa, (ATCC 15442)
Listeria monocytogenes, (ATCC 35152)
Staphylococcus aureus, (ATCC 6538)
Yersinia enterocolitica, (ATCC 23715)
Salmonella (choleraesuis) enterica, (ATCC 10708)
Enterococcus faecium, (ATCC 6569)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 43895)
Corynebacterium ammoniagenes, (ATCC 6871)
Salmonella (typhi) enterica, (ATCC 6539)
Streptococcus pyogenes, (Necrotizing Fasciitis-Group A) (V.A. Medical Center Isolate, 04001)

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: (Using approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence food serum and a **3 minute** contact time, unless otherwise noted, Parvosol II RTU kills the following organisms on hard non-porous inanimate surfaces:)
Methicillin resistant *Staphylococcus aureus* (MRSA), (ATCC 33593)
Vancomycin resistant *Enterococcus faecalis* (VRE), (ATCC 51575)
Vancomycin intermediate resistant *Staphylococcus aureus* (VISA), (CDC Isolate 99287)
Methicillin resistant *Staphylococcus epidermidis* (MRSE) (ATCC 51625)

(continued)

PARVOSOL II RTU DISINFECTANT

CLEANER • DEODORIZER • VIRUCIDE

Fungicide (against pathogenic fungi) • Mildewstat (on hard inanimate surfaces)
Kills Pandemic 2009 H1N1 Influenza A Virus (formerly called swine flu)

Active Ingredients:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	0.105%
Inert Ingredients:	99.790%
Total.....	100.000%

EPA Reg. No. 1839-83-61282

EPA Est. 61282-WI-01

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for additional Precautionary Statements.



(Directions for Use, continued)

Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 123) Genotype USA400
Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), (NRS 384) Genotype USA300

TUBERCULOCIDAL ACTIVITY: Parvosol II RTU Disinfectant exhibits disinfectant efficacy against *Mycobacterium tuberculosis* BCG (*Mycobacterium bovis*) at 20°C when the treated surface is allowed to remain wet for **5 minutes**.

***VIRUCIDAL ACTIVITY:** Parvosol II RTU Disinfectant kills on hard, non-porous, inanimate surfaces when allowed to remain wet for a **10-minute** contact time against: Canine Parvovirus (ATCC VR-2017)
Hepatitis A Virus (HAV) (University of Ottawa)
Poliovirus Type 1 (ATCC VR-1000)

Parvosol II RTU Disinfectant, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Human Coronavirus** (ATCC VR-740), **SARS associated Coronavirus** (ZeptoMetrix), **Avian Influenza A strain H3N2** (ATCC VR-2072) and **Avian Influenza A strain H9N2** (Charles River Laboratories) (Avian Influenza strains H3N2 and H9N2) for **2 minutes**; **Paramyxovirus** (Mumps) (ATCC VR-1438), **Rhinovirus type 39** (ATCC VR-340) and Rotavirus (University of Ottawa) for **3 minutes**; **HIV-1 (associated with AIDS)** (Advanced Biotechnologies, Inc.) for **1 minute**; **Hepatitis B Virus (HBV)** (Hepadnavirus Testing, Inc.), **Hepatitis C Virus (HCV)** and **Bovine Viral Diarrhea Virus (BVDV)** (NVSL) when the treated surface is allowed to remain wet for **1 (or) (5) minute(s)**; and **Norovirus (Norwalk virus)**, **Feline Calicivirus** (University of Ottawa) and **Rabies Virus** (CDC) for **30 seconds**.

KILLS HIV-1, HBV, AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

Item No. 190416



Net Contents: 1 gal. (3.785 L)

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE: Store in a dry place no lower in temperature than 50°F or higher than 120°F.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill container. Clean container promptly after emptying.

Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.

Container disposal for Refill Spray Bottle: Do not reuse empty spray bottle except with refill bottle, otherwise discard in trash. Do not reuse empty refill bottle. Wrap refill bottle and discard in trash. Offer for recycling or reconditioning, if available.

For non-emergency (e.g., current product information) call: 1-800-621-8829

You may contact 1-800-498-5743 for emergency medical treatment information

Manufactured for:

HACCO, Inc.
110 Hopkins Drive
Randolph, WI 53956 U.S.A.

Parvosol® is a registered trademark of Neogen® Corporation.

HACCO, Inc. is a wholly-owned subsidiary of Neogen® Corporation LA220268-0420