

# PURE HARD SURFACE

## TECHNICAL REPORT



---

## PURE HARD SURFACE: TECHNICAL REPORT

---

### PRODUCT DESCRIPTION:

PURE Hard Surface is a colorless, odorless, ready-to-use disinfectant and sanitizer for use on hard non-porous, environmental surfaces, including food contact surfaces.

---

### INGREDIENTS:

|                            |             |           |
|----------------------------|-------------|-----------|
| <b>Active ingredients:</b> | Silver Ion† | 0.003 %   |
|                            | Citric Acid | 4.846 %   |
| <b>Other ingredients:</b>  |             | 95.151 %  |
| <b>TOTAL:</b>              |             | 100.000 % |

† Electrolytically generated silver ions stabilized in citric acid as Silver Dihydrogen Citrate

---

### REGISTRATION:

PURE Hard Surface is registered with the U.S. Environmental Protection Agency.

EPA REG. NO.: 72977-5-73912

U.S. Patent(s) 6,197,814; 6,583,176

Other patents pending

---

### NSF International NonFood Compounds:

PURE Hard Surface has been listed on the NSF International Nonfood Compounds list (D2, 144518). Products eligible for NSF Registration include all compounds used in and around food establishments (nonfood compounds), such as disinfectants and lubricants, and those used for pre-processing of food proprietary substances, such as fruit/vegetable washing agents. NSF Registration assures inspection officials and end users that formulations and labels meet appropriate food safety regulations. NSF Registration is based on the NSF Registration Guidelines (formerly USDA Guidelines for Obtaining Authorization of Compounds to be Used in Meat and Poultry Plants).



### Kosher Certified, Pareve:

PURE Hard Surface has been certified kosher by KSA, Kosher Supervision of America.



# PURE HARD SURFACE: TECHNICAL REPORT

## DIRECTION FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### DISINFECTION:

#### AREA OF APPLICATION:

Homes, offices, hospitals, restaurants, schools, hotels, restrooms, recreational facilities, public transport vehicles.

Use on painted, glazed tile, plastic, non-porous vinyl, polyurethane, plasticized PVC, EDPM, neoprene, Viton®, Teflon®, silicone, metal, glass, acrylic, sealed fiberglass, linoleum and glazed porcelain.

For other areas, test in an inconspicuous area before use.

#### TO DISINFECT HARD, NON-POROUS SURFACES:

Pre-clean surfaces prior to disinfecting.

Apply PURE Hard Surface to the surface until visibly wet for the contact time as listed. Wipe dry with a clean towel.

Refer to the product label for full use instructions.

| ORGANISM   | KILL TIME  |
|--|------------|
| Adenovirus Type 2<br>Avian Influenza A<br>HIV type 1<br>Human Coronavirus<br>Influenza A (H1N1)<br>Influenza A<br>Pseudomonas aeruginosa<br>Respiratory Syncytial Virus<br>Rotavirus<br><i>Salmonella enterica</i><br>SARS-CoV-2 (COVID-19 virus)<br>Swine Influenza A (H1N1)  | 30 SECONDS |
| Hepatitis B Virus (HBV)<br>Hepatitis C Virus (HCV)<br>Herpes Simplex Type 1<br>Murine Norovirus<br>Norovirus<br>Polio Type 2<br>Rhinovirus   | 60 SECONDS |
| <i>Acinetobacter baumannii</i><br><i>Campylobacter jejuni</i><br>Carbapenem resistant <i>Escherichia coli</i><br>Carbapenem resistant <i>Klebsiella pneumoniae</i><br>Carbapenem resistant <i>Klebsiella pneumoniae</i> , NDM-1 +<br>Community Associated Methicillin resistant <i>Staphylococcus aureus</i> (CA-MRSA)<br>Community Associated Methicillin resistant <i>Staphylococcus aureus</i> (CA-MRSA-PVL)<br><i>Escherichia coli</i> O157:H7<br><i>Listeria monocytogenes</i><br>Methicillin resistant <i>Staphylococcus aureus</i> (MRSA)<br><i>Staphylococcus aureus</i><br>Vancomycin resistant <i>Enterococcus faecium</i> (VRE) | 2 MINUTES  |
| Trichophyton mentagrophytes (Athlete's Foot Fungus)  | 5 MINUTES  |

Kill times are based on the EPA accepted label dated May 25, 2021.

---

## **PURE HARD SURFACE: TECHNICAL REPORT**

---

### **SANITIZATION OF FOOD CONTACT SURFACES**

#### **AREAS OF APPLICATION:**

Restaurants, homes, food processing facilities, food storage areas, supermarkets, kitchens schools, hotels, and dining halls. Use on painted, glazed tile, plastic, metal, glass and glazed porcelain. For other areas, test in an inconspicuous area before use.

#### **TO SANITIZE FOOD CONTACT SURFACES:**

Pre-clean surfaces prior to using this product. Do not use this product on utensils, dishes or glassware.

#### **CONSUMER APPLICATIONS:**

Spray, pour or spread PURE Hard Surface on surface until visibly wet. Let stand for 60 seconds and wipe with a clean towel or allow to air dry. No rinsing is required. This product kills 99.999% of *Escherichia coli* and *Staphylococcus aureus*.

---

### **COMMERCIAL APPLICATIONS:**

To sanitize food processing equipment and other hard surfaces in food processing locations, dairies, restaurants and bars:

#### **CLEAN, RINSE SANITIZE:**

Prior to application, remove gross food particles and soil by pre-flush or pre-scrape and when necessary, pre-soak. Thoroughly wash objects to be sanitized with a good detergent or cleaner followed by a potable water rinse prior to applying sanitizer. No potable water rinse is allowed after application as a sanitizer.

Apply this product by spraying or by total immersion. Surfaces must remain wet for 60 seconds. If the surface cannot be washed and rinsed, clean thoroughly in an appropriate fashion prior to sanitizing.

This product is a ready to use sanitizer that eliminates 99.999% of the following bacteria in 60 seconds: *Escherichia coli*, *Staphylococcus aureus*.

#### **FEDERALLY INSPECTED FACILITIES:**

Prior to use in a federally inspected meat and poultry plants and dairies, food products and packaging materials must be removed from the room or carefully protected. A potable water rinse is not permitted following the use of this product as a sanitizer on previously cleaned hard, non-porous surfaces, provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

Apply product to pre-cleaned hard surfaces thoroughly wetting surfaces with a cloth, mop, sponge, sprayer or by immersion. Surfaces should remain wet for 1 minute followed by adequate draining and air drying.

This product is a ready to use sanitizer that eliminates 99.999% of the following bacteria in 60 seconds: *Escherichia coli*, *Staphylococcus aureus*.

#### **BEVERAGE DISPENSING EQUIPMENT SANITIZER DIRECTIONS:**

For sanitizing of bottling or pre-mixed dispensing equipment after cleaning thoroughly rinse equipment with a potable water rinse. Fill equipment with this product and allow it to remain in the equipment for at least 60 seconds. Sanitizing solution should be drained from the system. To insure the removal of flavors, it is suggested that during changeover between products the system should be cleaned, rinsed and flushed with the sanitizing solution for at least 1 minute. Drain thoroughly and allow to air dry before reuse. No potable water rinse is allowed.

#### **FOR SANITIZING IN FISHERIES, MILK, WINE, CITRUS, POTATO & ICE CREAM PROCESSING PLANTS:**

For use as a sanitizer on conveyor belts and equipment to reduce or eliminate odors in the processing area. Also for use on filling equipment to reduce bacteria. Follow directions for sanitizing food contact surfaces.

## PURE HARD SURFACE: TECHNICAL REPORT

### MECHANISM OF ACTION

#### BACTERICIDAL/FUNGICIDAL ACTION:

The active ingredient in PURE Hard Surface is Silver Dihydrogen Citrate (SDC), a worldwide patented technology. SDC provides silver ions stabilized in citric acid.

The bacterial outer membrane is called the cell wall. Bacterial cell walls are made of peptidoglycan which provides protection and rigidity to the organism. The exact membrane constitution depends on the type of bacteria. SDC utilizes a multiple prong attack against microorganisms. SDC targets an organism's cell wall. Silver ions are highly attracted to sulfur-containing thiol groups found in metabolic and structural proteins bound to the membrane surface. SDC targets these critical proteins and destroys their structure. This disruption of the organism's membrane function and integrity lyses the membrane and the organism dies.

Unlike traditional antimicrobials, bacteria are actually attracted to SDC because they recognize citric acid as a food source. This "Trojan Horse" attack allows SDC to easily enter the microorganism through membrane transport proteins. Once inside the organism, SDC binds to DNA and intracellular proteins causing irreversible damage to the DNA and protein structure. Metabolic and reproductive functions halt, and the organism dies.

#### VIRUCIDAL ACTION:

Viruses are much smaller than bacterial and fungal cells and do not have metabolic activity. Viruses present fewer targets sites on which a biocide can act. Silver targets the viral envelope or capsid and the viral nucleic acid. Silver not only destroys the viral envelope or capsid, preventing the virus from attaching to a host cell, it also destroys the infectious component of the virus, the nucleic acid.

### MICROBIOLOGICAL TEST DATA

#### FOOD CONTACT SURFACE SANITIZER TEST METHODS:

Sanitizers applied to food contact surfaces are defined as incidental food additives under the Federal Food, Drug, and Cosmetic Act, as amended (21 U.S.C. 201 et seq.), and require establishment of a food additive tolerance or an exemption for the need for a tolerance. In addition, to support registration with the US EPA, the product must show efficacy following the AOAC Germicidal and Detergent Sanitizing Action of Disinfectants (AOAC 960.09) test method.

In this method, cultures of each test organism are grown in an appropriate broth and grown to an acceptable concentration. In duplicate for each test organism, 1 mL of the concentrated culture is inoculated into a flask containing 99 mL of the test product for the contact time of 30 seconds. After the contact time has elapsed, aliquots of the organism/test substance are neutralized and plated to analyze for surviving microorganisms as compared to the untreated control. Acceptable results must demonstrate a 99.999% reduction in the number of microorganisms within 30 seconds. The EPA requires that labels directions state a minimum contact time of 1 minute for sanitization of food contact surfaces.

| ORGANISM                     | DRIED VIRUS CONTROL | LOG REDUCTION | CONTACT TIME | % REDUCTION |
|------------------------------|---------------------|---------------|--------------|-------------|
| <i>Staphylococcus aureus</i> | $8.0 \times 10^7$   | $\geq 10^7$   | 30 seconds   | >99.999     |
| <i>Escherichia coli</i>      | $9.4 \times 10^7$   | $\geq 10^7$   | 30 seconds   | >99.999     |

Note: No surviving organisms were detected for all three lots in all replicates for both test organisms.

#### FUNGAL TEST METHODS:

For registration with the US EPA, efficacy against pathogenic fungi is determined by following the AOAC Fungicidal Test Method or modifications of either the AOAC Use Dilution Test Method modified or the AOAC Germicidal Spray Products Test Method which meet the criteria of the AOAC Fungicidal Test Method. The carrier method is outlined below:

The disinfectant is placed in a water bath and allowed to equilibrate to a temperature of  $20.0 \text{ }^\circ\text{C} \pm 0.5^\circ\text{C}$ . Carriers will be inoculated with test culture. Carriers must have a minimum concentration of  $10^4$  after drying. Each contaminated and dried carrier is placed into a test tube containing test substance for the specified contact time and then transferred to test tubes containing growth medium and a neutralizing agent to stop the action of the disinfectant. The carriers are incubated for an appropriate time based upon the test organism. The tubes are examined for growth or no growth. To pass a 60 carrier test, two batches are tested and 59 out of 60 carriers must show no growth for each product batch. To pass a 10 carrier test, two batches are tested and all 10 carriers must show no growth.

| ORGANISM  | # OF CARRIERS EXPOSED | # OF CARRIERS SHOWING GROWTH | CONTACT TIME |
|---|-----------------------|------------------------------|--------------|
| Trichophyton mentagrophytes (Athlete's Foot Fungus) ATCC#9533 | 10                    | 0                            | 5 minutes    |

## PURE HARD SURFACE: TECHNICAL REPORT

### MICROBIOLOGICAL TEST DATA (CONT.)

#### BACTERIAL TEST METHODS:

For registration with the US Environmental Protection Agency (US EPA), disinfection efficacy is tested following specific test methods. In each method, cultures of bacteria are dried onto a number of small carriers (stainless steel penicylinders in the Use-Dilution test or glass slides in the AOAC Germicidal Spray Products Test). Once dried, the carriers must contain a bacteria concentration of at least  $10^4$ . These carriers are exposed to the disinfectant for a specified contact time and then transferred to test tubes containing growth medium and a neutralizing agent to stop the action of the disinfectant. The carriers are incubated for 48 hours. The tubes are then examined for growth or no growth. To pass a 60 carrier test, three batches are tested and 59 out of 60 carriers must show no growth for each product batch. To pass a 10 carrier test, two batches are tested and all 10 carriers must show no growth.

To make general broad spectrum claims a disinfectant must show efficacy against a *Staphylococcus aureus* (Gram positive bacteria) and *Salmonella enterica* (Gram negative bacteria). To make claim for use in and hospital/medical environments a disinfectant must show efficacy against *Pseudomonas aeruginosa* (nosocomial bacteria). Other bacteria may be tested to obtain additional claims.

| ORGANISM  | # OF CARRIERS EXPOSED | # OF CARRIERS SHOWING GROWTH | CONTACT TIME | CARRIER POPULATION |
|---|-----------------------|------------------------------|--------------|--------------------|
| <i>Pseudomonas aeruginosa</i> (ATCC#15442)                              | 180                   | 0                            | 30 seconds   | $10^6$             |
| <i>Staphylococcus aureus</i> (ATCC#6538)                                | 180                   | 2                            | 2 minutes    | $10^6$             |
| <i>Salmonella enterica</i> (ATCC#10708)                                 | 180                   | 0                            | 30 seconds   | $10^5$             |
| <i>Listeria monocytogenes</i> (ATCC#19111)                              | 20                    | 0                            | 2 minutes    | $10^6$             |
| Vancomycin resistant <i>Enterococcus faecium</i> (VRE) (ATCC#700221)    | 20                    | 0                            | 2 minutes    | $10^{4.5}$         |
| Methicillin resistant <i>Staphylococcus aureus</i> (MRSA) (ATCC#700698) | 20                    | 0                            | 2 minutes    | $10^5$             |
| Community Associated MRSA (CA MRSA) (NRS123, USA 400)                   | 20                    | 0                            | 2 minutes    | $10^6$             |
| Community Associated MRSA (CA MRSA-PVL) (NRS 192)                       | 20                    | 0                            | 2 minutes    | $10^6$             |
| <i>Escherichia coli</i> O157:H7 (ATCC#43888)                            | 20                    | 0                            | 2 minutes    | $10^{4.5}$         |
| <i>Campylobacter jejuni</i> (ATCC#29428)                                | 20                    | 0                            | 2 minutes    | $10^6$             |
| <i>Acinetobacter baumannii</i> (ATCC#19606)                             | 20                    | 0                            | 2 minutes    | $10^6$             |
| Carbapenem resistant <i>Escherichia coli</i>                            | 20                    | 0                            | 2 minutes    | $10^5$             |
| Carbapenem resistant <i>Klebsiella pneumoniae</i>                       | 20                    | 0                            | 2 minutes    | $10^5$             |
| Carbapenem resistant <i>Klebsiella pneumoniae</i> , NDM1+               | 20                    | 0                            | 2 minutes    | $10^5$             |

## PURE HARD SURFACE: TECHNICAL REPORT

### MICROBIOLOGICAL TEST DATA (CONT.)

#### VIRAL TEST METHODS:

The US EPA accepts carrier based virucidal test methods to support virucidal activity of a disinfectant which are modifications of the AOAC Use Dilution Test or the AOAC Germicidal Spray Products Test. Each virus claimed must be tested in an appropriate test system using a cell line which supports the growth of the virus. The method as outlined in the EPA Disinfectant Technical Science section (DIS-TSS 07) states:

To simulate in-use conditions, the specific virus to be treated must be inoculated onto hard surfaces, allowed to dry, and then treated with the product according to the directions for use on the product label. One surface for each of two different batches (or more when required) of disinfectant must be tested against a recoverable virus titer of at least  $10^4$  from the test surface (petri dish, glass slide, steel cylinder, etc.) for a specified exposure period at room temperature. The virus is then assayed by an appropriate virological technique.

In order for the data to be considered valid, the following criteria must be met:

- 1: Virus concentration after drying must be at least  $10^4$ .
- 2: Complete inactivation of the viruses at all dilutions is required. If cytotoxicity is evident, at least a 3 log reduction of virus concentration must be demonstrated beyond the cytopathic effect.
- 3: Cell controls must be negative for infectivity.

| ORGANISM   | DRIED VIRUS CONTROL | LOG REDUCTION | CONTACT TIME |
|--|---------------------|---------------|--------------|
| HIV type 1- Strain HTLV IIIB                           | $10^{5.25}$         | $\geq 3.75$   | 30 seconds   |
| Rotavirus (Strain WA, Ottawa)                          | $10^{4.5}$          | $\geq 4.0$    | 30 seconds   |
| Human Coronavirus (ATCC VR-740)                        | $10^{4.5}$          | $\geq 4.0$    | 30 seconds   |
| Influenza A (H1N1) (ATCC VR-1469)                      | $10^{6.5}$          | $\geq 6.0$    | 30 seconds   |
| Swine Influenza A (H1N1) (ATCC VR-333)                 | $10^{6.75}$         | $\geq 6.25$   | 30 seconds   |
| Respiratory Syncytial Virus (ATCC VR-26)               | $10^{4.75}$         | $\geq 4.25$   | 30 seconds   |
| Adenovirus Type 2 (ATCC VR-846)                        | $10^{6.0}$          | $\geq 5.5$    | 30 seconds   |
| SARS CoV2 (COVID 19 virus) (USA-WA1/2020)              | $10^{5.25}$         | $\geq 4.75$   | 30 seconds   |
| Herpes Simplex Type 1 VR-733 F(1) Strain (ATCC VR-733) | $10^{6.0}$          | $\geq 5.5$    | 60 seconds   |
| Murine Norovirus (MNV-1.CW1)                           | $10^{6.5}$          | $\geq 6.0$    | 60 seconds   |
| Norovirus -as Feline Calicivirus (ATCC VR-782)         | $10^{6.0}$          | $\geq 5.88$   | 60 seconds   |
| Avian Influenza A (ATCC VR-2072)                       | $10^{4.75}$         | $\geq 4.25$   | 30 seconds   |
| Influenza A (ATCC VR-544)                              | $10^{6.5}$          | $\geq 6.0$    | 30 seconds   |
| Rhinovirus (ATCC VR-1147)                              | $10^{4.5}$          | $\geq 4.0$    | 60 seconds   |
| Polio Type 2 (ATCC VR-1002)                            | $10^{4.5}$          | $\geq 3.0$    | 60 seconds   |
| Hepatitis B Virus (HBV)                                | $10^{6.0}$          | $\geq 5.79$   | 60 seconds   |
| Hepatitis C Virus (HCV)                                | $10^{5.25}$         | $\geq 4.93$   | 60 seconds   |

All efficacy studies performed on the product to substantiate the efficacy claims are performed following Good Laboratory Practices at third party laboratories which are recognized for their expertise in antimicrobial test methods. The methods described above were the current accepted methods and criteria at the time of testing.



## PURE HARD SURFACE: TECHNICAL REPORT

### SAFETY DATA:

#### FROM THE EPA LABEL REVIEW MANUAL:

The EPA determines the Toxicity Category based on the Acute Toxicity Data provided by the registrant. Here is the rating system utilized by the EPA:

| STUDY                         | CATEGORY I  | CATEGORY II   | CATEGORY III   | CATEGORY IV  |
|-------------------------------|---|---|--|--|
| Acute Oral                    | Up to and including 50 mg/kg  | > 50 thru 500 mg/kg   | > 500 thru 5000 mg/kg  | > 5000 mg/kg   |
| Acute Dermal                  | Up to and including 200 mg/kg   | > 200 thru 2000 mg/kg   | > 2000 thru 5000 mg/kg   | > 5000 mg/kg   |
| Acute Inhalation <sup>1</sup> | Up to and including 0.05 mg/liter   | > 0.05 thru 0.5 mg/liter  | > 0.5 thru 2 mg/liter  | > 2 mg/liter   |
| Primary Eye Irritation        | Corrosive (irreversible destruction of ocular tissue) or corneal involvement or irritation persisting for more than 21 days | Corneal involvement or other eye irritation clearing in 8-21 days | Corneal involvement or other eye irritation clearing in 7 days or less | Minimal effects clearing in less than 24 hours                           |
| Primary Skin Irritation       | Corrosive (tissue destruction into the dermis and/or scarring)  | Severe irritation at 72 hours (severe erythema or edema)          | Moderate irritation at 72 hours (moderate erythema)                    | Mild or slight irritation at 72 hours (no irritation or slight erythema) |
| <b>SIGNAL WORD</b>            | <b>DANGER</b>   | <b>WARNING</b>  | <b>CAUTION</b>   | <b>NONE REQUIRED</b>   |

<sup>1</sup> 4 hr exposure

A Signal Word is required based on the Toxicity category assigned to a pesticide by the EPA. For Category IV products, no Signal Word is required and no first aid statements are required. The Signal Word is determined by the most severe toxicity category assigned to the five acute toxicity studies or by the presence of methanol in concentrations of 4% or more.

**PURE Hard Surface falls into US EPA Category IV based on all of the toxicity data.  
The product does not require a Signal word or First Aid statements to be listed on the label.**


#### HMIS RATING:

Hazardous Material  
Identification System

|              |   |
|--------------|---|
| Health       | 0 |
| Flammability | 0 |
| Reactivity   | 0 |

#### NFPA:

National Fire Protection  
Association





---

## PURE HARD SURFACE: TECHNICAL REPORT

---

### PHYSICAL DATA:

|                                       |                          |
|---------------------------------------|--------------------------|
| Appearance                            | Clear, colorless liquid  |
| Odor                                  | Practically odorless     |
| pH                                    | 2                        |
| Specific Gravity (H <sub>2</sub> O=1) | 1                        |
| Solubility                            | Water soluble            |
| VOC Content (% Wt.)                   | 0.00% (0.000 lbs/gallon) |
| Flash Point                           | > 212°F                  |

---

### STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal.

*PESTICIDE STORAGE:* Store in a cool, dry area away from direct sunlight at temperatures above freezing.

*PESTICIDE DISPOSAL:* To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

*CONTAINER HANDLING:* Nonrefillable\* Container. Do not reuse or refill except as described in the directions for use. Refill only with this product. If empty: Place in trash or offer for recycling if available.

*\*As defined by the EPA, a nonrefillable container is one that is not intended to be refilled with pesticide for sale. This term does not apply to bottles which are intended to be refilled with the same product more than once for use but not for sale or distribution.*

---

### QUESTIONS?

For Customer Service please call: (619) 596-8600

Customer Service Hours of Operation Mon-Fri 8am-5pm PST

Distributed by PURE Bioscience, Inc.

---

### OTCQB: PURE



VOICE: [619.596.8600](tel:619.596.8600) | FACSIMILE: [619.596.8790](tel:619.596.8790)  
771 JAMACHA ROAD, #512 | EL CAJON, CALIFORNIA 92020  
OTCQB: [PURE](https://www.purebio.com)

©2021 PURE Bioscience, Inc. [PUREBIO.COM](https://www.purebio.com)

