

QUALI-PRO[®]

GROUP 4 INSECTICIDE

IMIDACLOPRID 2F

Turf & Ornamental Insecticide



Turf and Ornamental and Nursery and Greenhouse Uses

- Foliar and Systemic Insect Control for use on Turfgrass including Sod Farms, Landscape Ornamentals, Fruit and Nut Trees and Interior Plantscapes
- Foliar and Systemic Insect Control for use in Greenhouses, and Nurseries on Ornamentals, Fruit and Nut Trees and Vegetable Plants

ACTIVE INGREDIENT: % BY WT.

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]

-N-nitro-2-imidazolidinimine 21.8%

OTHER INGREDIENTS: 78.2%

TOTAL: 100.0%

Contains 2 pounds of imidacloprid per gallon.

Flowable insecticide.

Shake well before using.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

EPA Reg. No. 53883-514
EPA Est No. 53883-TX-002

CONTENTS:
1 GALLON



Manufactured for:
Control Solutions, Inc.
5903 Genoa Red Bluff Road
Pasadena, TX 77507

EPA 120313/Rev A

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of soap and water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: In case of emergency, contact SafetyCall at 1-866-897-8050. Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
NOTE TO PHYSICIAN: No specific antidote is available. Treat patient symptomatically.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash clothing before reuse.

Keep children or pets away from treated area until dry.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Applicators and other handlers must wear:

Worker Protection Standard Uses:

Applicators and other handlers (mixers and loaders) who handle this product for uses covered by the Worker Protection Standard (40 CFR Part 170) – such as sod farms, must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.
- Shoes plus socks

Non- Worker Protection Standard Uses:

Applicators and other handlers must wear:

- Shirt and pants
- Gloves
- Shoes plus socks

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(4)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. Wash thoroughly and change into clean clothing. Wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Apply this product only as specified on this label. Extreme care must be taken to avoid runoff. Apply only to soil or other fill substrate that will accept the solution at the specified rate. Do not treat soil that is water-saturated or frozen or in any conditions where run-off or movement from the treatment area (site) is likely to occur.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- o Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- o Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- o Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- o Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See individual sites for specific pollinator protection application restrictions. If none exist under the specific site, for foliar applications, follow these application directions for food/feed crops and commercially grown ornamentals that are attractive to pollinators, and for non-agricultural uses:



FOR FOOD/FEE CROPS AND COMMERCIALLY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.



Non-Agricultural Products:

Do not apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE while bees are foraging. Do not apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to plants that are flowering. Only apply after all flower petals have fallen off.

Read entire label before using this product. This label must be in the possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

STATE SPECIFIC RESTRICTIONS

The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to be agricultural uses.

APPLICATION ON TURFGRASS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE may be used to control listed insect pests on turfgrass in residential lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms.

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE controls listed soil inhabiting pests such as Northern & Southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs *Spherophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass ateniens, *Ataenius spretulus* and *Aphodius* spp.; European Crane Fly, *Tipula paludosa*; and mole crickets, *scapteriscus* spp. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE can also be used for suppression of cutworms and chinch bugs.

For optimum control, make applications preceding or during the egg laying period of the target pest. The active ingredient in QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE has enough residual activity so that applications can be made preceding the egg laying activity. Application timing can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Most favorable control will be achieved when applications are made prior to egg hatch of the target pests. Follow applications with sufficient irrigation or rainfall to move the active ingredient through the thatch.

RESTRICTIONS: Do not make applications when turfgrass areas are waterlogged or the soil is saturated with water. Sufficient distribution of the active ingredient cannot be achieved under these conditions. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

Do not exceed a total of 1.6 pt (0.4 lb of active ingredient)/A per year.

APPLICATION EQUIPMENT FOR USE ON TURFGRASS

Apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

RESTRICTION: Do not apply through any irrigation system.

APPLICATIONS

TURF GRASSES

PEST	RATE	APPLICATION INSTRUCTIONS
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbug Black turfgrass atanius Cutworms (suppression) European chafer European crane fly Green June Beetle Japanese beetle Northern Masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	1.25 to 1.6 pt/A or 0.46 to 0.6 fl. oz. (14 to 17 mL) per 1000 sq. ft.	For best control of grubs, billbugs, annual bluegrass weevil, and European Crane Fly, apply prior to egg hatch of the target pest. Read APPLICATION EQUIPMENT section of this label.
Chinch bugs (suppression) Mole crickets	1.6 pt /A or 0.6 fl. oz. (17 mL) per 1000 sq. ft.	For suppression of chinch bugs, apply before hatching of the first instar nymphs. To control mole crickets apply before or during the peak egg hatch period. Use a curative insecticide in addition to QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE when adults or large nymphs are present and actively tunneling. Follow label instructions for other insecticides when tank-mixing.
Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialists for more specific information regarding timing of application.		
Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. Do not mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.		
RESTRICTION: <ul style="list-style-type: none"> • Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year. • Do not apply this product in a way that will contact people or pets. • Do not allow children or pets to enter treated areas until sprays have dried. • Do not allow this product to contact plants in bloom if bees are foraging in the treatment area. 		

APPLICATION TO ORNAMENTALS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE is for use on ornamentals in commercial and residential landscapes and interior plantscapes. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE is a systemic product and will be taken up into the plant system from root uptake. The product must be placed where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is taken up throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, apply prior to anticipated pest infestation to achieve optimum levels of control.

RESTRICTIONS:

- For outdoor applications, do not exceed a total of 1.6 pt (0.4 lb of active ingredient) /A per year.
- Not for use on grass grown for seed or on commercial fruit and nut trees.
- Do not apply this product, by any application method, to linden, basswood, or other *Tilia* species.

Ant Management Programs

Use QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE applications can be then supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population.

APPLICATION EQUIPMENT FOR FOLIAR APPLICATIONS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE mixes readily with water and may be used in many types of application equipment. Mix product with the required amount of water and apply as desired dependent upon the selected use pattern.

When making foliar applications on hard to wet foliage such as holly, pine or ivy, the addition of a spreader/sticker is recommended. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed as would be used in a dilute application.

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

RESTRICTION:

- Do not apply through any irrigation system.
- Do not apply this product in a way that will contact people or pets.
- Do not allow children or pets to enter treated areas until sprays have dried.

APPLICATIONS

FOR USE ONLY IN AND AROUND THE PERIMETER OF INDUSTRIAL AND COMMERCIAL BUILDINGS AND RESIDENTIAL AREAS

CROP	PEST	RATE	APPLICATION INSTRUCTIONS
Trees Shrubs Evergreens Flowers Foliage plants Groundcovers Interior plantscapes	Adelgids Aphids Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Mealybugs Psyllids Sawfly Larvae Thrips (suppression) Whiteflies	1.5 fl. oz. (45 mL) per 100 gal of water	Foliar Applications: Begin applications before the onset of high pest populations and reapply as needed.
	White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle)	0.46 to 0.6 fl. oz. (14 to 17 mL) per 1000 sq. ft.	Broadcast Applications: Use enough water to mix the product and thoroughly apply to the treatment area. Do not use less than 2 gallons of water per 1000 sq ft. Irrigate after application to incorporate QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE into the upper soil layer. For additional use directions, refer to the FLOWERS and GROUND COVERS section of this label.

SOIL INJECTION* AND SOIL DRENCH APPLICATIONS IN AND AROUND THE PERIMETER OF INDUSTRIAL AND COMMERCIAL BUILDINGS AND RESIDENTIAL AREAS, AND STATE, NATIONAL AND PRIVATE WOODED AND FORESTED AREAS

*No Soil injection Applications Allowed in Nassau or Suffolk Counties of New York.

PEST	CROP/RATE	APPLICATION INSTRUCTIONS	REMARKS
<p>Adelgids Aphids Armored scales (suppression) Black vine weevil larvae Eucalyptus longhorned borer Flatheaded borer (including bronze, alder and emerald ash) Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine tip moth larvae Psyllids Roundheaded borers (including Asian longhorned beetles) Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies</p>	<p>TREES</p> <p>Use the following rates as a function of tree Diameter at Breast Height (D.B.H.):</p> <p>Apply 0.1 to 0.4 fl. oz. (3 to 12 mL) per inch of trunk diameter (D.B.H.)</p> <p>You may use the higher rate (0.3 – 0.4 fl. oz.) only for trees greater than 15 D.B.H. to control the following pests: Asian longhorned beetle, Emerald Ash Borer, Eucalyptus longhorned borer, Bronze birch borer, Alder borer</p> <p>RESTRICTION:</p> <p>Do not exceed a total of 1.6 pt (0.4 lb of active ingredient) /A per year.</p> <p>Diameter at Breast Height (D.B.H.) = is measured at 4.5 feet from the ground.</p>	<p>SOIL INJECTION:</p> <p>Grid System: Space holes in a grid pattern on 2.5 foot centers extending to the drip line of the tree.</p> <p>Circle System: Apply in holes evenly spaced in circles (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.</p> <p>Basal System: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p>Soil Drench: Apply uniformly as a drench around the base of the tree in not less than 10 gallons of water per 1000 square feet. Direct application to the root area. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>	<p>Use enough water to mix the product and inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment area. Keep the treated area moist for 7 to 10 days.</p> <p>Do not use less than 4 holes per tree.</p> <p>For Control of Specified Borers: Trees with existing insect damage and stress may not recover after treatment with QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE.</p>

(continued)

PEST	CROP/RATE	APPLICATION INSTRUCTIONS	REMARKS
	<p>SHRUBS</p> <p>0.1 to 0.2 fl. oz. (3 to 6 mL) per foot of shrub height</p>	<p>Soil Injection: Apply at the specified dosage to each plant.</p> <p>Soil Drench: Apply uniformly as a drench around the base of the tree in not less than 10 gallons of water per 1000 square feet. Direct application to the root area. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>	<p>Use enough water to mix the product and inject an equal amount of solution in each hole. Use low pressure and sufficient solution for distribution of the liquid into the treatment area. Keep the treated area moist for 7 to 10 days.</p> <p>Do not use less than 4 holes per shrub.</p>
	<p>FLOWERS AND GROUNDCOVERS</p> <p>0.46 to 0.6 fl. oz. (14 to 17 mL) per 1000 sq. ft.</p>	<p>Apply as a broadcast treatment before or after planting, prior to bloom or after all flower petals have fallen off for established plants. Mix into soil. On established plants, irrigate thoroughly after application.</p>	

FOLIAR APPLICATIONS FOR USE ONLY IN AND ON RESIDENTIAL AREAS

CROP	PEST	RATES	APPLICATION INSTRUCTIONS
<p>POME FRUITS Apple Crabapple Loquat Mayhaw Pear Pear (oriental) Quince</p>	<p>Aphids (except Wooly apple aphid) Leafhoppers (including glassy-winged sharpshooter) Leafminer Mealybugs¹ San Jose scale¹</p>	<p>1.5 fl. oz. (45 mL) per 100 gal or 6.0 fl. oz./A³</p>	<p>Apply specified dosage as foliar spray as needed after petal-fall is complete.</p> <p>For control of rosy apple aphid, apply prior to leafrolling caused by the pest.</p> <p>For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE will not control late stage larvae.</p> <p>For San Jose Scale, time applications to the crawler stage. Treat each generation.</p> <p>For late season (preharvest) control of leafhopper species, apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE while most leafhoppers are in the nymphal stage.</p> <p>For control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybug.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • Do not apply more than 6.0 fl. oz./A in a single application. • Do not make more than 5 applications /A per year. • Do not apply more than 1.6 pt (0.4 lb of active ingredient) per year. • Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.

(continued)

FOLIAR APPLICATIONS FOR USE ONLY IN AND ON RESIDENTIAL AREAS (continued)

CROP	PEST	RATES	APPLICATION INSTRUCTIONS
Pecan ²	Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	1.5 fl. oz. (45 mL) per 100 gal or 6.0 fl. oz./A ³	Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14 day interval may be required to achieve control. Scout and re-treat if needed. Thorough uniform coverage of foliage is necessary for optimal control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's use rate may improve coverage. RESTRICTIONS: <ul style="list-style-type: none"> Do not apply more than a total of 18.0 fl. oz. of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE /A per year. Do not make more than 3 applications. Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.
<p>¹ Not for use in California for control on pears. ² Use on pecans not permitted in California unless directed by state-specific 24(c) supplemental labeling. ³ The amount of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE required /A will depend on tree size and volume of foliage present. The rate /A is based on a standard of 400 gallons of dilute spray solution /A for large trees.</p>			

FOLIAR APPLICATION FOR USE ONLY IN AND ON INDUSTRIAL AND COMMERCIAL BUILDINGS AND RESIDENTIAL AREAS

CROP	PEST	RATE	APPLICATION INSTRUCTIONS
Grapes	Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1.5 fl. oz. (45 mL) per 100 gal or 3.0 fl. oz/A (90 mL/A)	Apply specified dosage as a foliar spray using 200 gallons of water /A. RESTRICTIONS: <ul style="list-style-type: none"> Do not apply more than a total of 6.0 ounces of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE /A per year. Allow at least 14 days between applications. Applications may be applied up to and including day of harvest. Pre-Harvest Interval 0 days.

Restrictions

- Keep children and pets off treated area until dry.
- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not allow runoff or puddling of irrigation water following application. Do not apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year.

APPLICATION IN GREENHOUSES, NURSERIES, ORNAMENTALS, FRUIT AND NUT TREES AND VEGETABLE PLANTS

APPLICATION TO ORNAMENTALS AND VEGETABLE PLANTS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE may be used to control listed insect pests on ornamental and vegetable plants in nurseries and greenhouses. Insect protection is achieved because QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE is a systemic product and the active ingredient moves upward into the plant system. Apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to the growing part of the plant for more absorption of the active ingredient. Nitrogen containing fertilizer may be added to the solution to aid in the uptake of the active ingredient where applicable. Apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE as a foliar spray or by soil applications such as soil injection, drenches, chemigation and broadcast sprays.

Soil applications to plants with woody stems will require applications of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE before expected pest infestations due to the delay in the uptake of the active ingredient and the time until the product is taken up throughout the plant.

Restriction: For outdoor applications, do not exceed a total of 1.6 pt (0.4 lb of active ingredient) /A per year.

Bark Media: The length of protection after treatment with QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE may be shortened if the media has 30% or more bark content.

APPLICATION EQUIPMENT FOR ORNAMENTALS AND VEGETABLE PLANTS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE mixes with water and may be applied with different types of application equipment. After mixing with the correct amount of water, follow the application directions for the selected use pattern.

For applications on hard to wet foliage such as holly, pine or ivy, the use of a spreader/ sticker is recommended. For application by concentrate or mist type spray equipment, use the same amount as would be used in a dilute application.

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE is compatible with frequently used fungicides, miticides, liquid fertilizers. Compatibility may be tested in a small jar by using the correct proportion of products if compatibility information is not available.

APPLICATION THROUGH IRRIGATION SYSTEMS

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE may be applied alone or as a tank mixture with other chemicals or pesticides registered for application through irrigation systems. The normal dilution ratio is 1:100 to 1:200 depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system.

Apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE only through micro irrigation (individual spaghetti tubes), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non uniform distribution of treated water.

If you have any questions about calibration, contact your State Extension Service specialist, equipment manufacturers or other experts in this area.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:

If the source of water for your irrigation system is a public water supply, follow the instructions below:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of material that is compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

APPLICATION TO GRASSY AREAS IN NURSERIES

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE may be used on nursery grass in areas such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries.

QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE controls listed soil inhabiting pests of grassy areas of nurseries, such as Northern and Southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotroqus majalis*; Green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; Billbugs, *Spherophorus* spp.; Annual bluegrass weevil, *Hyperodes* spp.; Black turfgrass atenius, *Ataenius spretulus* and *Aphodius* spp. and mole crickets, *Scapteriscus* spp. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE can also be used for suppression of cutworms and chinch bugs.

For optimum control, make applications preceding or during the egg laying period of the target pest. The active ingredient in QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE has enough residual activity so that applications can be made preceding the egg laying activity. Application timing can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Most favorable control will be achieved when applications are made prior to egg hatch of the target pests. Follow application with sufficient irrigation or rainfall to move the active ingredient through the thatch.

RESTRICTIONS

- Do not make applications when grassy areas are waterlogged or the soil is saturated with water. Sufficient distribution of the active ingredient cannot be achieved under these conditions. The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.
- Do not apply more than a total of 1.6 pt (0.4 lb of active ingredient) /A per year.

APPLICATION EQUIPMENT FOR USE ON GRASSY AREAS IN NURSERIES

Apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE in enough water to provide sufficient distribution in the treated area. Use accurately calibrated equipment typically used for the application of soil insecticides which will produce a uniform, course droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

APPLICATION SITES

GRASSY AREAS OF FIELD AND FOREST NURSERIES

PEST	RATES	APPLICATION INSTRUCTIONS
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass atenius <i>Phyllophaga</i> spp. Cutworms (suppression) European chafer Green June Beetle Japanese beetle Northern masked chafer Oriental beetle Southern masked chafer	19.2 to 25.6 fl. oz. /A or 0.45 to 0.6 fl. oz. (13 to 17 mL) per 1,000 sq. ft.	For best control of grubs, billbugs and annual bluegrass weevil, make application prior to egg hatch of the target pest. Make sure to read APPLICATION EQUIPMENT section of this label. For suppression of chinch bugs, make application prior to the hatching of the first instar nymphs. For control of mole crickets make application before or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE application should be accompanied by a curative insecticide. Follow label instructions for other insecticides when tank-mixing.
Chinch bugs (suppression) Mole crickets	25.6 fl. oz./A or (17 mL) per 1,000 sq. ft.	Consult your local turf, State Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application. Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. Do not mow grass area until after adequate irrigation or rainfall has occurred so that evenness of application will not be affected. RESTRICTION: <ul style="list-style-type: none"> · Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year. · Do not apply this product in a way that will contact people or pets. · Do not allow children or pets to enter treated areas until sprays have dried. · Do not allow this product to contact plants in bloom if bees are foraging in the treatment area.

ORNAMENTALS

FOLIAR AND SYSTEMIC APPLICATION IN OR ON FIELD-GROWN NURSERY AND CONTAINER STOCK, GREENHOUSE ORNAMENTALS, AND ORNAMENTALS GROWN IN FLAT BENCHES OR BEDS

PEST	CROP	RATES	APPLICATION INSTRUCTIONS
Adelgids Aphids Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips (suppression) Whiteflies	Trees (including non-bearing fruit and nut), Shrubs Evergreens Flowers Ground covers Vegetable plants*	1.7 fl. oz. (50 mL) per 100 gal. of water	<p>Foliar Applications: Start treatments before high pest pressure is observed and reapply as needed.</p> <p>For resistance management purposes, do not make a QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE foliar application following a soil application in the same crop.</p> <p>RESTRICTIONS</p> <ul style="list-style-type: none"> • For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato. • Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year. Minimum interval between applications: 5 days.
White grub larvae (such as Japanese beetle larvae, Chafer, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle)		0.45 to 0.6 fl. oz. (13 to 17 mL) per 1,000 sq. ft.	<p>Broadcast Applications: Mix required amount of product in enough water to uniformly and exactly cover the treatment area. Do not use less than 2 gallons of water per 1000 sq. ft. Irrigate to integrate QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE into the upper soil level.</p> <p>Refer to REMARKS section for use directions specific for FLOWERS AND GROUND COVERS concerning additional use directions.</p> <p>RESTRICTION: Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year. Minimum interval between applications: 5 days.</p>

* For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

SOIL INJECTION, SOIL DRENCH AND BROADCAST APPLICATIONS IN NURSERY AND GREENHOUSE

PEST	CROP/RATES	APPLICATION INSTRUCTIONS
Adelgids Aphids Armored scales Black vine weevil larvae Eucalyptus longhorned borers Flatheaded borers (including bronze birch and alder borers) Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Pine Tip moth larvae Psyllids Royal palm bugs Sawfly larvae Soft scales Thrips (suppression) White grub larvae Whiteflies	<p style="text-align: center;">TREES</p> Apply 0.1 to 0.2 fl oz (3 to 6 mL) per inch of trunk diameter. Diameter at Breast Height (D.B.H.) is measured at 4.5 feet from the ground.	<p>Soil Injections: Grid System: Space holes on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. Circle System: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. Basal System: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.</p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.</p> <p>Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p>For Control of Specified Borers: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p> <p>RESTRICTIONS</p> <ul style="list-style-type: none"> • No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York. • Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year.

(continued)

PEST	CROP/RATES	APPLICATION INSTRUCTIONS
	<p>SHRUBS 0.1 to 0.2 fl. oz. (13 to 17 mL) per foot of shrub height</p>	<p>Soil Injection: Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.</p> <p>Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p> <p>RESTRICTIONS</p> <ul style="list-style-type: none"> • No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York. • Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year.
	<p>FLOWERS AND GROUND COVERS 0.45 to 0.6 fl. oz. (13 to 17 mL) per 1,000 sq. ft.</p>	<p>Apply as a broadcast treatment before or after planting, prior to bloom or after all flower petals have fallen off for established plants. After application to established plants, irrigate thoroughly.</p> <p>RESTRICTION: Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year.</p>

EBB & FLOOD APPLICATION

Prior to treatment, to ensure accurate uptake by the plants, at least 10 plants must be brought up to a known field capacity and allowed to dry out for one or two days. Once dry, re-wet these plants to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. This should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

PEST	POT SIZE (inches)	Herbaceous species including vegetable plants* (1 or 2 plants per pot)	Woody perennials, Herbaceous species including vegetable plants* (3 or more plants per pot)	APPLICATION INSTRUCTIONS
		ML per 100 Plants	ML per 100 Plants	
Adelgids	2	1.6 mL	2.5 mL	<p>¹ Fungus gnat larvae: Control in the soil by drench or incorporation. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE will not control adult Fungus Gnats.</p> <p>² Root Mealybug: To obtain control, thoroughly drench the containerized media but do not allow leaching from the bottom of the container. Use the following rate of 1.7 fl oz (50 mL) in 150 gallons of water.</p> <p>³ Citrus Root Weevil: For use on non-bearing citrus nursery stock.</p> <p>⁴ Thrips: For suppression on foliage only. Thrips in buds and flowers will not be suppressed.</p> <p>Foliar insect control is accomplished by the uptake of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE from a healthy root system. This allows the active ingredient to move up into the plant.</p> <p>RESTRICTIONS</p> <ul style="list-style-type: none"> • For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato. • Do not apply more than 1.6 pt (0.4 lb of active ingredient) /A per year.
Aphids	2	1.6 mL	2.5 mL	
Armored scales	3	2.5 mL	3.7 mL	
Fungus Gnats (larvae only) ¹	3	2.5 mL	3.7 mL	
Japanese beetles (adults)	4	3.3 mL	5 mL	
Lacebugs	4	3.3 mL	5 mL	
Leaf beetles (including elm and viburnum leaf beetles)	5	4.2 mL	6.3 mL	
Leafhoppers (including glassy-winged sharpshooter)	6	5 mL	7.7 mL	
Leafminers	7	5.9 mL	9.1 mL	
Mealybugs	7	5.9 mL	9.1 mL	
Psyllids	8	6.6 mL	10 mL	
Root mealybugs ²	8	6.6 mL	10 mL	
Root Weevil Complex (such as Apopka Weevil, Black Vine Weevil, Citrus Weevil ³)	9	7.4 mL	11.1 mL	
Soft scales	9	7.4 mL	11.1 mL	
Thrips (suppression) ⁴	10	8.3 mL	12.5 mL	
Whiteflies	11	9 mL	14.3 mL	
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)	12	10 mL	16.7 mL	

DRENCH AND IRRIGATION APPLICATIONS

For use only on greenhouse and nursery ornamentals, vegetable plants*, and interiorscape plants using soil drenches, micro irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held or motorized calibrated irrigation equipment.

PEST	POT SIZE (inches)	Herbaceous species including vegetable plants* (1 or 2 plants per pot)	Woody perennials, Herbaceous species including vegetable plants* (3 or more plants per pot)	APPLICATION INSTRUCTIONS
		No. pots treated with 1.7 fl. oz. (50 mL)	No. pots treated with 1.7 fl. oz. (50 mL)	
Adelgids	2	3,000	2,000	Thoroughly wet most of the potting medium but do not allow runoff or leaching from the bottom of the container.
Aphids				
Fungus Gnats (larvae only) ¹	3	2,000	1,350	Follow the application with moderate irrigation. During the next 10 days, carefully irrigate to avoid the loss of the active ingredient due to leaching. ¹ Fungus gnat larvae: Control in the soil by drench or incorporation. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE will not control adult Fungus Gnats. ² Root Mealybug: To obtain control, thoroughly drench the containerized media but do not allow leaching from the bottom of the container. Use the following rate of 1.7 fl oz (50 mL) in 150 gallons of water. ³ Citrus Root Weevil: For use on non-bearing citrus nursery stock. ⁴ Thrips: For suppression on foliage only. Thrips in buds and flowers will not be suppressed. Foliar insect control is accomplished by the uptake of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE from a healthy root system. This allows the active ingredient to move up into the plant.
Japanese beetles (adults)	4	1,500	1,000	
Lacebugs				
Leaf beetles (including elm and viburnum leaf beetles)	5	1,200	800	
	6	1,000	650	
Leafhoppers (including glassy-winged sharpshooter)	7	850	550	
Leafminers	8	750	500	
Mealybugs				
Psyllids	9	675	450	
Root mealybugs ²				
Root Weevil Complex (such as Apopka Weevil, Black Vine Weevil, Citrus Weevil ³)	10	600	400	
	11	550	350	
Soft scales				
Thrips (suppression) ⁴	12	500	300	
Whiteflies				
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafers, Oriental Beetle, Asiatic Garden Beetle)				

(continued)

DRENCH AND IRRIGATION APPLICATIONS *(Continued)*

For use only on greenhouse and nursery ornamentals, vegetable plants*, and interiorscape plants using soil drenches, micro irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held or motorized calibrated irrigation equipment. *(continued)*

PEST	POT SIZE (inches)	Herbaceous species including vegetable plants* (1 or 2 plants per pot)	Woody perennials, Herbaceous species including vegetable plants* (3 or more plants per pot)	APPLICATION INSTRUCTIONS
		No. pots treated with 1.7 fl. oz. (50 mL)	No. pots treated with 1.7 fl. oz. (50 mL)	
	Ornamental and vegetable plants* grown in flats, benches, or beds		1.7 fl. oz. (50 mL) per 3,000 sq. ft.	<p>Mix the appropriate amount of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE in sufficient water to evenly cover the treatment area.</p> <p>RESTRICTION: Do not use less than 2 gallons of mixture per 1000 sq. ft.</p> <p>Apply as a broadcast treatment. Before planting, mix into the potting medium or apply after to established plants. Lightly irrigate after application to established plants for best control.</p> <p>RESTRICTION: Do not allow leaching or runoff for 10 days after application.</p>
<p>* For use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.</p>				

(continued)

DRENCH AND IRRIGATION APPLICATIONS (Continued)

PEST	Containerized Plants		APPLICATION INSTRUCTIONS
	Container Size	No. pots treated with 1.7 fl. oz. (50 mL)	
Adelgids	1 gallon	340 to 244	<p>Apply in sufficient water to wet the potting medium. For best control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.</p> <p>To prevent leaching, use 1.7 fl. oz. (50 mL) of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE in the appropriate amount of water to treat the number of pots based on the pot size as stated in the table.</p> <p>Foliar insect control is accomplished by the uptake of QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE from a healthy root system. This allows the active ingredient to move up into the plant.</p> <p>¹ Fungus gnat larvae: Control in the soil by drench or incorporation. QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE will not control adult Fungus Gnats.</p> <p>² Root Mealybug: To obtain control, thoroughly drench the containerized media but do not allow leaching from the bottom of the container. Use the following rate of 1.7 fl oz (50 mL) in 150 gallons of water.</p> <p>³ Citrus Root Weevil: For use on non-bearing citrus nursery stock.</p> <p>⁴ Thrips: For suppression on foliage only. Thrips in buds and flowers will not be suppressed.</p>
Aphids			
Fungus Gnats (larvae only) ¹	2 gallon	280 to 210	
Japanese beetles (adults)	3 gallon	220 to 165	
Lacebugs			
Leaf beetles (including elm and viburnum leaf beetles)	5 gallon	160 to 110	
	7 gallon	100 to 75	
Leafhoppers (including glassy-winged sharpshooter)	10 gallon	60 to 45	
Leafminers	15 gallon	40 to 30	
Mealybugs			
Psyllids	20 gallon	20 to 15	
Root mealybugs ²			
Root Weevil Complex (such as Apopka Weevil, Black Vine Weevil, Citrus Weevil ³)			
Soft scales			
Thrips (suppression) ⁴			
Whiteflies			
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)			

(continued)

PEST	Containerized Plants		APPLICATION INSTRUCTIONS
	Container Size	No. pots treated with 1.7 fl. oz. (50 mL)	
Field and Forest Nurseries			
White grub larvae (such as Japanese Beetle, Masked Chafers, European Chafer, Oriental Beetle, Asiatic Garden Beetle)	1.7 fl. oz. (50 mL) per 1,000 ft. of row or 3,000 sq. ft.		Before application, mow the vegetation in the treatment area to a height of 3 inches or less. Mow to the lowest height possible. Applications must be made May through July. Treatment must be followed by rainfall or irrigation. Do not use less than 2 gallons of spray volume per 1000 square feet. Apply as a uniform band on either side of the row using a band width six (6) inches wider than the actual root ball diameter to be dug. Do not overlap bands in adjacent rows. For grub control in areas of turf, apply as a broadcast application using 1.35 to 1.7 fl oz (40 to 50 mL) per 3000 sq. ft.

RESTRICTIONS

Do not graze treated areas or use clippings for treated areas for feed or forage. Do not allow runoff or puddling of irrigation water following application.

Do not apply QUALI-PRO IMIDACLOPRID 2F TURF & ORNAMENTAL INSECTICIDE to soils which are water logged or saturated, which will not allow penetration into the root zone of the plants.

Do not allow leachate run out for the first 10 days after application, in order to retain the product and facilitate full plant uptake of the active ingredient.

For outdoor ornamentals, do not apply more than a total of 1.6 pt (0.4 lb of active ingredient imidacloprid) /A per year.

Food Crops: Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval must be observed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children, preferably in a locked storage area. Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Open dumping is prohibited. Pesticide wastes are toxic. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid, Nonrefillable containers small enough to shake (i.e. with capacities equal to less than five gallons). Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Once container is rinsed, offer for recycling if available, or puncture and dispose of in a sanitary landfill.

Rigid, Nonrefillable containers that are too large to shake (i.e. with capacities greater than 5 gallons or 50 lbs).

Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

(continued)

STORAGE AND DISPOSAL *(continued)*

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Once container is rinsed, offer for recycling if available, or puncture and dispose of in a sanitary landfill.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.**

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc. election, the replacement of product.

QUALI-PRO[®]

IMIDACLOPRID 2F

Turf & Ornamental Insecticide



Turf and Ornamental and Nursery and Greenhouse Uses

- Foliar and Systemic Insect Control for use on Turfgrass including Sod Farms, Landscape Ornamentals, Fruit and Nut Trees and Interior Plantscapes
- Foliar and Systemic Insect Control for use in Greenhouses, and Nurseries on Ornamentals, Fruit and Nut Trees and Vegetable Plants

ACTIVE INGREDIENT:	% BY WT.
Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	21.8%
OTHER INGREDIENTS:	78.2%
TOTAL:	100.0%

Contains 2 pounds of imidacloprid per gallon.
Flowable insecticide.
Shake well before using.

CONTENTS:
1 GALLON

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

EPA Reg. No. 53883-514
EPA Est No. 53883-TX-002

Manufactured for:
Control Solutions, Inc.
5903 Genoa Red Bluff Road
Pasadena, TX 77507

EPA 120313/Rev A