

RESTRICTED USE PESTICIDE
Toxic to fish and aquatic organisms.
For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

Product Bulletin



EPA Reg. No. 34704-858

SUPPLEMENTAL LABELING

For use on Soybeans, Fruiting Vegetables, Peanuts, Root Crops and Mayhaws.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions in the registered product label.

DIRECTIONS FOR USE

SOYBEANS

CROP	PEST	DOSAGE		REMARKS
		LBS AI/A	FL OZ/A	
Soybean	Soybean aphid	0.033-0.10	2.1-6.4	Apply foliar treatments in at least 10 gallons per acre at the rate of 6.4 fl. oz. (0.1 lb. active) per acre at a minimum of 30 days intervals. Do not apply more than 0.31b ai per acre per season. Do not apply within 18 days of harvest.
	Spider mites	0.08-0.10	5.12-6.4	

FRUITING VEGETABLES

CROP	PEST	DOSAGE		REMARKS
		LBS AI/A	FL OZ/A	
Eggplant, Pepper (bell & non-bell), Groundcherry, Pepino.	Armyworms including Beet armyworm, Fall armyworm, Southern yellowstriped armyworm, Cabbage looper, Colorado potato beetle, Corn earworm, Cucumber beetle, Cutworms, European corn borer, Flea beetle, Leafminers, Loopers, Pepper weevil, Plant bug, Stink bug, Thrips, Tomato Hornworm, Tomato pinworm, Vegetable leafminer, Whitefly	0.033-0.10	2.1-6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not make applications less than 7 days apart. Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season. Do not apply within 7 days of harvest.
	Banks grass mite, Broad mite, Carmine mite, <i>Lygus</i> spp., Pacific spider mite, Twospotted spider mite,	0.08-0.10	5.12-6.4	

ROOT CROPS

CROP	PEST	DOSAGE		REMARKS
		LBS AI/A	FL OZ/A	
Burdock (edible), Carrot, Celeriac, Chervil, Turnip (rooted), Chicory, Ginseng, Horseradish, Parsley, Parsnip, Radish, Radish (oriental), Rutabaga, Salsify, Salsify (black), Salsify (Spanish), Skirret, Turnip	Aphids, Beet armyworm, Celery leaf tier, Corn earworm, Cross-striped Cabbageworm, Cutworms, Diamondback moth, European corn borer, Fall armyworm, Fire ants, Flea beetles, Green cloverworm, Hornworms, Imported cabbageworm, Loopers, Southern armyworm, Spider mites, Tobacco budworm, Velvetbean caterpillar, Whitefly, Yellowstriped armyworm	0.08-0.10	5.12-6.4	Apply foliar treatments in at least 25 gallons water per acre. Apply no more than once every 7 days. Do not apply more than 0.5 lb. ai per acre per season. Do not apply within 21 days of harvest.
Garden beet	Aphids, Fire ants, Flea Beetles, Lepidopterous larvae, Spider mites, Whitefly	0.08-0.10	5.12-6.4	Apply foliar treatments in at least 25 gallons water per acre. Apply no more than once every 7 days. Do not apply more than 0.4 lb. ai per acre per season. Do not apply within 1 day of harvest.

PEANUT

CROP	PEST	DOSAGE		REMARKS
		LBS AI/A	FL OZ/A	
Peanut	Beet armyworm, Corn earworm, Cutworm spp., Fall armyworm, Grasshoppers, Green cloverworm, Leafhoppers, Lesser cornstalk borer, Loopers, Rednecked peanut worm, Southern armyworm, Southern corn rootworm, Stink bugs, Threecornered alfalfa hopper, Velvetbean caterpillar, Yellowstriped armyworm	0.033-0.10	2.1-6.4	Apply foliar treatments in at least 10 gallons per acre at the rate of 6.4 fl. oz./A (0.1 lb. active) per acre at a minimum of 30 days intervals. Do not apply more than 0.31b ai per acre per season. Do not apply within 18 days of harvest.
	Aphids, Spider mites, Thrips, Whitefly	0.08-0.10	5.12-6.4	

MAYHAW

CROP	PEST	DOSAGE		REMARKS
		LBS AI/A	FL OZ/A	
Mayhaw	Plum curculio	0.08-0.10	5.12-6.4	Apply foliar treatments in at least 28 gallons per acre. Apply no more than once every 7 days. Do not apply more than 0.2 lb. ai per acre per season. Do not apply within 30 days of harvest.

