SPECIMEN LABEL

Rev 04/22/2022



FOR INDOOR AND OUTDOOR USE -

- Modifies the exposure profile of the insects enhancing the bioavailability of the active ingredient(s) in insecticides
- Provides an even deposition of the pesticide active ingredient on the treated surface/insect
- Provides increased effectiveness of insecticide sprays
- · Acts to improve spray coverage and enhance the pest's exposure to the insecticide used in the tank mix
- · Disperses spray droplets for more even coverage of insect and surfaces
- · Enhances deposition and uniform coverage of herbicides, fungicides, insecticides, fertilizers, and plant growth regulators

Stik-Kote* can be applied on agricultural, turf and ornamental, as well as indoor and outdoor non-cropland sites. Not for aquatic use.



Principal Functioning Agents:

ingredients are exempt from the requirements of a lerance under 40 CFR 180.

NET CONTENTS: 32 FL. OZ

STOP - READ THE LABEL BEFORE USE KEEP OUT OF REACH OF CHILDREN Warning/Aviso

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY
STATEMENTS, SEE INSIDE BOOKLET FOR DIRECTIONS FOR USE. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)



— FOR INDOOR AND OUTDOOR USE

GENERAL INFORMATION

Stik-Kote" is a nonionic organosilicone adjuvant useful in the application of herbicides, fungicides, insecticides, fertilizers, and plant growth regulators. It lowers the surface tension of spray solutions.

Lower surface tension will result in more effective wetting and more uniform coverage on foliar surfaces, resulting in a reduction in water requirements.

Stik-Kote* is formulated to modify the exposure profile of insects to enhance the bioavailability of the active ingredient(s) in insecticides. It also works to disperse the spray droplets and provide even deposition of the pesticide active ingredient(s) on the insects.

Stik-Kote" acts to improve spray coverage and enhance the pest's exposure to the insecticide used in the tank mix. It can be applied to a wide variety of agricultural crops and ornamental plants as well as indoor and outdoor non-cropland sites. Not for aquatic use.

COMPATIBILITY

Stik-Kote" is compatible with most herbicides, fungicides, insecticides, fertilizers, and plant growth regulators that are not highly alkaline. If separation of spray mixture occurs, do not use.

DIRECTIONS FOR USE READ ENTIRE LABEL PRIOR TO USE.

Stik-Kote* will give best results in low volume spray applications, which result in around 50% of the normal volume of water when sprayed to run-off. High water volumes with Stik-Kote* will result in excessive run-off of spray liquid and may lead to poor efficiency.

Adjust either the rates of Stik-Kote" or the water volumes to obtain thorough coverage without excessive run-off.

Buffer spray water to pH 5.0-7.5 if necessary.

- Fill spray tank half full with water and begin agitation.
 If foaming is expected, add silicone-based antifoam to the tank before adding the other products.
- Add pesticides and/or nutrients as directed on their respective labels; maintain agitation. Continue filling with water until the tank is -90% full
- Add the required amount of Stik-Kote® and mix thoroughly
- Add the remaining water to 100% full.

It is the user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture

MANUFACTURED BY: ICB Pharma ul. Mozdzierzowcow 6a 43-602 Jaworzno, Poland

DISTRIBUTED BY: 3D BioSciences, LLC. 2101 Mystic Ridge Rd Knoxville, TN 37922

CA Reg. No.: 92035-50001-AA



USE RATES

Stik-Kote" is recommended to be used with herbicides, fungicides, insecticides, fertilizers, and plant growth regulators registered for the following uses

INDOOR & OUTDOOR TREATMENT OF NONCROPLAND SITES

Adjuvant Concentration: 1.0% - 2.0% (1 1/4 - 2 1/2 fl. oz of the product per gallon).

Care should be taken while using Stik-Kote" mixture on surfaces susceptible to damage/stains (e.g., mattresses, carpet, furniture). Treat a small area to ensure damage/stain does not occur before full application

INDOOR & OUTDOOR USE ON CROPS (INCLUDING, BUT NOT LIMITED TO: VEGETABLES, FRUIT TREES, BERRIES CEREAL GRAINS, AND HERBS) AND ORNAMENTAL PLANTS (INCLUDING, BUT NOT LIMITED TO: TREES, SHRUBS, FLOWERS, FOLIAGE PLANTS AND TURF)

Adjuvant Concentration: Due to varying phytotoxic risk of the Stik-Kote[®] on different plants, the adjuvant concentration must be adjusted according to plant being treated. Please refer to the table for the concentrations to be used on the following plants.

Agricultural Crop Application Rates:

Agricultural Crop Application Rates:

Mix as directed using sufficient volumes to ensure thorough foliage coverage.

Fruit trees, berries, and vegetables: 0.10% - 0.15% v/v (12 ¾ -19 ¼ fl. oz of the product per 100 gallons).

Cucumbiers: 0.10% v/v (12 ½ fl. oz of the product per 100 gallons).

Tomatoes, peppers, eggplants: 0.05% - 0.07% v/v (6 ½ - 9 fl. oz of the product per 100 gallons).

Cereal grains: 0.15% v/v (19 ¼ fl. oz of the product per 100 gallons). Herbs: 0.05% v/v (6 1/2 fl. oz the product per 100 gallons)

Ornamental Application Rates:

Mix as directed using sufficient volumes to ensure thorough foliage coverage Omamental trees and shrubs: 0.10% - 0.15% $_{\rm V}$ /v (12 % - 19 $_{\rm V}$ fi n. oz of the product per 100 gallons) Roses: 0.08% - 0.10% $_{\rm V}$ /v (10 $_{\rm V}$ 4 - 12 $_{\rm V}$ fi, oz of the product per 100 gallons) Other omamental plants: 0.04% - 0.05% $_{\rm V}$ /v (5 $_{\rm V}$ 4 - 6 $_{\rm V}$ fi, oz, of the product per 100 gallons)

For greenhouse grown ornamental, flowering and foliage plants: Do not apply more frequently than once every 5 days.

NEVER EXCEED LABEL RATES DUE TO PHYTOTOXICITY RISK.

IMPORTANT NOTE:

Stik-Kote* has demonstrated good plant safety when applied at the specified dilution rates. However, not all plants and spray mixtures have been tested.

As it is not possible to evaluate the phytotoxicity of Stik-Kote* towards all plant varieties that may react differently to spray applications in different growth stages or under varying environmental conditions.

Before making widespread applications of Stik-Kote", the user or applicator advisor must have experience with the tank mix or have conducted a phytotoxicity trial by treating a limited number of plants and observing for phytotoxicity over a

For sensitive crops or plant stages (e.g., seedlings, tomatoes, peppers, herbs, cucumber), avoid spraying when drying times may be significantly prolonged beyond average drying times.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container at 32-86°F. Keep away from open flame, sparks, heated surfaces, and avoid heat (>86°F). Store in a cool, dry place inaccessible to children and pets.

CONTAINER DISPOSAL: DO NOT puncture or incinerate! Empty container by using the product according to the label directions. Non-refillable container. If empty: Offer container for recycling, if available, or place in trash if allowed by state and local regulations. If partly filled: Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but remporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

| FIRST AID | |
|-----------|--|
|-----------|--|

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- · Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information on this product, call the poison control center at 1-800-222-1222.

WARRANTY

Read the entire Directions for Use and Warranty before using this product. By using this product, user or buyer accepts head on the strict Directions for use and warranty determine using this product, by using product, part or object accepts the following warranty. The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer. The seller is committed to providing high quality products. To the extent consistent with applicable law, the seller makes no warranties, express or implied of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, the manufacturer or the seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.

In Case of Emergency, call: 1-800-222-1222 Non-Emergency call: 1-800-858-7378

Distributed by: 3D BioSciences, LLC. 2101 Mystic Ridge Rd Knoxville, TN 37922

This specimen label is intended for use only as a guide in providing general information regarding the directions, warning and cautions associated with the use of this product. As with any pesticide, always follow the label instructions on the package before using.