DIRECTIONS FOR USE

General Use: Use 1 to 15 gallons per acre. Shake well before use.

MIXING INSTRUCTIONS

- 1. Add BIOAG TM-7 L to water and start agitation
- 2. Add pesticides and other nutrients.
- 3. Fill the tank and agitate thoroughly

For further details on direction for use and application rates, please visit www.bioag.com or consult with a local crop advisor or agronomist.

Appx 30 ml = 1 fl. oz. 1:100 dilution = 0.5 fl. oz/gal

TM-7 is a registered trademark of Faust-Bio-Agricultural Services, Inc.

WARNING

KEEP OUT OF REACH OF CHILDREN

CAUTION NOTICE: Buyer assumes all risk for use not in accordance with labeled directions. Harmful if swallowed. Avoid contact with skin, eyes, open cuts, and sores. Wash hands after use.



sales@bioag.com • (855) 844-4632





Distributed and Guaranteed by: Faust Bio-Agricultural Services, Inc. 6080 Wigrich Road, Independence, OR 97351









ØBIOAG

HUMIC ACIDS WITH MICRONUTRIENTS N-N-6

CHADANTEED AMAINCIC.

OMRI

For Organic Use

GUARANTEED ANALYSIS:			
Soluble Potash (K ₂ 0)	6%	Iron (Fe)	0.4%
Boron (B)	0.65%	Manganese (Mn)	0.1%
Cobalt (Co)	. 0.03%	Molybdenum (Mo)	0.08%
Copper (Cu)	0.1%	Zinc (Zn)	0.35%
DERIVED FROM: Boric Acid	, Cobalt Sulfate, Cop	per Sulfate, Manganese S	Sulfate,
Sodium Molyhdate Ferrous	Sulfate 7inc Sulfate	and Potassium Hydroxi	de

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

DERIVED FROM: Leonardite. PURPOSE: May increase micronutrient uptake.

NET CONTENTS: 1 GAL (3.8L) 10.08LB Information regarding the contents and levels of metals in this product is available on the internet at: http://www.aapfco.org/metals.html

WARNING: Contains boron. Do not use boron sensitive crops. Use only according to the manufacturer's directions. The use of this material on any crops other than those recommended may result in injury to the crop(s). Contains molybdenum. Forage crops high in molybdenum may be toxic to grazing ruminant animals or other livestock. Use only according to the manufacturer's directions.

GENERAL INFORMATION

BIOAG TM-7 L is generally compatible with most pesticide, herbicide, fungicide, and fertilizer solutions. Add to water before adding pesticides. Test for compatibility by means of a jar test prior to large scale tank mixing. BIOAG TM-7 L is intended for use as a plant nutrient supplement for the purpose of enhancing micronutrient uptake in plants and soil. BIOAG TM-7 L is intended for use on all agricultural, horticultural, and ornamental crops that require and respond to humic acid and micronutrient supplements. BIOAG TM-7 L is to be applied as a foliar application through conventional, low-volume, or aerial sprayers. BIOAG TM-7 L may be formulated into NPK fertilizers. This product should be stored in a cool, dry area out of direct sunlight. For proper agronomic application rates for your crop, soil type, and climate, please consult with a local crop advisor, farm advisor, or agronomist.

FIRST AID IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. Call a poison control center or doctor for treatment advice. IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by

mouth to an unconscious person. IF INHALED: Move the person to fresh air. If a person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

