



Product Bulletin

RESTRICTED USE PESTICIDE

Due to toxicity to Fish and Aquatic Organisms. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

TOMBSTONE™ Emulsifiable Pyrethroid Insecticide
EPA Reg. No. 34704-912
For use on Soybeans to Control Bean Leaf Beetles

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF CONNECTICUT, DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MAINE, MARYLAND, MASSACHUSETTS, MINNESOTA, MISSOURI, NEBRASKA, NEW JERSEY, NORTH DAKOTA, OHIO, PENNSYLVANIA, RHODE ISLAND, SOUTH DAKOTA, VERMONT, VIRGINIA, WEST VIRGINIA AND WISCONSIN.

FIFRA Section 2(ee) Recommendation: This recommendation is made and permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U. S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

RECOMMENDED APPLICATION		
Use Site	Insect	Rate of TOMBSTONE™
Soybeans	Bean Leaf Beetles	1.0 – 1.6 fl. oz/A

Apply specified dosage per acre as a foliar spray when bean leaf beetles begin to migrate into soybean fields. **TOMBSTONE™** should be applied when soybeans are between growth stage VC-V2. To control infestations occurring in soybean past the V2 growth stage or in areas where Bean Pod Mottle Virus transmission is a concern, refer to application rate information contained on the **TOMBSTONE™** label. A maximum of 2.8 fluid ounces/acre may be applied within a 7 day period. Total applications may not exceed 11.2 fluid ounces of **TOMBSTONE™** per season.