

ULTIMATUM Pest

FIPRONIL

ROACH GEL BAIT



FIRST AID

IF ON SKIN OR CLOTHING

- Take off contaminated clothing
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00 AM to 12:00 PM Pacific Standard Time. For medical emergencies, call the poison control center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read entire label. Use strictly in accordance with label precautionary statements and directions.

INDOOR & OUTDOOR USE

Use **ULTIMATUM PEST ROACH GEL BAIT** as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches in residential areas and the nonfood/non-feed areas of institutional, warehousing and commercial establishments, including warehouses,

restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, apartment buildings, bakeries, beverage plants, breweries, bottling facilities, candy plants, canneries, cereal processing and manufacturing plants, dairies and dairy product processing plants, frozen food plants, schools, laboratories, computer facilities, sewers, aircraft, buses, boats/ships, trains, homes, houses, industrial buildings, office buildings, kennels, kitchens, manufacturing facilities, spice plants, stores, wineries and similar structures, mausoleums, meat processing and packaging plants, meat and vegetable canneries, pet shops, and zoos. Non-food/non-feed areas include bedrooms, living rooms, bathrooms, closets, media rooms, garbage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

[in the non-food/non-feed areas of institutional, warehousing and commercial establishments, including warehouses, restaurants (non-public areas), food processing plants (including meat or vegetable processing plants or canneries), supermarkets (non-public areas), hospitals (non-public areas), laboratories, computer facilities, sewers, animal housing facilities (such as but not limited to poultry houses, cattle, sheep, swine, goat and dairy facilities, horse stables, dog kennels). Non-food/non-feed areas are areas such as garbage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).]

Product may stain textiles, fabrics, and or porous surfaces.

USE RESTRICTIONS

USE ONLY IN AREAS NOT EASILY ACCESSIBLE TO CHILDREN AND PETS.

- Do not treat food preparation surfaces. Keep gel away from foods and food contact surfaces.
- Do not apply bait gel to areas which have been recently sprayed with insecticide, and do not spray insecticide over gel as it may cause the bait to become repellent.
- Place bait out of reach of pets and zoo animals.

ULTIMATUM Pest

ROACH GEL BAIT

FIPRONIL



USE **ULTIMATUM PEST ROACH GEL BAIT** as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches in residential areas and the nonfood/non-feed areas of institutional, warehousing and commercial establishments.

This product may also be used in food/feed areas of food/feed handling establishments to crack and crevice treatment only.

Pest population reduction will be visible within 1 day for German roaches and 3-4 days for American roaches.



Manufactured for & Distributed by:

MYLVA S.A.

Sant Galderic 23 -08395 Sant Pol de Mar (Barcelona) Spain

Tel: + 34 93 4153226

www.mylva.com

NET WT:

4 cartridges of 30 g / 0,71 oz

ACTIVE INGREDIENTS:

ACTIVE INGREDIENT: Fipronil (CAS NO. 120068-37-3) 0.05%

OTHER INGREDIENTS:

TOTAL: 99.95%

TOTAL: 100.00%

EPA Est. No. 92418-ESP-1

EPA Reg. No. 92028-4-92418 Batch No. Batch and expiry date printed on the cartridge.

KEEP OUT
OF REACH
OF CHILDREN
CAUTION



8 437026 137007

ULTIMATUM Pest

ROACH GEL BAIT

FIPRONIL



APPLICATION DIRECTIONS

Application rate for all uses: Application rate varies according to the level of infestation and species to be controlled. Apply gel as round spots or thin beads into cracks and crevices or as a spot treatment. Place bait close to cockroach harborages, foraging, and feeding areas such as corners and cracks and crevices. Numerous smaller bait placements provide faster control than fewer larger placements. Use the highest rate of application in cases of severe infestation and areas with excessive clutter or alternative food sources that cannot be eliminated.

German Cockroaches, Brown Banded Cockroaches, and stored product pests: For use in maintenance programs where regular monthly applications are made for continuous control of cockroach populations, use 0.75 grams of bait per square yard. This rate will not provide fast control of existing infestations. For light to moderate infestations (or for retreatments), use up to 1.5 grams of bait per square yard. To clean out severe or heavy infestations, use up to 3.0 grams of bait per square yard.

Large Cockroaches such as American, Smokey-Brown or Oriental: 1 to 2 grams of bait per square yard.

BAIT QUANTITY REFERENCE TABLE	
Approximate weight per spot diameter	Approximate weight per bead length
0.1 gm = 1/4 inch diameter spot	1/3 gram = 1.5-inch bead
1.0 gm = Dime size spot	2/3 gram = 3.0-inch bead
1.5 gm = Quarter size spot	1 1/3 gram = 6-inch bead

Place bait under baseboards, sinks, countertops, seats, around water pipes, water heaters, dishwasher equipment, in crack and crevices as well as voids where insects harbor.

PANTRY PESTS: To control exposed, adult stages of beetles: cadelle, cigarette beetle, confused flour beetle, dermestid, drugstore beetle, flat grain beetle, granary weevil, lesser grain borer, lesser mealworm, mealworm, merchant grain beetle, red flour beetle, rice weevil, rusty grain beetle, sawtoothed grain beetle, spider beetle, or warehouse beetle.

Applications of this product will kill adult stages of beetles remaining in area after removing infested foodstuffs and cleaning up spilled foodstuffs and dust. Apply in voids, cracks and crevices, behind and under cupboards, and other difficult to reach areas where insects may hide. Application rate is dependent on level of infestation.

APPLICATIONS IN FOOD/FEED HANDLING ESTABLISHMENTS: Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. If gel contacts an exposed surface where food is handled, remove gel and wash exposed surface with an effective cleaning compound followed by a potable water rinse prior to use. Apply with bait injector directly into cracks and crevices. Place applicator tip 1/2 inch into cracks, crevices, holes and other small openings where roaches may be a problem. Apply small amounts of gel.

USE RESTRICTIONS IN FOOD/FEED HANDLING ESTABLISHMENTS:

- Do not apply bait to areas where food/feed, utensils, or processing surfaces may become contaminated.
- Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation, and prepared food holding surfaces as bait may be removed by washing.
- Do not apply bait to surfaces where the temperature exceeds 130°F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, fume hoods, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers, and hot water pipes.
- Do not deposit gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves, and other equipment where cockroaches hide. During follow-up visits inspect bait placements and reapply when necessary.

Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted.

Gel placements should be at or near harborages or aggregation areas, such as corners or cracks and crevices. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control.

ULTIMATUM Pest

ROACH GEL BAIT

FIPRONIL



For outdoor use (adjacent to home -or- structure): Place bait in areas adjacent to structures where roaches may nest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints. Apply bait in protected areas whenever possible. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness. Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushing.

Refillable stations: NOUVEL SALES Fipronil Roach Bait Gel may be placed into refillable bait stations designed for interior or exterior use; when placing bait in stations do not exceed 1 gm (1/5 teaspoon) and do not exceed the capacity of the stations. Close the station securely after application. During follow up visits inspect placements and reapply as needed.

DIRECTIONS FOR LOADING CARTRIDGES AND OPERATION:

1. With the teeth of the drive rod facing up, pull the drive rod all the way back. If the drive rod is initially hard to move, simply un-screw (turn counter-clockwise) the [black] [or other color] portion of the applicator that holds the gel cartridge until the drive rod moves easily. **2.** Attach a pre-filled cartridge of gel with a twist-locking motion. Remove the outlet cap and attach a dispensing tip. After use, the cartridge may be resealed with the enclosed [black][or other color] outlet cap. The [black] [or other color] outlet cap is reusable. **3.** Turn the knob of the drive rod so that the teeth are facing down and move it slightly forward so that the trigger mechanism will engage. Squeeze the trigger until the drive plunger sits against the piston in the gel cartridge. Gently squeeze the trigger until the gel flows out. **4.** If gel continues to flow after the trigger is released, simply un-screw the portion of the bait applicator that holds the cartridge to a point where gel does not continue to flow after the trigger is released.

DIRECTIONS FOR CHANGING CARTRIDGES: With the arrow on the drive rod pointing up, pull the drive rod out as far as it will go. Remove the empty cartridge and follow instructions in the Storage and Disposal box. Cartridges that still contain gel can be sealed with the end cap and outlet cap for future use. To attach a new cartridge, follow the above directions.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CAUTION: Keep gel away from foods and food contact surfaces. Causes moderate eye irritation. Avoid contact with eyes and clothing. Wash thoroughly with soap and water after handling.

CONTAINER HANDLING:

If empty: Do not reuse container. Place in trash or offer for recycling if available.

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of MYLVA S.A.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold MYLVA S.A. harmless for any claims relating to such factors. MYLVA S.A. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of MYLVA S.A., and to the extent consistent with applicable law, Buyer and User assume the risk of any such use. MYLVA S.A. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

IT IS MYLVA S.A. INTENTION THAT to the extent consistent with applicable law, in no event shall MYLVA S.A. be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF MYLVA S.A. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCTOR, AT THE ELECTION OF MYLVA S.A., THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law, MYLVA S.A. offers this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except

by written agreement signed by a duly authorized representative of MYLVA S.A..